




MEAD/McCULLOUGH- VICTORVILLE/ADELANTO

TRANSMISSION PROJECT

TECHNICAL REPORT
VOLUME II
NATURAL ENVIRONMENT

MEAD/McCULLOUGH-VICTORVILLE/ADELANTO
TRANSMISSION PROJECT
TECHNICAL REPORT

VOLUME II
NATURAL ENVIRONMENT

Prepared by:

Dames & Moore

Prepared for:

Los Angeles Department
of Water and Power
U.S. Department of the Interior
Bureau of Land Management

June 1985

TABLE OF CONTENTS

	PAGE
CHAPTER 1 - METEOROLOGY AND AIR QUALITY	1- 1
Introduction	2- 1
Secondary Hazards	2- 1
Meteorology	2- 1
Air Quality	2- 3
Impact Assessment/Mitigation Planning	2- 7
Effects of Construction	2- 7
Effects of Operation	2- 7
References	2- 7

CHAPTER 2 METEOROLOGY/AIR QUALITY

TABLE OF CONTENTS

	PAGE
CHAPTER 2 - METEOROLOGY AND AIR QUALITY	2- 1
Introduction	2- 1
Inventory Results	2- 1
Meteorology	2- 1
Air Quality	2- 3
Impact Assessment/Mitigation Planning	2- 7
Effects of Construction	2- 7
Effects of Operation	2- 7
References	2- 9

LIST OF TABLES

TABLE NUMBER	TITLE	FOLLOWS PAGE
2-1	Wind Frequency Distribution	2- 9
2-2	Percentage Frequency of Wind Direction and Speed	2- 9
2-3	Wind Direction Frequency Distribution	2- 9
2-4	Mean Temperature	2- 9
2-5	Precipitation	2- 9
2-6	Ambient Air Quality Standards	2- 9
2-7	Oxidant Concentrations and Exceedances	2- 9
2-8	Carbon Monoxide Concentrations and Exceedances	2- 9
2-9	Maximum Measured Nitrogen Dioxide Concen- trations in Project Area	2- 9
2-10	Particulate Concentrations and Exceedances	2- 9

METEOROLOGY AND AIR QUALITY

INTRODUCTION

This report describes the meteorology and air quality of the study area for the proposed Mead/McCullough-Victorville/Adelanto Transmission Project. The study was conducted by Ms. Cynthia Folks, Assistant Engineer for Dames & Moore. Ms. Folks holds a B.S. degree in Environmental Engineering, and works in the Environmental and Permitting Group of Dames & Moore's Santa Barbara, California office.

INVENTORY RESULTS

Meteorology

Regional Climate

The project area lies within the southwest portion of Nevada, across the Nevada/California state border and through the Mojave Desert parallel to Interstate 15. The meteorology of the region is typical of the Mojave Desert (High Desert) region of Nevada and California with hot dry summers and cool dry winters. Large daily and seasonal temperature differences are in evidence due primarily to the coastal mountain ranges which inhibit the penetration of marine air into the High Desert.

Precipitation is sparse and infrequent. The semi-permanent Pacific High Pressure Cell, which resides over the west coast during much of the year, keeps most storm systems north of the area. Extreme temperatures range from well over 100°F during summer to below freezing (32°F) during winter. Temperatures vary considerably with altitude.

Wind

Seasonal and diurnal wind regimes markedly affect the horizontal transport of air in the project area. Atmospheric stability associated with these winds determines the air dispersion of pollutants. Wind data sites at Edwards Air Force Base, Daggett, Nipton, and Cadiz were selected to represent wind data of the project area.

Table 2-1 shows the frequency distribution of wind speed and direction for the period of time between January 1968 and December 1970 from Edwards Air Force Base. The predominant wind direction for this area is west-southwest. The mean wind speed is approximately 8.8 miles per hour.

Table 2-2 shows the percentage frequency of wind direction and speed for the years 1955 through 1964 from Daggett, California. From the data, it is evident that the prevailing wind direction is from the west and west-northwest. The annual mean wind speed is approximately 11.3 miles per hour.

Table 2-3 shows the wind direction frequency distribution for the data sites of Nipton and Cadiz, California. The data collected (June 1979 to August 1979 in Nipton) shows that, during nighttime hours (0000 - 0500 PST), easterly drainage flow is dominant; otherwise, southerly winds dominate during the summer season. During the winter season, northerly winds will periodically dominate the flow regime in the Ivanpah Valley.

The Cadiz wind data collected during the summer months of July and August 1979 show that the prevailing wind direction is from the southeast with approximately 50 percent of the hours reporting winds from the east through south directions. Taking into account the terrain and other data from similar locations, it appears that the wind flow patterns in the Cadiz area would be from the northwest the majority of the time, and from the south or southeast during the summer.

Temperature and Humidity

Diurnal temperature ranges (defined as the difference between the daily maximum and minimum temperature) are large, up to 30°F to 40°F, due to the efficient heating and cooling of the land surface under generally clear skies. Temperatures frequently exceed 100°F during summer and occasionally dip below 32°F during winter.

Temperature data sites at Barstow, Daggett, Las Vegas, Yucca Grove (about 26 miles southwest of Ivanpah), and Victorville, were used to evaluate the monthly average temperatures of the project area. These monthly average temperatures for the various sites are shown in Table 2-4. From the data, it is evident that the warmest temperatures will be experienced in July and the coolest in January.

Highest relative humidities are recorded during winter nights due to the low temperatures, whereas spring and summer afternoons experience the driest conditions due to excessively hot temperatures at this time.

Precipitation

Precipitation varies with altitude and is affected by local mountain ranges. In general, annual precipitation on the desert floor ranges between 3.0 inches (Baker) and 5.4 inches (Victorville). Table 2-5 shows average monthly precipitation recorded for various representative sites: Baker, Barstow, Daggett, Las Vegas, and Victorville. Precipitation is higher in the mountains, such as the Clark Range, as is shown by changes in vegetation. Precipitation is highly variable from year to year. Much of a year's precipitation may fall in a few storms. Occasional violent thunderstorms cause much localized runoff, filling local washes and washing out local roads. During wet years the playas (dry lakes) may remain filled for months, then stay dry for several consecutive years.

Thunderstorms occur infrequently over the study area. Thunderstorms are observed an average of 15 days per year in Las Vegas. Occurrences of lightning striking the ground can vary greatly in an area but, in general, depends on elevation and latitude.

Air Quality

Effects of ambient air quality within an area depend on the characteristics of the receptors (population, crop, etc.) and the type, amount, and duration of exposure. Air quality standards specify upper limits of concentrations and durations of pollutants in the ambient air consistent with the management goal of preventing specific harmful effects. Air quality standards and background air quality information pertinent to the project area are described below.

Ambient Air Quality Standards

The U.S. Environmental Protection Agency (EPA), the California Air Resources Board (ARB), and the Nevada Department of Environmental Protection (NDEP) have established ambient air quality standards. National ambient air quality standards (NAAQS) specify acceptable pollutant concentrations which may be equaled and, in the case of short-term standards, exceeded up to once per year. California ambient air quality standards (CAAQS) specify unacceptable pollutant concentrations which should never be equaled or exceeded. Nevada ambient air quality standards specify pollutant concentrations that should not be exceeded. Table 2-6 summarizes the current Federal and state standards.

On December 9, 1982, the California ARB approved new ambient air quality standards for fine particulate matter (less than 10 μm diameter) of 50 $\mu\text{g}/\text{m}^3$ (24-hour average) and 30 $\mu\text{g}/\text{m}^3$ (annual average). These standards have been formally adopted and replace the previous standards for particulate matter of 100 $\mu\text{g}/\text{m}^3$ (24-hour averaging period) and 60 $\mu\text{g}/\text{m}^3$ (annual averaging period). The replacement of the NAAQS for particulate matter with a fine particulate matter standard was proposed on March 20, 1984 (49 FR 10408), but final action has not yet been taken.

Existing Air Quality

The project site is located within the Southeast Desert Air Basin of California and within the portion of Nevada southwest of Lake Mead to the California/Nevada border. The available California air quality data (California ARB, annual summary) supplied information for Victorville and Barstow, the nearest locations in the project area. Clark County Health Department supplied air quality data for Las Vegas and Henderson, Nevada, also located in the project area. The air contaminants of interest in characterizing the existing air quality in the project area are oxidants (O_3), carbon monoxide (CO), oxides of nitrogen (NO_x , measured as NO_2), sulfur dioxide (SO_2), total

suspended particulates (TSP), sulfates (SO_4), lead (Pb), and nitrates (NO_3). A review of available air quality data for the three-year period 1981 through 1983 is presented in the following sections.

Attainment/Nonattainment Areas

The air quality of the project area encompasses part of southwest Nevada and part of San Bernardino County, California. For the Las Vegas Valley the region is classified by the EPA as "nonattainment" for oxidants, carbon monoxide, and particulate matter. For the remaining major pollutants (NO_2 , SO_2), this portion of Nevada is classified as "unclassified or better than national standards." For the Nevada portion of the project area near Lake Mead southwest to the California/Nevada border, excluding Las Vegas Valley, the region is classified by EPA as an "attainment" area for all major pollutants.

That portion of the project area located in San Bernardino County, part of the Southeast Desert Air Basin (from the California/Nevada border parallel or nearly parallel to Interstate 15 southwest to and including Victorville and the surrounding area), is classified as "nonattainment" for oxidants and particulate matter. All other major pollutants (SO_2 , NO_x , CO) are classified as "unclassified" or "unclassified or better than national standards." It is generally believed that violations of the oxidant and particulate standards result from the transport of pollutants into the area from the South Coast Air Basin to the southwest as well as wind-blown fugitive dust (40 CFR 81).

Oxidants

Oxidants are formed in the atmosphere through a series of complex reactions, primarily involving reactive hydrocarbons and nitrogen oxides in the presence of sunlight. Ozone is the principal oxidant compound in the California atmosphere (California ARB 1976). For this reason, and the fact that measured oxidant concentrations are often reported as ozone, oxidant and ozone are often considered synonymous.

Maximum hourly ozone concentrations generally occur during afternoon hours between April and September. Concentrations are generally higher in rural areas than in adjacent urban areas, probably because of the scavenging effect of urban nitric oxide emissions on ozone concentrations.

Maximum one-hour concentrations measured during 1981-1983 in the project vicinity are shown in Table 2-7. Also shown are the number of hours and calendar days when the CAAQS were reached and/or exceeded and when the NAAQS were exceeded. These data indicate that both Barstow and Victorville, in San Bernardino County, exceed the Federal standard for ozone (0.12 ppm) and the state standard (0.10 ppm) a few times a year. Measured ozone levels in Victorville reached pollution levels high enough to warrant a Stage I episode (0.20 ppm) during two of the three years. Data collected at

Las Vegas and Henderson, in Clark County, indicate infrequent exceedances of the Nevada and Federal ozone standard (0.12 ppm).

Carbon Monoxide

Most carbon monoxide emissions occur in urban areas, primarily due to motor vehicles, coal combustion sources, and industrial processes. In nonurban areas, carbon monoxide emissions result primarily from forest fires and agricultural burning.

Maximum one-hour and eight-hour average CO concentrations for the project area are shown in Table 2-8. Also shown are the number of exceedances of the CO standards. There were no violations of the one-hour or eight-hour standards for CO in the past three years for the California monitoring stations presented. The eight-hour CO standard (9 ppm) was exceeded in Las Vegas all three years (1981-1983). The one-hour standard was not exceeded during this period.

Nitrogen Dioxide

Nitrogen dioxide (NO_2) is an indirect product of fuel combustion by industrial sources and motor vehicles. NO_x emissions are typically emitted largely as nitric oxide (NO) and are subsequently oxidized to nitrogen dioxide (NO_2) in the atmosphere. Diurnal NO_x concentration trends show peaks correlated with morning and evening commuter traffic. NO_2 concentrations increase steadily after the morning NO peak and remain relatively high until after midnight. This phenomenon is attributed to the oxidation of NO to NO_2 .

The maximum one-hour average and annual average NO_2 concentrations measured in the project area are shown on Table 2-9. The one-hour CAAQS for NO_2 (0.25 ppm or $470 \mu\text{g}/\text{m}^3$) was violated in 1981 at Barstow. The annual average NAAQS and Nevada standard for NO_2 is 0.05 ppm ($100 \mu\text{g}/\text{m}^3$). The annual standard has not been violated in the project area.

Sulfur Dioxide

Sulfur dioxide (SO_2) is primarily a product of coal and fuel oil combustion processes. In the project area, most SO_2 emissions result from fuel oil burning sources, such as cement kilns, scattered throughout the county. The maximum SO_2 concentrations generally occur near these sources.

The only monitoring data available for sulfur dioxide are the SO_2 concentrations measured at Southeast Desert Air Basin, Trona-Market St. The SO_2 ambient standards were not equaled or exceeded.

Particulate Matter

Wind-blown fugitive dust is a major source of ambient particulate concentrations in the project area. The maximum TSP concentrations measured in Barstow, Victorville, Las Vegas, and Henderson for 1981 to 1983 are shown in Table 2-10. Also shown are the number of secondary 24-hour NAAQS exceedances for each site. There were no exceedances of the primary NAAQS or secondary annual NAAQS. The CAAQS for particulates are based on ambient concentrations of fine particulates (particles less than 10 microns in diameter). There are no monitoring data available in the project area for comparison to the new California standards.

Lead

Ambient concentrations of lead are primarily the result of lead additives in automobile fuel. Lead is measured as a particulate pollutant in the same manner as TSP or sulfates. The CAAQS for a 30-day averaging period is $1.5 \mu\text{g}/\text{m}^3$, identical to the calendar quarter NAAQS and Nevada state standard.

The maximum measured suspended lead concentrations for 1981-1983 measured at Barstow, Victorville, and Las Vegas were below regulated standards. The maximum calendar quarter mean suspended lead concentration in the project area was $1.29 \mu\text{g}/\text{m}^3$ in Las Vegas. The measured suspended lead concentrations at Barstow and Victorville were well below the standard ($1.5 \mu\text{g}/\text{m}^3$).

Sulfate

Suspended particulates containing sulfate ions (SO_4) include both manmade and natural sources. Sulfates can result from the oxidation of SO_2 , but they are also natural components of soils.

The maximum measured 24-hour sulfate concentrations measured at Barstow and Victorville from 1981 to 1983 indicate no violations of the SO_4 CAAQS. There are no sulfate data available for Clark County. There is no national standard for sulfates.

Nitrates

Nitrate (NO_3) concentrations have been measured at Barstow and Victorville, but there are no data available for the Nevada portion of the project area. The maximum 24-hour concentration measured during 1981-1983 is $27.9 \mu\text{g}/\text{m}^3$ at the Victorville monitoring site. There are no Federal or state standards for nitrates.

IMPACT ASSESSMENT/MITIGATION PLANNING

This section presents a discussion of the emissions attributable to construction and operation activities and the potential air quality impacts associated with the proposed Mead/McCullough-Victorville/Adelanto 500kV transmission line.

Emission sources during construction will be associated with the following activities:

- Access road construction
- Clearing right-of-way and tower sites
- Installing foundations and towers (including construction yards and batch plants)
- Assembling and erecting towers
- Stringing, tensioning, and clipping conductors
- Installing of counterpoise
- Clean-up and restoration

These activities result in temporary land disturbances and require the operation of miscellaneous heavy-duty mobile diesel-powered construction equipment.

Emission sources during operation are expected to include maintenance vehicle exhausts, emergency maintenance vehicle and equipment exhausts, and high voltage transmission line ozone and nitrogen oxides generation.

Effects of Construction

Construction activities could occur on a six days per week, eight hours per day schedule over a period of approximately 24 months. The construction equipment to be used during the various phases of construction is described in the Project Description section. Emissions during construction would be associated primarily with fugitive dust and exhaust from gasoline and diesel-fired construction equipment.

Construction activities would result in temporary emissions increases at localized areas as construction proceeds along the approximately 200-mile-long transmission line route over a period of about 24 months. Fugitive dust emissions would be mitigated using water sprays. Criteria pollutant emissions would be minimized through conformance with existing tailpipe exhaust standards for applicable equipment. Proper maintenance of construction equipment and vehicles would help minimize emissions. Consequently, no long-term significant impacts on air quality would occur.

Effects of Operation

Operation activities would occur throughout the life of the proposed transmission line project. These activities include: transmission line inspection performed on a regular basis, maintenance and emergency maintenance performed as needed, and operation of the transmission line. Emissions during operation would be primarily associated with airborne dust from unpaved access routes and repair crew vehicle traffic. Small amounts of ozone and nitrogen oxides would be generated from the high voltage line. Emissions associated with the operation phase should be minimal and would not have a significant impact on regional air quality. The emissions associated with the operation of high voltage transmission lines, even during worst-case conditions, have been shown to be relatively small (Droppo 1979). Criteria pollutant emissions would be minimized through conformance with existing tailpipe exhaust standards for applicable equipment. Proper maintenance of construction equipment and vehicles would also help minimize emissions.

REFERENCES

California Air Resources Board. 1984. California air quality data, summary of 1983 air quality data, gaseous and particulate pollutants, annual summary, Technical Services Division. Vol. XV.

_____. 1983. California air quality data, summary of 1982 air quality data, gaseous and particulate pollutants, annual summary, Technical Services Division. Vol. XIV.

_____. 1982. California air quality data, summary of 1981 air quality data, gaseous and particulate pollutants, annual summary, Technical Services Division. Vol. XIII.

California Department of Water Resources. 1970. Wind in California. Bulletin No. 185.

Droppo, J.G. February 1979. "Field Determinations of HVDC Ozone Production Rates." Battelle Pacific Northwest Laboratory.

Federal Register. March 20, 1984. Proposed revisions to the National Ambient Quality Standards for particulate matter.

National Oceanic and Atmospheric Administration. 1982. Annual summary with comparative data, Las Vegas, Nevada. National Climatic Center, Asheville, North Carolina.

TABLE 1-1
WIND SPEED FREQUENCY
Frequency in Percent of Total Characteristic

Estimate for Force from
Table 1-1

Wind Speed	Wind Speed Class (mph)						General Notes	Total	Mean Speed
	2.00 to 2.99	3.00 to 3.99	4.00 to 4.99	5.00 to 5.99	6.00 to 6.99	7.00 to 7.99			
10-12	1.15	.73	.29	.14	.07	.03	0.86	1.00	2.72
13-15	1.25	.76	.31	.14	.07	.04	0.89	1.25	2.80
16-18	1.40	.81	.33	.15	.07	.05	0.90	1.40	2.85
19-21	1.55	.87	.35	.16	.08	.06	0.91	1.55	2.90
22-24	1.70	.93	.37	.17	.09	.07	0.92	1.70	2.95
25-27	1.85	.99	.39	.18	.10	.08	0.93	1.85	3.00
28-30	2.00	1.05	.41	.19	.11	.09	0.94	2.00	3.05
31-33	2.15	1.11	.43	.20	.12	.10	0.95	2.15	3.10
34-36	2.30	1.17	.45	.21	.13	.11	0.96	2.30	3.15
37-39	2.45	1.23	.47	.22	.14	.12	0.97	2.45	3.20
40-42	2.60	1.29	.49	.23	.15	.13	0.98	2.60	3.25
43-45	2.75	1.35	.51	.24	.16	.14	0.99	2.75	3.30
46-48	2.90	1.41	.53	.25	.17	.15	1.00	2.90	3.35
49-51	3.05	1.47	.55	.26	.18	.16	1.01	3.05	3.40
52-54	3.20	1.53	.57	.27	.19	.17	1.02	3.20	3.45
55-57	3.35	1.59	.59	.28	.20	.18	1.03	3.35	3.50
58-60	3.50	1.65	.61	.29	.21	.19	1.04	3.50	3.55
61-63	3.65	1.71	.63	.30	.22	.20	1.05	3.65	3.60
64-66	3.80	1.77	.65	.31	.23	.21	1.06	3.80	3.65
67-69	3.95	1.83	.67	.32	.24	.22	1.07	3.95	3.70
70-72	4.10	1.89	.69	.33	.25	.23	1.08	4.10	3.75
73-75	4.25	1.95	.71	.34	.26	.24	1.09	4.25	3.80
76-78	4.40	2.01	.73	.35	.27	.25	1.10	4.40	3.85
79-81	4.55	2.07	.75	.36	.28	.26	1.11	4.55	3.90
82-84	4.70	2.13	.77	.37	.29	.27	1.12	4.70	3.95
85-87	4.85	2.19	.79	.38	.30	.28	1.13	4.85	4.00
88-90	5.00	2.25	.81	.39	.31	.29	1.14	5.00	4.05
91-93	5.15	2.31	.83	.40	.32	.30	1.15	5.15	4.10
94-96	5.30	2.37	.85	.41	.33	.31	1.16	5.30	4.15
97-99	5.45	2.43	.87	.42	.34	.32	1.17	5.45	4.20
100-102	5.60	2.49	.89	.43	.35	.33	1.18	5.60	4.25
103-105	5.75	2.55	.91	.44	.36	.34	1.19	5.75	4.30
106-108	5.90	2.61	.93	.45	.37	.35	1.20	5.90	4.35
109-111	6.05	2.67	.95	.46	.38	.36	1.21	6.05	4.40
112-114	6.20	2.73	.97	.47	.39	.37	1.22	6.20	4.45
115-117	6.35	2.79	.99	.48	.40	.38	1.23	6.35	4.50
118-120	6.50	2.85	1.01	.49	.41	.39	1.24	6.50	4.55
121-123	6.65	2.91	1.03	.50	.42	.40	1.25	6.65	4.60
124-126	6.80	2.97	1.05	.51	.43	.41	1.26	6.80	4.65
127-129	6.95	3.03	1.07	.52	.44	.42	1.27	6.95	4.70
130-132	7.10	3.09	1.09	.53	.45	.43	1.28	7.10	4.75
133-135	7.25	3.15	1.11	.54	.46	.44	1.29	7.25	4.80
136-138	7.40	3.21	1.13	.55	.47	.45	1.30	7.40	4.85
139-141	7.55	3.27	1.15	.56	.48	.46	1.31	7.55	4.90
142-144	7.70	3.33	1.17	.57	.49	.47	1.32	7.70	4.95
145-147	7.85	3.39	1.19	.58	.50	.48	1.33	7.85	5.00
148-150	8.00	3.45	1.21	.59	.51	.49	1.34	8.00	5.05
151-153	8.15	3.51	1.23	.60	.52	.50	1.35	8.15	5.10
154-156	8.30	3.57	1.25	.61	.53	.51	1.36	8.30	5.15
157-159	8.45	3.63	1.27	.62	.54	.52	1.37	8.45	5.20
160-162	8.60	3.69	1.29	.63	.55	.53	1.38	8.60	5.25
163-165	8.75	3.75	1.31	.64	.56	.54	1.39	8.75	5.30
166-168	8.90	3.81	1.33	.65	.57	.55	1.40	8.90	5.35
169-171	9.05	3.87	1.35	.66	.58	.56	1.41	9.05	5.40
172-174	9.20	3.93	1.37	.67	.59	.57	1.42	9.20	5.45
175-177	9.35	3.99	1.39	.68	.60	.58	1.43	9.35	5.50
178-180	9.50	4.05	1.41	.69	.61	.59	1.44	9.50	5.55
181-183	9.65	4.11	1.43	.70	.62	.60	1.45	9.65	5.60
184-186	9.80	4.17	1.45	.71	.63	.61	1.46	9.80	5.65
187-189	9.95	4.23	1.47	.72	.64	.62	1.47	9.95	5.70
200-202	10.10	4.29	1.49	.73	.65	.63	1.48	10.10	5.75
203-205	10.25	4.35	1.51	.74	.66	.64	1.49	10.25	5.80
206-208	10.40	4.41	1.53	.75	.67	.65	1.50	10.40	5.85
209-211	10.55	4.47	1.55	.76	.68	.66	1.51	10.55	5.90
212-214	10.70	4.53	1.57	.77	.69	.67	1.52	10.70	5.95
215-217	10.85	4.59	1.59	.78	.70	.68	1.53	10.85	6.00
218-220	11.00	4.65	1.61	.79	.71	.69	1.54	11.00	6.05
221-223	11.15	4.71	1.63	.80	.72	.70	1.55	11.15	6.10
224-226	11.30	4.77	1.65	.81	.73	.71	1.56	11.30	6.15
227-229	11.45	4.83	1.67	.82	.74	.72	1.57	11.45	6.20
230-232	11.60	4.89	1.69	.83	.75	.73	1.58	11.60	6.25
233-235	11.75	4.95	1.71	.84	.76	.74	1.59	11.75	6.30
236-238	11.90	5.01	1.73	.85	.77	.75	1.60	11.90	6.35
239-241	12.05	5.07	1.75	.86	.78	.76	1.61	12.05	6.40
242-244	12.20	5.13	1.77	.87	.79	.77	1.62	12.20	6.45
245-247	12.35	5.19	1.79	.88	.80	.78	1.63	12.35	6.50
248-250	12.50	5.25	1.81	.89	.81	.79	1.64	12.50	6.55
251-253	12.65	5.31	1.83	.90	.82	.80	1.65	12.65	6.60
254-256	12.80	5.37	1.85	.91	.83	.81	1.66	12.80	6.65
257-259	12.95	5.43	1.87	.92	.84	.82	1.67	12.95	6.70
260-262	13.10	5.49	1.89	.93	.85	.83	1.68	13.10	6.75
263-265	13.25	5.55	1.91	.94	.86	.84	1.69	13.25	6.80
266-268	13.40	5.61	1.93	.95	.87	.85	1.70	13.40	6.85
269-271	13.55	5.67	1.95	.96	.88	.86	1.71	13.55	6.90
272-274	13.70	5.73	1.97	.97	.89	.87	1.72	13.70	6.95
275-277	13.85	5.79	1.99	.98	.90	.88	1.73	13.85	7.00
278-280	14.00	5.85	2.01	.99	.91	.89	1.74	14.00	7.05
281-283	14.15	5.91	2.03	1.00	.92	.90	1.75	14.15	7.10
284-286	14.30	5.97	2.05	1.01	.93	.91	1.76	14.30	7.15
287-289	14.45	6.03	2.07	1.02	.94	.92	1.77	14.45	7.20
290-292	14.60	6.09	2.09	1.03	.95	.93	1.78	14.60	7.25
293-295	14.75	6.15	2.11	1.04	.96	.94	1.79	14.75	7.30
296-298	14.90	6.21	2.13	1.05	.97	.95	1.80	14.90	7.35
299-301	15.05	6.27	2.15	1.06	.98	.96	1.81	15.05	7.40
302-304	15.20	6.33	2.17	1.07	.99	.97	1.82	15.20	7.45
305-307	15.35	6.39	2.19	1.08	1.00	.98	1.83	15.35	7.50
308-310	15.50	6.45	2.21	1.09	1.01	.99	1.84	15.50	7.55
311-313	15.65	6.51	2.23	1.10	1.02	1.00	1.85	15.65	7.60
314-316	15.80	6.57	2.25	1.11	1.03	1.01	1.86	15.80	7.65
317-319	15.95	6.63	2.27	1.12	1.04	1.02	1.87	15.95	7.70
320-322	16.10	6.69	2.29	1.13	1.05	1.03	1.88	16.10	7.75
323-325	16.25	6.75	2.31	1.14	1.06	1.04	1.89	16.25	7.80
326-328	16.40	6.81	2.33	1.15	1.07	1.05	1.90	16.40	7.85
329-331	16.55	6.87	2.35	1.16	1.08	1.06	1.91	16.55	7.90
332-334	16.70	6.93	2.37	1.17	1.09	1.07	1.92	16.70	7.95
335-337	16.85	6.99	2.39	1.18	1.10	1.08	1.93	16.85	8.00
338-340	17.00	7.05	2.41	1.19	1.11	1.09	1.94	17.00	8.05
341-343	17.15	7.11	2.43	1.20	1.12	1.10	1.95	17.15	8.10
344-346	17.30	7.17	2.45	1.21	1.13	1.11	1.96	17.30	8.15
347-349	17.45	7.23	2.47	1.22	1.14	1.12	1.97	17.45	8.20
350-352	17.60	7.29	2.49	1.23	1.15	1.13	1.98	17.60	8.25
353-355	17.75	7.35	2.51	1.24	1.16	1.14	1.99	17.75	8.30
356-358	17.90	7.41	2.53	1.25	1.17	1.15	2.00	17.90	8.35
359-361	18.05	7.47	2.55	1.26	1.18	1.16	2.01	18.05	8.40
362-364	18.20	7.53	2.57	1.27	1.19	1.17	2.02	18.20	8.45
365-367	18.35	7.59	2.59	1.28	1.20	1.18	2.03	18.35	8.50
368-370	18.50	7.65	2.61	1.29	1.21	1.19	2.04	18.50	8.55
371-373	18.65	7.71	2.63	1.30	1.22	1.20	2.05	18.65	8.60
374-376	18.80	7.77	2.65	1.31	1.23	1.21	2.06	18.80	8.65
377-379	18.95	7.83	2.67	1.32	1.24	1.22	2.07	18.95	8.70
380-382	19.10	7.89	2.69	1.33	1.25	1.23	2.08	19.10	8.75
383-385	19.25	7.95	2.71	1.34	1.26	1.24	2.09	19.25	8.80
386-388	19.40	8.01	2.73	1.35	1.27	1.25	2.10	19.40	8.85
389-391	19.55	8.07	2.75	1.36	1.28	1.26	2.11	19.55	8.90
392-394	19.70	8.13	2.77	1.37	1.29	1.27	2.12	19.70	8.95
395-397	19.85	8.19	2.79	1.38	1.30	1.28	2.13	19.85	9.00
398-400	20.00	8.25	2.81	1.39	1.31</				

TABLE 2-1
WIND FREQUENCY DISTRIBUTION
(Frequency in Percent of Total Observations)

Edwards Air Force Base
1968-1970

Wind Sector	Wind Speed Class (mps)								Greater Than 15.00	Total	Mean Speed
	.76 to 2.00	2.01 to 3.00	3.01 to 4.00	4.01 to 5.00	5.01 to 7.00	7.01 to 10.00	10.1 to 15.00				
NNE	2.30	1.13	.70	.24	.16	.09	.03	0.00	4.65	2.22	
NE	1.95	.67	.36	.21	.26	.14	.06	0.00	3.65	2.46	
ENE	1.40	.51	.41	.24	.48	.21	.05	0.00	3.29	2.98	
E	1.14	.42	.28	.14	.35	.18	.02	0.00	2.54	2.93	
ESE	.86	.24	.18	.07	.06	.03	.00	0.00	1.44	2.09	
SE	.75	.18	.09	.04	.05	.02	.02	0.00	1.14	1.97	
SSE	.71	.26	.10	.04	.02	.02	.00	0.00	1.16	1.88	
S	1.22	.51	.34	.10	.05	.02	.00	0.00	2.24	2.01	
SSW	1.93	1.29	1.40	.87	.53	.16	.05	0.00	6.23	2.91	
SW	3.06	2.66	3.14	2.69	3.07	2.08	.71	0.00	17.40	4.24	
WSW	3.06	1.94	2.35	2.55	5.34	4.71	1.08	0.00	21.03	5.13	
W	3.27	1.28	1.07	1.35	3.03	2.81	.71	.01	13.54	4.76	
WNW	3.36	1.41	.49	.43	1.29	1.75	1.19	.05	9.95	4.65	
NW	3.09	.98	.32	.25	.52	.85	.50	.01	6.52	3.61	
NNW	2.03	.36	.17	.06	.08	.07	.01	0.00	2.79	1.77	
N	1.64	.43	.21	.06	.05	.02	.00	0.00	2.41	1.71	
CALM (5.75 mps)								0.00			
TOTAL	31.76	14.26	11.61	9.356	15.34	13.15	4.45	0.08	100.00	3.92	

Source: National Weather Service, summary of hourly surface data.

TABLE 2-2
PERCENTAGE FREQUENCY OF WIND DIRECTION AND SPEED
(From Hourly Observations)

Daggett, California
1955-1964

Wind Sector	Wind Speed Class (knots)						%	Mean Wind Speed (knots)
	1-3	4-6	7-10	11-16	17-21	21		
N	0.1	0.6	0.5	0.2	0.0	0.0	1.4	7.5
NNE	0.1	0.4	0.4	0.2	0.1	0.0	1.2	8.5
NE	0.1	1.0	1.0	0.2	0.0		2.4	7.4
ENE	0.1	0.7	0.9	0.3	0.0	0.0	1.9	7.7
E	0.1	1.3	1.6	0.6	0.1	0.0	3.7	8.0
ESE	0.1	0.7	0.7	0.3	0.1	0.0	1.9	8.1
SE	0.1	0.8	0.4	0.1	0.0	0.0	1.4	6.7
SSE	0.0	0.3	0.1	0.0	0.0	0.0	0.5	6.8
S	0.0	0.3	0.1	0.0	0.0	0.0	0.6	6.9
SSW	0.0	0.2	0.1	0.1	0.1	0.0	0.5	9.9
SW	0.1	0.5	0.6	0.8	1.0	0.4	3.3	14.1
WSW	0.1	0.9	2.1	3.4	3.8	1.3	11.6	15.1
W	0.1	2.7	8.7	6.3	3.3	1.4	22.5	12.1
WNW	0.1	3.2	11.0	8.1	2.0	0.5	24.9	10.8
NW	0.1	1.9	4.2	2.2	0.2	0.0	8.6	9.2
NNW	0.1	0.6	0.7	0.3	0.0	0.0	1.7	8.0
CALM							11.8	
TOTAL	1.3	16.0	33.4	23.0	10.8	3.7	100.0	9.8

Source: Wind in California, Bulletin No. 185.

TABLE 2-3
WIND DIRECTION FREQUENCY DISTRIBUTION
NIPTON: 6/19/79 - 8/31/79
CADIZ: 7/6/79 - 8/31/79
(Percent)

Wind Sector	Hour (PST)									
	0000-0500		0600-1100		1200-1700		1800-2300		All Hours	
	Nipton	Cadiz	Nipton	Cadiz	Nipton	Cadiz	Nipton	Cadiz	Nipton	Cadiz
N	1.1	2.9	2.1	3.6	1.0	0.5	1.2	1.9	1.3	3.2
NNE	0.4	2.6	0.9	2.8	0.5	0.3	0.6	1.6	0.6	1.7
NE	6.1	7.0	3.6	4.8	0.5	0.9	2.6	4.4	3.2	5.3
ENE	17.7	7.7	3.7	4.8	0.7	2.5	5.3	5.3	6.7	4.8
E	21.8	7.8	6.5	9.8	1.1	3.2	10.8	2.9	10.0	5.9
ESE	17.0	13.9	4.8	14.2	1.2	8.0	14.4	4.9	9.3	11.7
SE	8.9	15.5	7.5	15.4	2.6	24.4	9.8	22.2	7.2	15.8
SSE	9.2	9.9	10.9	7.6	13.5	16.5	14.1	14.6	11.9	11.7
S	10.7	5.6	11.4	5.6	13.7	14.1	11.6	11.1	11.8	7.6
SSW	3.4	4.6	14.1	4.7	18.9	5.2	9.6	9.2	11.5	3.1
SW	0.9	4.5	8.7	5.8	26.6	8.0	9.1	5.4	11.5	5.6
WSW	0.3	5.1	7.0	6.1	6.6	6.1	4.7	6.0	4.7	4.5
W	0.3	3.1	7.4	3.9	5.5	2.1	2.3	3.7	3.8	6.1
WNW	0.8	4.7	6.8	3.3	4.4	3.0	0.8	3.4	3.2	7.1
NW	0.6	3.3	2.9	2.9	2.8	1.3	2.4	1.3	2.2	2.6
NNW	0.3	2.0	0.9	1.7	0.2	1.8	0.9	1.1	0.6	1.3
CALM		1.9		3.0		2.2		1.3		2.3

Source: Onsite Monitoring Program - Southern California Edison Company.

TABLE 2-4
MEAN TEMPERATURE
(degrees F)

Location	No. of Years Recorded	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec	Annual
Barstow	36	45.9	50.1	55.7	62.4	68.9	78.0	84.4	82.6	75.6	64.7	53.6	46.3	64.0
Daggett	17	46.8	51.4	56.5	56.2	72.1	80.4	87.2	85.0	80.1	68.1	55.0	48.2	66.3
Las Vegas	30	44.2	49.1	54.8	63.8	73.3	82.3	89.6	87.4	80.1	67.1	53.3	45.2	65.8
Yucca Grove	21	39.3	42.9	48.0	55.2	63.5	71.4	79.7	78.0	71.0	59.8	49.0	42.7	58.5
Victorville	21	42.4	45.6	50.4	57.8	64.0	71.5	78.6	77.4	71.7	60.9	49.2	43.9	59.5

Source: Climatic Summary of the United States, California.
Climatography of the United States, No. 86-4.
Local Climatological Data - Las Vegas, Nevada.

TABLE 2-5
PRECIPITATION

Location	No. of Years Recorded	Average Precipitation (inches)												
		Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec	Annual
Baker	7	0.47	0.22	0.29	0.19	0.11	0.01	0.38	0.31	0.13	0.38	0.36	0.13	2.98
Barstow	48	0.80	0.54	0.64	0.19	0.08	0.09	0.26	0.23	0.20	0.34	0.32	0.58	4.27
Daggett	17	0.60	0.33	0.30	0.21	0.03	NA ^a	0.31	0.50	0.33	0.19	0.32	0.48	3.60
Las Vegas	30	0.45	0.30	0.33	0.27	0.10	0.09	0.44	0.49	0.27	0.22	0.43	0.37	3.76
Victorville	22	1.17	0.86	0.85	0.42	0.06	NA	0.06	0.13	0.12	0.29	0.52	0.93	5.41

^a NA = Data not available.

Source: Climatology of the United States, No. 86-4.
Local Climatological Data - Las Vegas, Nevada.

TABLE 2-6
 AMBIENT AIR QUALITY STANDARDS^a

Pollutant	Averaging Time	California Standards ^{b,c,d}	Nevada Standards ^{d,e}	National Standards ^a	
				Primary ^{d,f}	Secondary ^{d,g}
Oxidant (Ozone)	1-hour	0.10 ppm (200 $\mu\text{g}/\text{m}^3$)	235 $\mu\text{g}/\text{m}^3$ (0.12 ppm)	235 $\mu\text{g}/\text{m}^3$ (0.12 ppm)	Same as primary standard
Carbon monoxide	8-hour	9 ppm (10 mg/m^3)	10 mg/m^3 (9 ppm)	10 mg/m^3 (9 ppm)	Same as primary standard
	1-hour	20 ppm (23 mg/m^3)	40 mg/m^3 (35 ppm)	40 mg/m^3 (35 ppm)	Same as primary standard
Nitrogen dioxide	Annual average	--	100 $\mu\text{g}/\text{m}^3$ (0.05 ppm)	100 $\mu\text{g}/\text{m}^3$ (0.05 ppm)	Same as primary standard
	1-hour	0.25 ppm (470 $\mu\text{g}/\text{m}^3$)	--	--	--
Sulfur dioxide	Annual average	--	80 $\mu\text{g}/\text{m}^3$ (0.03 ppm)	80 $\mu\text{g}/\text{m}^3$ (0.03 ppm)	--
	24-hour	0.05 ppm ^h (131 $\mu\text{g}/\text{m}^3$)	365 $\mu\text{g}/\text{m}^3$ (0.14 ppm)	365 $\mu\text{g}/\text{m}^3$ (0.14 ppm)	--
	3-hour	--	1300 $\mu\text{g}/\text{m}^3$ (0.5 ppm)	--	1300 $\mu\text{g}/\text{m}^3$ (0.5 ppm)
	1-hour	0.25 ppm (655 $\mu\text{g}/\text{m}^3$)	--	--	--
Suspended particulate matter	Annual geometric mean	--	75 $\mu\text{g}/\text{m}^3$	75 $\mu\text{g}/\text{m}^3$	60 $\mu\text{g}/\text{m}^3$ ⁱ
	24-hour	--	150 $\mu\text{g}/\text{m}^3$	260 $\mu\text{g}/\text{m}^3$	150 $\mu\text{g}/\text{m}^3$
Fine particulate matter ($\leq 10 \mu\text{m}$)	24-hour	50 $\mu\text{g}/\text{m}^3$	--	--	--
	Annual geometric mean	30 $\mu\text{g}/\text{m}^3$	--	--	--
Sulfates	24-hour	25 $\mu\text{g}/\text{m}^3$	--	--	--

Table 2-6 (continued)
Ambient Air Quality Standards^a

Pollutant	Averaging Time	California Standards ^{b,c,d}	Nevada Standards ^{d,e}	National Standards ^a	
				Primary ^{d,f}	Secondary ^{d,g}
Lead	30-day	1.5 $\mu\text{g}/\text{m}^3$	--	--	--
	Calendar quarter	--	1.5 $\mu\text{g}/\text{m}^3$	1.5 $\mu\text{g}/\text{m}^3$	Same as primary standard
Hydrogen sulfide	1-hour	0.03 ppm (42 $\mu\text{g}/\text{m}^3$)	--	--	--
Vinyl Chloride	24-hour	0.010 ppm (26 $\mu\text{g}/\text{m}^3$)	--	--	--
Ethylene	8-hour	0.1 ppm	--	--	--
	1-hour	0.5 ppm	--	--	--
Visibility reducing particles	1 observation	In sufficient amount to reduce the prevailing visibility to less than 10 miles when the relative humidity is less than 70 percent ^j	In sufficient amount to reduce the prevailing visibility to less than 10 miles when the relative humidity is less than 70 percent ^j	--	--

^a National standards, other than those based on annual averages or annual geometric means, are not to be exceeded more than once per year.

^b Standards from California Air Resources Board (1984).

^c California standards are values that are not to be equaled or exceeded.

^d Concentration expressed first in units in which it was promulgated. Equivalent units given in parentheses are based upon a reference temperature of 25°C and a reference pressure of 760 mm of Hg (1,013.2 millibars); ppm in this table refers to ppm by volume, or micromoles of pollutant per mole of gas.

^e Nevada standards are values that should not be exceeded.

^f National Primary Standards: The level of air quality necessary, with an adequate margin of safety, to protect public health.

^g National Secondary Standards: The level of air quality necessary to protect the public welfare from any known or anticipated adverse effects of a pollutant.

^h 0.05 ppm (131 $\mu\text{g}/\text{m}^3$) (conductimetric) in the presence of oxidant in excess of state 1-hour standard or in presence of particulates in excess of state 24-hour standard (established June 29, 1977).

ⁱ For guidance to achieve 24-hour standard.

^j Prevailing visibility is defined as the greatest visibility which is attained or surpassed around at least half of the horizon circle, but not necessarily in continuous sectors.

TABLE 2-7
OXIDANT CONCENTRATIONS AND EXCEEDANCES

Maximum Measured 1-Hour Oxidant Concentrations
in Project Area
(concentrations in ppm)

Year	Monitoring Site			
	Barstow	Victorville-Fairgrounds	Las Vegas	Henderson
1981	0.16	0.21	0.113	0.148
1982	0.16 ^a	0.20	0.114	0.141
1983	0.15	0.23	0.138	0.120

Number of Hours and Calendar Days of Oxidant
Standards Exceedances in Project Area

Standard/Year	Monitoring Site							
	Barstow		Victorville-Fairgrounds		Las Vegas		Henderson	
	Hours	Days	Hours	Days	Hours	Days	Hours	Days
CAAQS (0.10 ppm)								
1981	163	50	459	104	--	--b	--	--b
1982	58	25	367	85	--	--b	--	--b
1983	318	70	505	91	--	--b	--	--b
NAAQS (0.12 ppm)								
1981	20	13	87	39	0	0	18	14
1982	19	12	118	35	0	0	4	4
1983	9	5	60	21	4	2	0	0

^a Statistics do not meet California ARB criteria for representativeness.

^b Standard does not exist in Nevada.

TABLE 2-8
CARBON MONOXIDE CONCENTRATIONS AND EXCEEDANCES

Maximum Measured CO Concentrations
in Project Area
(concentrations in ppm)

Averaging Period/ Year	Monitoring Site			
	Barstow	Victorville-Fairgrounds	Las Vegas	Henderson
1-Hour Average				
1981	6.0	6.0 ^a	21.1	NA
1982	5.0 ^a	5.0 ^a	14.1	7.8
1983	8.0	NA ^b	24.0	4.7
8-Hour Average				
1981	2.10	6.00 ^a	11.4	--c
1982	3.30 ^a	3.50 ^a	18.8	--c
1983	2.90	NA	16.0	--c

Number of Hours and Calendar Days of
CO Standards Exceedances in Project Area

8-Hour NAAQS (9 ppm)	Monitoring Site							
	Barstow		Victorville-Fairgrounds		Las Vegas		Henderson	
	Hours	Days	Hours	Days	Hours	Days	Hours	Days
Year								
1981	0	0	0	0	NA	34	NA	NA
1982	0	0	0	0	NA	29	NA	NA
1983	0	0	NA	NA	NA	35	NA	NA

^a Statistics do not meet California ARB criteria for representativeness.

^b NA = Data not available.

^c Data are not collected if the high 1-hour concentration does not exceed 3.7 ppm (10 mg/m³).

TABLE 2-9
MAXIMUM MEASURED NITROGEN DIOXIDE CONCENTRATIONS
IN PROJECT AREA

Averaging Period/ Year	Monitoring Site (Concentrations in ppm)		
	Barstow	Victorville-Fairgrounds	Las Vegas
1-Hour Average			
1981	0.30	0.00 ^a	0.203
1982	0.20	0.12 ^a	0.206
1983	0.15 ^a	0.13	0.138
Annual Average			
1981	0.025	0.000 ^a	NA ^b
1982	0.025	0.020 ^a	0.025
1983	0.023 ^a	0.021	0.024

^aStatistics do not meet California ARB criteria for representativeness.

^bNA = Data not available.

TABLE 2-10
PARTICULATE CONCENTRATIONS AND EXCEEDANCES

Maximum Measured Particulate Concentrations
in Project Area
(concentrations in $\mu\text{g}/\text{m}^3$)

Averaging Period/ Year	Monitoring Site			
	Barstow	Victorville-Fairgrounds	Las Vegas	Henderson
24-Hour Average				
1981	300	172	295	NA ^b
1982	116	237	267	201
1983	87	125	379	673
Annual Geometric Average				
1981	70.9	73.5	95.6	NA
1982	46.0 ^a	59.0 ^a	87.3	73.5
1983	64.2 ^a	61.6 ^a	89.6	80.0

Number of Measured Exceedances and Observations
of 24-Hour PM Standard

Standard/ Year	Monitoring Site							
	Barstow		Victorville- Fairgrounds		Las Vegas		Henderson	
	Exc. ^c	Obs. ^d	Exc.	Obs.	Exc.	Obs.	Exc.	Obs.
24-Hour Secondary NAAQS ($150 \mu\text{g}/\text{m}^3$)								
1981	2	55	1	61	NA	NA	NA	NA
1982	0	21	1	57	NA	NA	NA	NA
1983	0	20	0	56	NA	NA	NA	NA

^a Statistics do not meet California ARB criteria for representativeness.

^b NA = Data not available.

^c Exc. = Exceedances

^d Obs. = Observations

TABLE OF CONTENTS

	PAGE
CHAPTER 1 - EARTH RESOURCES	1-1
Study Area Overview	1-1
Study Purpose	1-2
Inventory Methods	1-2
Data Collection	1-2
Data Compilation	1-3
Inventory Results	1-3
Regional Physiographic Setting	1-3
Regional Geologic Setting	1-4
Paleontology	1-4
Minerals	1-4
Seismotectonic Setting	1-7
Hydrology	1-8
Soils	1-10
Geologic Hazards	1-12
Impact Assessment/Mitigation Planning Process	1-15
Impact Assessment Methods	1-15
Mitigation Planning Methods	1-16
Results	1-17
Appendix: Data and Methods Detail	1-20
References	1-21

CHAPTER 3 EARTH RESOURCES

TABLE OF CONTENTS

	PAGE
CHAPTER 3 - EARTH RESOURCES	3- 1
Study Area Overview	3- 1
Study Personnel	3- 2
Inventory Methods	3- 2
Data Collection	3- 2
Data Compilation	3- 3
Inventory Results	3- 3
Regional Physiographic Setting	3- 3
Regional Geologic Setting	3- 4
Paleontology	3- 6
Minerals	3- 6
Seismotectonic Setting	3- 7
Hydrology	3- 9
Soils	3-10
Geologic Hazards	3-12
Impact Assessment/Mitigation Planning Process	3-15
Impact Assessment Methods	3-15
Mitigation Planning Methods	3-19
Results	3-19
Agency Contacts (follows page)	3-20
References	3-21

LIST OF TABLES

TABLE NUMBER	TITLE	FOLLOWS PAGE
3-1	Mines and/or Mineral Deposits in Project Study Area	3-23
3-2	Routes Crossed by Quaternary Faults	3-23
3-3	Daily Discharges of Mojave River - Lower Narrows	3-23
3-4	Daily Discharges of Mojave River - Afton Canyon	3-23
3-5	Inventory of Geologic Resources	3-23
3-6	Inventory of Paleontological Resources	3-23
3-7	Soils Inventory Table	3-23
3-8	Soil Erosion Impact Assessment Model	3-23
3-9	Paleontological Resources Impact Assessment Model	3-23
3-10	Impact Assessment/Mitigation Planning Chart Soil Erosion	3-23
3-11	Impact Assessment/Mitigation Planning Paleontological Resources	3-23

CHAPTER 3 EARTH RESOURCES

This chapter contains the baseline inventory for Earth Resources for the proposed 500kV transmission line from the Mead or McCullough Switching Station in southern Nevada to the Victorville/Adelanto area in the southwestern portion of the Mojave Desert in California. Discussions presented in this report include geology, hydrology, soils, and geologic hazards.

STUDY AREA OVERVIEW

The proposed Mead/McCullough-Victorville/Adelanto Transmission Project lies within the Mojave Desert Physiographic Province of southeastern California and southern Nevada. The Mojave Desert is a wedge-shaped region southeast of the Sierra Nevada batholith and southwest of the Basin and Range Physiographic Province. The region is bounded by the Garlock Fault on the north and northwest, and by the San Andreas Fault on the southwest. The eastern boundary of the Mojave Desert is not well defined but, for the purposes of this report, is considered to be the Colorado River.

The study area is characterized by broad alluviated basins interrupted by isolated north- to northwest-trending mountain ranges. These ranges are generally flanked by alluvial fan complexes that grade into broad alluvial plains of little to no relief. Many of the intermontane valleys are closed drainage basins which contain playas at their lowest elevation.

The predominant types of basement rock in the study area are Precambrian, Paleozoic and Mesozoic intrusive igneous and metamorphic rocks. Other rock types in the region include Paleozoic, Mesozoic and Tertiary volcanic and sedimentary rocks. Most of the alternative transmission line routes traverse areas underlain by unconsolidated Quaternary-age alluvial, eolian and lacustrine (playa) deposits.

The tectonic setting of the western Mojave Desert is characterized by northwest-trending right-lateral faults which commonly offset Quaternary sediments. These faults trend generally parallel to the San Andreas Fault to the southwest. Minor east-west trending faults are present east and northeast of Barstow.

The eastern portion of the study area is seismically much less active than the western portion. With the exception of the Manix and Galway Lake events, only occasional, isolated, relatively small magnitude (less than 5.0 on the Richter Scale) earthquakes have historically occurred in the study area.

Economic mineral deposits are widespread in the study area. These deposits occur almost exclusively within the basement rocks, and include gold, silver, lead, zinc, copper, tungsten, fluorite, barite, bentonite, gypsum, borates, talc, and some rare earth elements.

STUDY PERSONNEL

This study was conducted under the supervision and guidance of Roy H. Patterson, with technical assistance from Robert E. Troutman and Jeff Jeter. Thomas W. Dibblee, who had previously mapped a large portion of the study area, made available to us unpublished geologic maps covering parts of the study area.

Mr. Patterson, a Senior Geologist in Dames & Moore's Los Angeles office, has eight years of experience in geological and geotechnical investigations in support of environmental impact analyses. He has extensive experience in environmental analysis for offshore oil development and was principal geologic investigator for environmental reports for oil development in the eastern and western Santa Barbara Channel and in the offshore Santa Maria Basin and for petrochemical facilities in the Los Angeles Basin. He participated in earth resources impact analyses for transmission lines associated with the Sundesert Nuclear Power Project for Wirth Associates. As geotechnical coordinator for the Belridge Cogeneration Project in Kern County, Mr. Patterson had overall responsibility for preparation of geotechnical, geologic and hydrogeologic aspects of an Application for Certification for a large cogeneration project. This project included route selection studies and impact analysis for a transmission line between the project site and an existing substation. Mr. Patterson is a registered geologist in the State of California and has an M.A. in Geology.

Mr. Troutman, a Staff Geologist in Dames & Moore's Santa Barbara office, has four years of experience in geological and geotechnical investigations in support of environmental impact analyses. Mr. Troutman has experience in environmental analyses for offshore oil development in the Santa Barbara Channel and Santa Maria Basin. In addition, he was a principal investigator for a transmission line project in the Tabuk region of Saudi Arabia and was also involved in the Belridge Cogeneration Project in Kern County. Mr. Troutman has participated in foundation investigations for a solar power plant near Daggett, California and performed an evaluation of potential hazardous waste disposal sites in the Colorado Desert region southeast of the project area.

Mr. Jeter, a recent graduate of Humboldt State University, was raised in the western Mojave Desert in Lancaster and has worked at Edwards Air Force Base immediately west of the project area.

INVENTORY METHODS

Data Collection

Compilation of the baseline inventory involved available geologic, hydrologic and soils literature and information. Data sources include published and unpublished reports, maps, and open-file reports of the United States Geological Survey (USGS), Bureau of Land Management (BLM), Soils Conservation Service (SCS); California Division of Mines and Geology (CDMG), Department

of Transportation; Nevada Bureau of Mines and Geology (NBMG); and publications from various scientific periodicals and journals. Representatives of the Federal and state agencies listed above and local governments were contacted as information sources. The geologic map was compiled using numerous sources, including published and unpublished maps from the USGS, CDMG, and NBMG. Data concerning historic seismicity were obtained from available CDMG and NBMG publications. Hydrologic information was obtained from the USGS Water Resources Division.

Data Compilation

Available geologic maps of portions of the study area at scales of 1:24,000, 1:62,500, 1:125,000, and 1:250,000 were transferred to maps at scales of 1:40,000 and 1:62,500 prior to final compilation at a scale of 1:126,720. The corridor width inventoried for geology and soils was generally two miles. The corridor width inventoried for mineral deposits was six miles.

INVENTORY RESULTS

Regional Physiographic Setting

The project study area traverses the Mojave Desert Physiographic Province of southern California and southern Nevada. The Mojave Desert is a wedge-shaped geomorphic province bounded by the Sierra Nevada and Garlock faults on the north, the San Andreas Fault on the southwest, and the Transverse Ranges and Colorado Desert on the south and southeast. The southern and eastern limits of the Mojave Desert are not well defined due to the lack of distinct geomorphic boundaries; however, most investigators recognize the eastern Transverse Ranges and Colorado Desert provinces as the southern and southeastern boundaries. Although some investigators include southernmost Nevada in the Basin and Range Physiographic Province, it is included, for the purposes of this report, in the Mojave Desert Physiographic Province.

The portion of the Mojave Desert that contains the alternative transmission line corridors is dominated by broad alluviated basins between relatively isolated mountain ranges and dissected uplands. The majority of the prominent mountain ranges in the project area trend north to northwest, and rise from less than 1,000 to as much as 4,000 feet above the surrounding lowlands. The average elevation of the study area (along the corridors) ranges from less than 1,500 feet in the central portion to over 3,000 feet in several places in the east-central and eastern portions. Local relief varies from extreme along the flanks of ranges in the central portion of the study area to very low in the alluvial valleys.

The entire study area is characterized by internal drainage, into the Mojave River, Kingston Wash, or one of many dry lakes or playas present in the region. The drainage characteristics of the project study area are discussed in more detail in the regional Surface Water Hydrology section.

Regional Geologic Setting

The alternative transmission line routes under consideration traverse a variety of geomorphic land forms. These landforms reflect, to a certain degree, the diverse nature of the underlying sediments and rocks that occur within the study area. The stratigraphy of the bedrock/sediment types traversed by the proposed routes is described below followed by a discussion of the geologic structure of the project area.

The geology of the proposed transmission line routes is described in the inventory table at the end of this report, and depicted on the geologic map on file in the Dames & Moore Project Files in Phoenix, Arizona.

Stratigraphy

Discussion of the stratigraphy of the study area is presented from oldest to youngest, beginning with a description of the consolidated rocks followed by descriptions of the relatively young unconsolidated to semi-consolidated sediments which underlie most of the alternative transmission line routes.

Consolidated Rocks

Consolidated rocks which outcrop within the transmission line corridors range in age from Precambrian to late Tertiary. Precambrian rocks are primarily metamorphic in origin and outcrop in several areas within the mapped corridors. These rocks are included with the Paleozoic and Mesozoic mapped units (PM series). Precambrian rocks exposed within the study area include gneiss, schist, quartzite and marble.

Relatively few rocks of Paleozoic age have been mapped in the study area. According to Dibblee (1980a), the Paleozoic sequence in the Mojave Desert consists of several thousand feet of slightly metamorphosed sedimentary rocks. These strata occur in the east-central portion of the study area. This sequence thins and becomes progressively more deformed, metamorphosed, and invaded by plutonic rocks toward the western part of the study area.

Rocks of Mesozoic age include marine and nonmarine metasedimentary rocks, metavolcanic rocks, and abundant granitic igneous rocks (Dibblee 1980a). A major portion of the Mojave Desert is floored by granitic intrusives similar in age and composition to the widespread granitic plutons present in the Sierra Nevada. Mesozoic sedimentary and metamorphic rocks are much less common in the study area.

Cenozoic-age strata are abundant in the study area and consist entirely of nonmarine sedimentary and volcanic rocks (Norris and Webb 1976). The sequence of volcanic and sedimentary deposits of early to middle Tertiary age is up to 10,000 feet thick in the study area (Dibblee 1980b). These strata consist primarily of intercalated volcanic, pyroclastic and clastic sedimentary

rocks which accumulated in the irregular eroded surface of the pre-Cenozoic basement rocks. Middle Tertiary strata include terrestrial clastic sediments as well as lacustrine clays (Dibblee 1980b). Late Tertiary rocks are relatively uncommon in the study area, except for Pliocene-age sediments exposed near Baker (Dibblee 1980b).

Quaternary Sediments and Volcanic Rocks

Most of the study area is underlain by semi-consolidated to unconsolidated Quaternary-age sediments. In most places on the geologic map, Quaternary alluvium has been differentiated as older alluvium, alluvium, and young alluvium. Other Quaternary units include dune sand and lacustrine (playa) deposits present beneath the dry lake beds. Older alluvium generally is associated with dissected alluvial fans or pediments while young alluvium has been mapped in the major washes or drainages including the channel of the Mojave River.

Quaternary volcanic rocks are present in the study area southeast of Baker, California (south of Link 10). No volcanic rocks were mapped within the alternative transmission line corridors.

Structure

The Mojave Desert is bounded on the northwest and southwest by the Garlock and San Andreas faults. These two major strike-slip faults bound a wedge-shaped structural block commonly referred to as the Mojave Block (Hewett 1954). Deformation within the block during the early to middle Tertiary formed several major roughly east-trending depositional basins in the western Mojave Block (Dibblee 1980c). These basins were the site of volcanic and sedimentary deposition during the early to middle Tertiary. During the late Tertiary, these basins were offset both vertically and laterally along northwest-trending faults in the western and central portions of the Mojave Block (Dibblee 1980c). In most cases, vertical offset appears to have preceded lateral displacements (Norris and Webb 1976). These faults trend roughly parallel to the San Andreas and at a high angle to the Garlock Fault. The more prominent of these include the Lenwood, Helendale, Camprock, Calico and Pishgah faults. Cenozoic sedimentary formations are deformed near these faults into folds that trend more westward than the faults, presumably due to right-lateral drag (Dibblee 1967). Although the northwest-trending faults within the Mojave Block partly bound ranges and valleys, they do not form impressive range fronts as do faults in the Basin and Range Physiographic Province. Vertical displacements on some of these faults show alternating in senses of apparent vertical separation along the fault, suggesting that movement has been predominantly lateral (Dibblee 1961).

Several east-west trending faults have been recognized in the central portion of the study area east and northeast of Barstow. The most prominent of these, the Manix Fault, displays evidence of left slip analogous to the Garlock Fault

to the north. This fault extends from the Calico Fault eastward (across Links 21 and 22) about 25 miles to the mouth of Afton Canyon (south of Link 17). These and other faults in the study area are discussed further in the section entitled Seismotectonic Setting.

The eastern portion of the study area contains little evidence of deformation in late Tertiary and Quaternary time. However, there is abundant evidence of Quaternary volcanic activity in the form of volcanic craters and cinder cones to the southeast of Baker and south of the alternative routes.

Paleontology

Most of the consolidated rock units exposed along the alternative transmission line routes are igneous or metamorphic in origin and are not fossiliferous. These include widespread occurrences of granitic igneous rocks and gneisses, slates, quartzites and phyllites of metamorphic origin.

Vertebrate mammalian fossils, including horse teeth, have been found in the Barstow Formation of middle Tertiary age in the central portion of the study area (Dibblee 1960). Localities in the Mud Hills have yielded one of the largest vertebrate faunas in North America. These strata are regarded as the type for the Barstovian Stage (Dibblee 1967). The richest fossil beds are generally found in the western portion of the Mud Hills, although some fossil remains are also found in the eastern Mud Hills near Owl, Solomon, and Rainbow canyons. Link 20b between Mileposts 3.0 and 5.3 may overlie such fossiliferous beds of the Barstow Formation. Another known fossil location within the Barstow Formation includes an area along Link 6 between Mileposts 79.7 and 82.0. Fossil mammal remains are also reported from the Barstow Formation in the vicinity of Daggett Ridge area (Dibblee 1970). Link 23 (Mileposts 11.0 to 12.2) and Link 25 (Mileposts 0.8 to 1.9) cross outcrops of the Barstow Formation in the Daggett Ridge area that may be fossiliferous.

Vertebrate mammalian fossils of late Pleistocene age occur within lacustrine deposits of ancient Lake Manix. These sediments are mapped along Link 18 (Mileposts 0.5 to 3.7), Link 19 (Mileposts 0.0 to 0.8), and Link 22 (Mileposts 0.0 to 2.0).

Invertebrate corals, brachiopods, and fusulinids are reported from the limestone of the Bird Hill limestone near Baker Hill. Link 9 between Mileposts 20.8 and 21.7 may cross portions of these sediments.

Minerals

A wide variety of minerals occur in the Mojave Desert. These can be divided into three groups: metallic minerals, nonmetallic minerals, and rare earth elements. For mineral locations and descriptions refer to Table 3-1.

Metallic Mineral Deposits

Metallic minerals deposits of the Mojave Desert include gold, silver, copper, lead, zinc, mercury, antimony and tungsten. These deposits were formed during the waning stages of magmatism, as metal-enriched fluids were expelled from a cooling magma and passed through fractures and other open cavities within outer margins of the plutons and surrounding country rock.

Nonmetallic Minerals

Nonmetallic mineral deposits within the study area include asbestos, barite, borates, zeolite, limestone and marl. Of these, the borate minerals are the most important. Large deposits of borate minerals occur in the Kramer District, located several miles northwest of Boron. The center of these operations is the large open pit mine operated by U.S. Borax and Chemical Corporation, located about 10 miles northwest of Link 38 in the northwestern portion of the study area. The East Kramer deposits are two smaller boratic deposits that occur 1.0 mile and 3.5 miles northwest of Kramer Junction (Link 38, Milepost 4.5). The borate deposits in the area occur within beds of an ancient playa (Dibblee 1967). Borates are used in large volumes by the glass and ceramics industries and by detergent manufacturers in Europe.

Rare Earth Elements

Rare earth elements found in the study area include cerium, lanthanum, neodymium, thorium and europium. These elements are found in Precambrian rocks of the Clark Mountains. This area is one of the major producers of these elements in the world.

Seismotectonic Setting

The seismotectonic setting of the project region is dominated by the San Andreas and Garlock faults. The San Andreas Fault extends from the Gulf of California northwestward to Cape Mendocino and forms the principal boundary between the Pacific and North American tectonic plates. The Garlock Fault is a major intraplate fault that extends eastward from its intersection with the San Andreas Fault and displays abundant evidence of left-lateral offset in Holocene time. Both of these faults are capable of generating major to great earthquakes which could result in significant levels of strong ground motion along the proposed transmission line routes.

In addition to these two major active structures, several lesser faults considered to be at least potentially active are present in the central and western portion of the project area. Most if not all of the faults listed in Table 3-2 are capable of generating moderate to large albeit less frequent events which, due to their closer proximity to the transmission line routes,

could result in significant ground motions as well as pose a potential surface fault rupture hazard.

Significant Faults

San Andreas Fault - The San Andreas Fault is seismically active and capable of generating earthquakes in excess of Magnitude 8. The distribution of seismic activity along the length of the fault is variable. Some segments have historically generated major or great earthquakes; other segments are characterized by more frequent, smaller magnitude events, and others are currently experiencing aseismic creep. At its closest approach to elements of the proposed project, the fault is about 17 miles southwest of the Adelanto and Victorville switching stations. This portion of the fault ruptured during the 1857 Magnitude ± 8 earthquake. Based on extensive studies by Kerry Sieh of the California Institute of Technology, this segment of the fault is inferred to generate major to great earthquakes with an average recurrence interval of between about 145 and 200 years (Sieh 1984).

Garlock Fault - The Garlock Fault extends from its junction with the San Andreas Fault eastward for approximately 150 miles. The fault is characterized by cumulative left slip of up to 30 to 40 miles (Dibblee 1980c) and is considered to be active based on abundant geomorphic evidence including offset stream channels. Seismic activity on the Garlock Fault could result in significant levels of ground motion in the western and central portions of the project area.

Northwest-Trending Faults - A number of northwest-trending faults have been mapped within the Mojave Block. Of these, the Helendale, Calico, Harper, Camprock, Lenwood, Spring, and South Lockhart faults cross one or more of the alternative transmission line links (Table 3-2). These faults are considered to be at least potentially active based on the ages of offset rock units. In addition, photo reconnaissance mapping along some of these faults by Morton et al. (1980) has identified numerous fault-related topographic features suggestive of late Quaternary activity. Dibblee (1961) presented evidence for right-lateral displacements ranging from hundreds of feet up to several miles on the Helendale and other faults. Most, if not all, of these faults may be capable of generating moderate to large magnitude earthquakes within the immediate project area.

Manix Fault - The Manix Fault is an east-west trending fault located about 25 miles east of Barstow, California. The east-west trend of the Manix Fault generally parallels that of the Garlock Fault to the north. The Manix Fault has been historically active and was the source of the 1947 Magnitude 6.2 Manix earthquake. About two inches of left-lateral displacement occurred along the Manix Fault during this event. South of Afton, the Manix Fault offsets Quaternary playa deposits about 10 feet in a vertical sense (Keaton 1977). This suggests that previous events may have contained an oblique or normal component of displacement. Where the western portion of the Manix Fault crosses Link 21 (Milepost 11.7) (Table 3-2), the surface trace of the fault

is readily apparent on aerial photographs. Here, the fault is marked by an abrupt south-facing scarp in older alluvium.

Historic Seismicity

The central and western portions of the project study area have experienced the effects of moderate to large earthquakes within historic times (post-1850). Earthquakes of Magnitude ± 8 in 1857 on the San Andreas Fault and in 1872 on the Sierra Nevada frontal fault system undoubtedly resulted in strong ground motion within the study area. Although there are no known faults within the Mojave Block capable of generating earthquakes of this size, moderate to large ($\leq$ Magnitude 7) events could potentially occur along faults within the block. The two most significant events that have occurred within the study area are the 1947 Manix and 1975 Galway Lake earthquakes. These events are discussed briefly below.

1947 Manix Earthquake - On April 10, 1947, a Magnitude 6.2 earthquake occurred on the Manix Fault east of Barstow, California. The earthquake was accompanied by two inches of left-lateral displacement for a distance of about two miles along the Manix Fault (Buwalda and Richter 1948). The earthquake was felt throughout the study area and as far away as San Diego, Bakersfield, Phoenix, and Las Vegas. Five minutes after the main event an after-shock measuring Magnitude 5.2 occurred followed by about 140 smaller magnitude after-shocks (Keaton 1977).

1975 Galway Lake Earthquake - On May 31, 1975, a Magnitude 5.2 earthquake occurred on a previously unmapped fault in the Galway Lake area south of the study area. The earthquake was accompanied by surface rupture in the form of left stepping en echelon cracks indicative of right-lateral displacement. Surface rupture occurred along an approximately four-mile-long fault segment, which is now referred to as the Galway Lake Fault. Right-lateral displacements ranging between 2 and 5 millimeters were measured along this segment (Beeby and Hill 1975). The Galway Lake Fault trends north to northwest and is at a slightly oblique angle to the more northwesterly trending Emerson Fault which appears to be related to the Camprock Fault to the northwest.

Hydrology

Surface Water Hydrology

The study area is characterized by internal drainage, into either the Mojave River or Kingston Wash, or into one of many dry lakes (playas) present in the region. Kingston Wash, in the east-central portion of the study area, drains northward into the Amargosa River which eventually terminates in Death Valley. The Mojave River originates in the San Bernardino Mountains southwest of the study area and flows generally northward through Victorville and then eastward through Barstow. The Mojave River eventually terminates

at Soda Lake (playa) in the central portion of the study area. Most of the Mojave River is characterized by subsurface flow; surface flow generally occurs only in areas where the channel is restricted such as at Lower Narrows near Victorville (Link 33, Milepost 14.6) and in Afton Canyon (south of Link 17). Peak, mean and minimum daily discharges in the Mojave River at these two localities for the years 1960-1978 are listed in Tables 3-3 and 3-4.

Due to the arid climate, all of the drainages within the study area are ephemeral except for portions of the Mojave River as discussed above. Streamflow occurs only in direct response to rainfall, although generally only a small fraction of the total precipitation results in surface runoff due to evapotranspiration and infiltration. According to Maxey (1968), recharge in typical arid basins occurs primarily as a result of infiltration of surface runoff from precipitation over the adjacent upland areas. Occasionally, intense storms generate enough precipitation to result in surface flow that reaches the local base level which, in many cases in the project area, is a dry lake bed. Storms resulting in inundation of the playa surfaces occur only infrequently, perhaps several times per century. The potential for flooding is discussed further in the section entitled Geologic Hazards.

Ground Water Hydrology

The Mojave Desert region is generally characterized by large ground water basins which are commonly interconnected resulting in interbasinal flow of ground water. In general, shallow ground water is present in the topographically lowest portion of the hydrologic basins which, in the study area, is visually a playa. However, shallow ground water beneath playa lakes is commonly of poor quality due to salt buildup resulting in high levels of total dissolved solids. Shallow ground water also is common beneath the distal portions of alluvial fan complexes (Maxey 1968).

Generally, the most productive aquifers in arid hydrologic basins consist of coarser alluvial deposits as well as conglomerates beneath the flanks of the basin. Consolidated rock units generally are not found to be productive aquifers unless highly fractured. Fine-grained deposits commonly present in the central portions of desert basins also generally yield only small amounts of water to wells.

Soils

Due to the large size and remoteness of the study area, detailed soil surveys have not been completed for most of the study area. Data are available covering the extreme western and eastern portions of the study area (USDA, unpublished data; HDR 1981). These data were compared to landforms and mapped geologic units within the remainder of the study area to provide generalized descriptions of the soils inferred to be present within the transmission line corridors.

The primary factors that have influenced soil development in the study area are climate, geology and topography. Low annual precipitation restricts vegetation cover, resulting in limited humus accumulation, and light (soil) colors (HDR 1981). Subsurface accumulation of calcium carbonate is common and often forms a caliche or K horizon. Soils formed on well-drained alluvium may also contain loamy horizons of clay accumulation. Precipitation in the area occurs during short periods of high intensity rainfall. These intensive storms typically result in high runoff from mountainous areas on the alluvial fans, and flooding of alluvial plains and playas. This periodical flooding and evaporation of the floodwaters causes an accumulation of salts, which greatly affect soil development on the alluvial plains and playas.

Topography influences soil formation primarily through its effects upon drainage, runoff and erosion, and secondarily through variations in exposure to the sun and wind (USDA 1951). Topographic features within the study area include mountain ranges, alluvial fans and terraces, and alluvial plains and playas. The steep slopes of the mountainous regions experience rapid to very rapid runoff and commonly high rates of erosion. Soil development is generally stunted because of rapid erosion as well as indurated parent material (bedrock). Lithosols and the lithosolic associations of other soils are characteristic of the steep slopes in mountain areas (USDA 1951). Moderate to slightly sloping alluvial fans and terraces experience medium to slow runoff. These areas are characterized by moderate to slow erosion rates, and may develop deep soils. Planosols are typically found in these areas. Alluvial plains and playas experience very slow to no runoff, and generally no natural erosion. These areas retain water that falls as rain and receive a considerable amount of runoff from adjacent uplands. Hydromorphic and halomorphie intrazonal soils are typically found in these areas (USDA 1951). In many places wind has removed the fine soil particles from old alluvium, creating a surface of "desert pavement." The resulting surface of desert pavement is resistant to water and wind erosion.

The study area contains soils belonging predominantly to the Aridisol and Entisol USDA taxonomic soil orders. Aridisols are characterized by a low content of organic matter in the A horizon and salt or silica accumulations at depth. The two suborders of Aridisols, Argids and Orthids, are based on the presence or absence of an argillic (clay-rich) B horizon (Birkeland 1974). Aridisols are dry more than six months of the year in all horizons. In the study area Aridisols have developed on Quaternary older alluvium, and Quaternary alluvium. Argids dominate the Quaternary older alluvium.

Entisols are young soils that are characteristically lacking developed subsurface horizons. The two suborders of Entisols are Orthents (loamy or clayey) and Psamments (textures of loamy fine sand or coarser). In the study area, Entisols have developed on Quaternary alluvium and young alluvium. Orthents dominate the playas and Psamments occur on young alluvial fans, plains and terraces.

A discussion of the potential for soil erosion is presented in the following section.

Geologic Hazards

Soil Erosion

Soils within the alternative study corridors generally coincide with the following sediment types and/or geomorphic surfaces: young alluvium in active stream channels and floodplains, eolian sand deposits, fine-grained playa deposits, undissected alluvial fans and plains, older (dissected) alluvial fans, and consolidated rocks.

The potential for soil erosion is generally low within most of the study area due to the relatively low amount of rainfall, the sediment/soil types present, and the predominantly gentle slopes along the proposed transmission line routes. However, the potential for accelerated soil erosion due to transmission line construction is significant for some of these soil types. Water is generally the most significant erosive agent although wind plays an important role in arid environments such as the study area.

As shown on Figure 4-1 (Map Volume), the following study area soils have the highest potential to experience accelerated erosion due to project activities: young alluvium in active stream channels and floodplains, and areas with eolian sand deposits (sand dunes). The potential for soil erosion in areas underlain by consolidated rocks with little or no soil development is considered to be low to nonexistent. The potential for soil erosion is considered to be moderate to low in all other areas along the proposed transmission line routes. Areas with high potential for soil erosion are noted in the inventory table for geology/hydrology.

Surface Fault Displacement

The potential for surface faulting along the proposed transmission line route exists at locations where mapped quaternary faults cross the proposed routes. The locations of these crossings are listed in Table 3-2 and are noted in the inventory table. Quaternary faults which cross proposed routes are discussed above in the section entitled Seismotectonic Setting. The frequency with which surface fault rupture occurs along these faults is not known. However, based on evidence such as displaced contacts, geomorphic expression, tectonic setting, and historical seismicity, it is likely that earthquake recurrence intervals on these faults is on the order of hundreds to thousands or even tens of thousands of years. Therefore, the potential for surface fault rupture to occur at any of the fault crossings during the lifetime of the project is considered to be low to extremely low.

Strong Ground Motion

Due to the proximity of the project area to major active structures such as the San Andreas and Garlock faults and the presence of Quaternary faults in the immediate project area, the alternative transmission line routes will likely be

subjected to significant levels of strong ground motion as a result of seismic activity within the project lifetime. The level of acceleration that can be expected at a given location depends on several factors including distance to the earthquake source, magnitude of the event, subsurface soil conditions, and attenuation of ground motion with distance from the source. Due to the highly variable nature of these factors, levels of acceleration that will result from a given seismic event will vary considerable along the proposed routes. However, based on the distribution of seismogenic faults within and adjacent to the study area, it is apparent that both the likelihood and strength of seismically induced strong ground motions increases from east to west across the study area.

Soil Liquefaction

In order for liquefaction to occur, loose, granular, saturated soils in the shallow subsurface must be subjected to repeated cycles of strong ground motion. The potential for liquefaction is present in several areas in the western and central portions of the study area where shallow groundwater underlies granular soils within and adjacent to the Mojave River. These areas are noted in the inventory table. Although shallow groundwater is likely present in the vicinity of the playa lakes, these areas are commonly underlain by silts and clays which would not be subject to liquefaction.

Flooding and Debris Flows

As discussed in the section entitled Surface Water Hydrology, surface water flow generally occurs only in direct response to rainfall from infrequent but intense storms. Such storms commonly trigger flash flooding in channelized drainages that receive runoff from large areas with high levels of precipitation. Drainages subject to flash flooding are most common in dissected alluvial fans adjacent to mountainous terrain. Debris flows most commonly occur on active alluvial fans (Map Units Qya and Qal, Figure 4-1) in response to episodes of intense precipitation. The potential for debris flows is higher in areas downslope from exposures of unconsolidated to semi-consolidated sediments. These areas are noted in the inventory table.

Slope Instability

No areas with potential serious slope instability problems such as major landslide terrains have been noted in the study area. Most of the alternative transmission line routes traverse areas with slopes less than five percent. Areas with the highest potential for slope instability consist mainly of exposures of weathered bedrock with relatively steep (>10 to 15 percent) slopes. These areas are noted in the inventory table.

Hydrocompaction (Soil Collapse)

Sediments susceptible to hydrocompaction are loose, low density, and open-textured soils which collapse upon wetting. Soils subject to hydrocompaction are typically found in young alluvial fan deposits above the zone of saturation, generally within a few hundred feet of the ground surface. Subsidence due to hydrocompaction could present a potential hazard to transmission line towers if ponding of water occurs beneath a tower footing underlain by susceptible soils. Subsidence related to hydrocompaction would not present a problem if its occurrence is restricted to areas between towers.

The potential for subsidence due to hydrocompaction is highest in the young alluvial fan deposits. The effects of hydrocompaction would only present a hazard to the transmission route if the soils underlying a support tower were affected. The potential for soil collapse would be evaluated at the location of transmission towers in high susceptibility areas prior to final engineering design and construction, and mitigative measures would be taken as appropriate.

Expansive Soils

In an arid environment any soil that contains expansive clay minerals will exhibit some shrink-swell potential. The amount of expansion-shrinkage that results is a function of confining stress and moisture content changes as well as percentage of the soil that is composed of expansive clay minerals. Soils containing expansive clay minerals such as montmorillonite are likely present in the vicinity of the playa lake beds as well as the distal portions of alluvial fans.

Expansiveness of soils is usually identified based on soil index tests such as dry density and liquid limit and by consolidation tests where actual soil expansion is measured in a consolidimeter after water flooding. Such tests should be performed on soil samples prior to final design of the transmission line towers.

IMPACT ASSESSMENT/MITIGATION PLANNING PROCESS

The impact assessment for Earth Resources was designed to determine which features, if any, may potentially be impacted by transmission line construction and/or operation. Of those features inventoried, only current soil erosion patterns and paleontological resources could be shown to potentially be significantly affected by the proposed transmission system. In addition, an analysis of geologic hazards was conducted to determine those hazards that could potentially affect the proposed project.

Based on the results of the impact assessment, mitigation recommendations were made to potentially reduce, where feasible, identified significant impact levels. The overall objective in mitigation planning is to create low residual impacts.

Impact Assessment Methods

The method for the impact assessment was derived by initially defining impact levels with respect to the impacted resources, namely, soil erosion potential and paleontological resources. Four qualitative terms or combinations thereof are used in this assessment to describe the magnitude of potential impacts: high, moderate, low and negligible. These impact levels are derived by applying a set of specific criteria (i.e., sensitivity, quality, and quantity and duration).

Soil Erosion Potential

Sensitivity--represents both the inherent value of the resource as well as its susceptibility to changes brought about by this project. With respect to soil erosion, sensitivity is defined in the inventory report as the qualitative estimate of erosion potential (i.e., high, moderate to low, and low).

The soils were classified on the basis of diagnostic morphologic characteristics. In addition to classifying each soil on the basis of diagnostic morphology, broad soil/geomorphic relationships were used to determine erosion potential (Figure 4-1).

Quality and Quantity--Quality represents the existing condition (i.e., disturbed or undisturbed) of the resource. Quantity represents the amount of area disturbed by the project. With respect to erosion, quality and quantity were defined in terms of access levels or amount of disturbance resulting from proposed tower pads, access roads, pulling stations, and splicing sites. The three access levels are defined below:

- Sufficient (existing access is sufficient)

Existing roads, including paved, improved, or those currently being used for utility right-of-way access will be fully utilized, with no blading or

substantive upgrading required. Approximately 0.25 mile of new road (spur road) construction would be required for every mile of transmission line.

- Upgrade (existing access requires upgrading)

Existing roads will be utilized but will require substantive upgrading (blading, culvert installation, etc.). Approximately 0.75 mile of new road (spur and centerline) construction would be required for every mile of transmission line.

- New (new access is required)

Existing roads are insufficient or nonexistent, requiring the construction of totally new access roads. Approximately 1.5 to 2.0 miles of new road (spur and centerline) construction would be required for every mile of transmission line.

Duration--represents the time during which the resource would be affected by the proposed project, either short- or long-term. For the purpose of this assessment, all impacts were assumed to be long-term.

Once the impact criteria were formulated, the impact model was used to predict the various impact levels. Table 3-8 presents the soil erosion impact assessment model with potential impact levels.

Paleontological Resources

The four impact criteria as they apply to paleontological resources are defined below:

Sensitivity--represents the value or significance of the fossil-bearing strata. In this case, only those areas underlain by fossil-bearing strata with a high or moderate significance level were evaluated with the impact assessment model.

Quality and Quantity--represent the access level or amount of disturbance resulting from the proposed tower pads, access roads, pulling stations, and splicing sites. The three access levels (sufficient, upgrade and new) are defined the same as outlined under soil erosion potential.

Duration--represents the time during which the resource would be affected by the proposed project, which in this case, all impacts are considered long-term.

Once the impact criteria were formulated, the impact model was used to predict the various impact levels. Table 3-9 presents the paleontological resources impact assessment model with potential impact levels.

Potential Geotechnical Hazards

Results of the inventory portion of this study have shown that some aspects of the geologic environment could affect the proposed steel lattice towers and attached conductors; these are identified as geotechnical hazards to the proposed project.

The purpose of this study was to combine the geotechnical data collected during the inventory portion of this study so that the cumulative effects of the different hazards would be easily recognized.

The nature and extent of the geotechnical hazards considered to be of potential significance to the proposed project are discussed in detail in the baseline inventory report. The following geotechnical features are considered in the hazard analysis:

- Surface Fault Displacement
- Strong Ground Motion
- Soil Liquefaction
- Slope Instability
- Expansive Soils
- Hydrocompaction (Soil Collapse)
- Flooding and Debris Flows

The purpose of this report was to summarize the potential hazards presented by these features. In this regard, the hazards could potentially present either siting or foundation problems, depending on the nature and extent of the hazard. (A siting problem is defined as a geotechnical hazard that could be avoided by proper route selection and/or tower spacing within the proposed corridors. A foundation problem is defined as a geotechnical hazard that could require special tower foundation design which would be developed during the specific tower foundation investigation.)

Surface Fault Displacement

Owing to the difficulty in designing foundations to accommodate differential movement along faults, active or potentially active fault zones are considered a siting problem. Faults are generally long, linear features with limited width; therefore, most fault traces can be avoided by properly routing and spacing transmission towers within the corridors. Most of the faults traversed by the corridors have dominant right-lateral movement; if this is considered when designing the crossings, the routes can be aligned so that movement will not overextend the transmission lines, causing rupture.

The locations of fault crossings are listed in Table 3-2. Link 22, between Mileposts 12.0 and 14.5, closely parallels the Manix Fault, the trace of which is concealed beneath alluvial deposits. It is recommended that this portion of Link 22 be slightly realigned such that the proposed route crosses the fault at a higher angle.

Strong Ground Motion

Most of the region traversed by the corridors is likely to be subjected to moderate to strong levels of ground motion during seismic events. However, transmission towers are particularly insensitive to primary ground shaking during seismic events. Therefore, no specific design measures to mitigate the effects of strong ground motion are recommended.

Soil Liquefaction

Soils that liquefy during an earthquake may undergo permanent differential displacement and such movements can present significant hazards to towers. Therefore, liquefaction is considered a foundation problem. The potential for liquefaction is present in several areas within the western and central portion of the study area where shallow groundwater may be present along with granular soils. These areas are noted on the inventory table for earth resources. Soil liquefaction potential can be mitigated by designing tower foundations to withstand the effects of liquefaction.

Slope Instability

Potential slope instability conditions are present in only a few areas within the proposed corridors. These areas were noted in the inventory table. Slope instability is considered to be a siting problem due to its localized nature and can be mitigated through selective siting of towers in areas where this problem may exist.

Expansive Soils

Expansive soils may undergo volumetric changes that can produce significant vertical movement. Because of the potential for vertical movements, expansive soils are considered to be a foundation problem. They also are so extensive locally that, in general, they cannot be routed around and are not considered a siting problem.

Hydrocompaction (Soil Collapse)

The potential for hydrocompaction occurs where loose, low density, and open-textured soils may be susceptible to collapse upon wetting. Hydrocompaction may be a potential hazard in areas underlain by young alluvial fan deposits. The potential for soil collapse should be evaluated at the transmission tower locations in high susceptibility areas prior to final engineering design and construction so that mitigative measures can be taken as appropriate.

Flooding and Debris Flow Potential

Flash floods are generally localized and are considered to be a siting problem. Proper route alignment and tower spacing should alleviate this concern.

In areas where gulleys are of substantial width and placement cannot be avoided (i.e., width greater than tower spacing), flash floods could be considered to present foundation problems; however, proper engineering measures can be used to protect tower sites. Areas where flooding and debris flows could potentially occur are noted on the inventory table. Flooding and/or inundation could occur within 100-year floodplains which coincide with young alluvium and playa deposits which have been mapped as part of the baseline inventory (Qya and QI map units on Figure 4-1).

Mitigation Planning Methods

The rationale used to develop specific mitigation recommendations is based on potential impacts as predicted in each impact model as well as existing conditions along the alternative corridors. The mitigation recommendations presented herein are designed with realistic constraints in mind (i.e., time, cost, construction). These recommended mitigation measures represent a reasonable attempt to reduce the amount of disturbed area to the greatest extent possible.

Taking into account the generically committed mitigation measures, the following specific mitigation measures are recommended for areas with potentially significant impact levels to soil erosion and paleontological resources. These areas are identified on the Impact Assessment/Mitigation Planning Charts (Tables 3-10 and 3-11).

12. Prior to construction, a geotechnical field review of tower and access road design will be conducted by an appropriately qualified professional to identify site-specific, soil erosion impacts and determine the most effective means of mitigating them. Possible mitigation measures could include minor adjustments in tower and road locations, restricting access during periods of high moisture and utilizing selective biodegradable soil stabilizing agents, etc.
- 12A. Prior to construction, a field review of tower and access road design and location will be conducted by a qualified professional to identify fossil localities that should not be disturbed. Minor realignment of tower and access road locations may be necessary.

Results

The potential for soil erosion is illustrated on Figure 4-1 and the results of impact assessment and mitigation planning analysis are presented on

Table 3-10. The results of the impact assessment and mitigation planning for paleontological resources are presented on Table 3-11.

While portions of some transmission line routes may be affected by potential geological hazards, it should be noted that with minor exception, the potential hazards can be mitigated by proper route selection, tower spacing, or foundation design.

AGENCY CONTACTS

EARTH RESOURCES (GEOLOGY/HYDROLOGY/SOILS)

Agency	Date	Type of Contact	Summary
United States Department of Agriculture, Soil Conservation Service (Apple Valley)			
Jim Earson	11/20/84	Meeting	Obtained unpublished soil survey material (Victorville to Barstow area only).
USDA, BLM			
Riverside Bill Collins	11/21/84	Meeting	Obtained literature on geology and mineral resources of the Mojave Desert.
San Bernardino County Geologist			
Lou Blanck	11/20 - 11/21/84	Meeting	Obtained general geology and slope stability maps for the Mojave Desert. Requested references for maps in study area.
California Department of Transportation			
Bob Austin	11/29/84	Telephone	Requested any information dealing with geohazards along I-15. Was told that CALTRANS uses railroad system and designs for 100-year flood.

Agency Contacts (continued)
Earth Resources (Geology/Hydrology/Soils)

<u>Agency</u>	<u>Date</u>	<u>Type of Contact</u>	<u>Summary</u>
California Department of Transportation			
Tim Griffin (District Drainage Engineer)	11/24/84	Telephone	Requested any material or data used in surface runoff calculations for I-15. Will be sent.
California Department of Transportation			
Bob Hott	12/16/84	Telephone	Requested specific information regarding magnitude of 100-year flood calculations used in design of I-15. He will look for and call back.
Nevada Bureau of Mines and Geology			
John Bell	11/20/84	Telephone	Received information on geologic maps covering the portions of transmission line corridors within Nevada. Mr. Bell also informed us that flood maps are not available for these areas.
Thomas W. Dibblee, Jr. (Retired Consulting Geologist)	11/20/84	Meeting	Mr. Dibblee referred us to published geologic maps which cover the entire project area southwest of the Baker area. In addition, Mr. Dibblee agreed to provide us with a complete and unpublished geologic map of the Cave Mountain 15-foot quadrangle.

REFERENCES

- Anderson, R.E. 1977. Geologic map of the Boulder City 15-Minute Quadrangle, Clark County Nevada: United States Geological Survey Map GQ-1395.
- Beeby, D.J. and R.L. Hill. 1975. Galway Lake Fault. California Geology, Volume 28, Number 10, pp. 219-221.
- Birkeland, P.W. 1974. Pedology, Weathering, and Geomorphological Research. Oxford Press, New York.
- Bowen, O.E., Jr. 1954. Geology and Mineral Deposits of Barstow Quadrangle, San Bernardino County, California: California Division Mines Bulletin 165, p. 7-185, map, scale 1:125,000.
- Buwalda, J.P. and C.F. Richter. 1948. Movement of the Manix (California) fault on April 10, 1947. Abstract: Geological Society of America, Bulletin v. 59, no. 12, p. 1367.
- Byers, F.M. 1963. Geology of the Alvord Mountain Quadrangle, San Bernardino County, California: U.S. Geol. Survey Bull. 1089a, 71 pp., map scale 1:62,500.
- Clary, M.R. 1976. Geology of the eastern part of the Clark Mountain Range, San Bernardino County, California. California Division of Mines and Geology, Map Sheet 6.
- Davis, J.F. and T.P. Anderson. 1980. Mineral resources of the California Desert. In Geology and Mineral Wealth of the Mojave Desert. p. 122.
- Dibblee, T.W., Jr. 1980a. Pre Cenozoic rock units of the Mojave Desert. In Geology and Mineral Wealth of the California Desert. p. 13.
- . 1980b. Cenozoic rock units of the Mojave Desert. In Geology and Mineral Wealth of the Mojave Desert. p. 41.
- . 1980c. Structure of the Mojave Desert. In Geology and Mineral Wealth of the Mojave Desert. p. 69.
- . 1970. Geologic Map of the Daggett Quadrangle, San Bernardino County, California: U.S. Geol. Survey Misc. Geol. Inv. Map I-592, scale 1:62,500.
- . 1968. Geology of the Fremont Peak and Opal Mountain Quadrangles, California: Calif. Div. Mines and Geology Bull. 188, 64 pp., map scale 1:62,500.
- . 1967a. Areal Geology of the Western Mojave Desert, California: United States Geol. Survey Prof. Paper 522, map scale 1:125,000. 153 pp.

TABLE 3-1
MINES AND/OR MINERAL DEPOSITS
IN PROJECT STUDY AREA*

Loc. Number	Description	Minerals/Commodity Present
1	Multiple claims	Cold, Silver, Copper
2		Gold, Silver
3		Gold
4		Iron, Manganese
5		Gold, Manganese
6	Delany Mine	Fe
7	Alford Mine	Gold
8	Shaff	
9	Multiple claims	
10		Copper
11	Multiple claims	Talc and Talcum
12		Talc, Copper
13		Gold
14		Gold
15		Copper
16	McIntosh Mine	Lead, Silver, Zinc
17		Lead, Silver, Zinc
18	Sulfide Canyon Mine	Gold, Silver
19	Partridge Mine	Rare earth elements
20	Timbercliff Mine	
21	State Line Mine	
22	Shire Gypsum Deposit	
23	Alderson Mine	Lead, Silver
24	Multiple claims	
25	Telegraph Mine	Gold, Silver
26	Landing Field Mine	
27	Pine Lake Mine	Montanite
Area 1		High potential for oil and gas
Area 2		High potential for oil and gas
Area 3		Rare earth elements
Area 4		Rare earth elements
Area 5		Gold, Silver
Area 6		Gold, Silver
Area 7		Gold, Silver

TABLES

TABLE 3-1
MINES AND/OR MINERAL DEPOSITS
IN PROJECT STUDY AREA*

Link Number	Description	Minerals/Commodity Present
1	Multiple claims	Gold, Silver, Copper
2		Gold, Silver
3		Gold
4		Gold, Barite
5		Gold, Barite
6	Daisy Mine	(?)
7	Alvord Mine	Gold
8	Shaft	
9	Multiple claims	
10		Copper
11		Talc and Tungsten
12		Talc, Copper
13		Gold
14	Mohawk Mine	Gold
15		Copper
16		Lead, Silver, Zinc
17		Lead, Silver, Zinc
18		Gold, Silver
19	Sulfide Queen Mine	Rare earth elements
20	Birthday Mine	
21	Umberci Mine	
22	State Line Mine	
23	Shire Gypsum Deposit	
24	Atkinson Mine	Lead, Silver
25	Multiple mines	
26	Telegraph Mine	Gold, Silver
27	Landing Field Mine	
27	Pink Lady Mine	Bentonite
Area 1		High potential for oil and gas
Area 2		High potential for oil and gas
Area 3		Rare earth elements
Area 4		Rare earth elements
Area 5		Gold, Silver, Zinc
Area 6		Gold, Silver, Zinc
Area 7		Gold, Silver, Zinc

*Includes mines/mineral deposits within a six-mile study corridor.

TABLE 3-2
 ROUTES CROSSED BY QUATERNARY FAULTS

<u>Fault</u>	<u>Link</u>	<u>Milepost</u>
State Line	6	2.0
State Line	5b	3.6
State Line	45	4.2
Calico	22	21.1
Calico	23	0.0
Calico	21b	8.0
Camprock	23	9.8
Harper	20	32.1
Helendale	24a	21.5
Helendale	32	4.8
Helendale	33	5.0
Helendale	27	5.8
Lenwood	25	0.5
Lenwood	24a	0.6
Lenwood	25	0.7
Lenwood	24a	1.1
Manix	22	13.5
Manix	22	8.8
Manix	21b	6.1
South Lockhart	37	0.2
South Lockhart	38	0.3
Spring	42	19.8
Spring	39	2.8
Unnamed	41	0.2
Unnamed	42	0.2
Unnamed	42	16.5
Unnamed	42	16.9
Unnamed	6	66.0

Note: Fault names as used by Jennings (1975).

TABLE 3-3
DAILY DISCHARGES OF MOJAVE RIVER - LOWER NARROWS

<u>Water-Year</u>		<u>Daily Discharge Rate (cubic feet/second)</u>		
		<u>Mean</u>	<u>Peak</u>	<u>Minimum</u>
1960-1961		26.1	284	8.7
1	2	37.0	1,760	8.7
2	3	23.5	242	11.0
3	4	23.6	135	8.7
4	5	23.2	100	8.1
5	6	70.5	5,060	6.7
6	7	103.0	6,860	10.0
7	8	25.9	54	8.7
8	9	402.0	21,000	14.0
1969-1970		31.9	102	9.6
0	1	28.3	81	8.4
1	2	31.4	1,540	10.0
2	3	48.0	1,420	7.9
3	4	24.5	70	7.9
4	5	23.0	48	3.4
5	6	27.8	200	11.0
6	7	39.0	128	14.0
7	8	289.0	9,700	15.0

TABLE 3-4
DAILY DISCHARGES OF MOJAVE RIVER - AFTON CANYON

<u>Water-Year</u>	<u>Discharge Rate (cubic feet/second)</u>		
	<u>Mean Daily Discharge</u>	<u>Peak Daily Discharge</u>	<u>Minimum Daily Discharge</u>
1960-1961	0.92	20.0	0.0
1 2	0.78	4.6	0.0
2 3	1.04	61.0	0.0
3 4	0.74	3.6	0.0
4 5	0.78	9.1	0.0
5 6	6.60	1.170	0.0
6 7	2.02	403.0	0.0
7 8	0.49	13.0	0.0
8 9	100.00	9,500	0.0
9 1970	0.75	3.2	0.0
1970-1971	0.50	1.6	0.0
1 2	0.82	139.0	0.0
2 3	0.43	3.9	0.0
3 4	0.60	33.0	0.0
4 5	0.22	1.0	0.0
5 6	0.41	79.0	0.0
6 7	1.24	156.0	0.0
7 8	64.6	5,870	0.14

TABLE 3-5
INVENTORY OF GEOLOGIC RESOURCES

Link No.	Length	Mileposts		Description
		Begin	End	
0	13.5	0.0	13.5	Sediment/ Rock Types: Alluvium adjacent to Eldorado Dry Lake Bed. Geologic Hazards: Corridor crosses wash at Mile-post 2.5. Possible shallow groundwater near Eldorado Dry Lake.
1	16.6	0.0	7.2	Sediment/ Rock Types: Alluvium Geologic Hazards: None
		7.2	10.6	Sediment/ Rock Types: Tertiary volcanic and Precambrian (metamorphic?) rocks. Geologic Hazards: Slope instability.
		10.6	16.6	Sediment/ Rock Types: Alluvium with outcrops of Precambrian and Paleozoic sedimentary and metamorphic rocks. Geologic Hazards: None
2	6.5	0.0	4.5	Sediment/ Rock Types: Alluvium and playa deposits of Roach Lake (from Mile-posts 2.5 to 3.0). Geologic Hazards: Possible inundation along edge of Roach Lake from infrequent intense storms; shallow groundwater.

Table 3-5 (continued)
Inventory of Geologic Resources

Link No.	Length	Mileposts		Description
		Begin	End	
3	8.4	0.0	8.4	Sediment/ Rock Types: Alluvium and playa deposits of Roach Lake (from Mileposts 2.0 to 4.8). Geologic Hazards: Possible inundation of Roach Lake following infrequent intense storms; shallow groundwater.
4	1.9	0.0	1.9	Sediment/ Rock Types: Alluvium. Geologic Hazards: Shallow groundwater.
5a	0.2	0.0	0.2	Sediment/ Rock Types: Alluvium. Geologic Hazards: None.
5b	4.8	0.0	4.8	Sediment/ Rock Types: Alluvium, playa deposits of Ivanpah Lake. Geologic Hazards: Possible inundation of playa surface following infrequent intense storms; cross State Line Fault at Milepost 3.6.
5c	12.3	0.0	6.8	Sediment/ Rock Types: Alluvium, playa deposits of Ivanpah Lake. Geologic Hazards: Inundation of playa surface following intense storms.

Table 3-5 (continued)
Inventory of Geologic Resources

Link No.	Length	Mileposts		Description
		Begin	End	
5c cont.		6.8	12.3	Sediment Rock Types: Pre-Tertiary sedimentary and metamorphic bedrock units, older alluvium. Geologic Hazards: Unstable slopes in bedrock areas, flash flooding in narrow drainage courses.
6	83.3	0	10.1	Sediment/ Rock Types: Alluvium and playa deposits (Mileposts 0.0 - 0.5). Geologic Hazards: Occasional gullies which may be susceptible to flash floods; State Line Fault crosses route at Milepost 2.0.
		10.1	12.0	Sediment/ Rock Types: Precambrian granitic gneiss. Geologic Hazards: Slope stability. High potential for soil erosion in drainages between Mileposts 9.0 and 13.5.
		12.0	29.4	Sediment/ Rock Types: Alluvium and possibly (undifferentiated) older alluvium. Geologic Hazards: High potential for soil erosion in drainages between Mileposts 28.0 and 30.2.
		29.4	31.2	Sediment/ Rock Types: Late Tertiary fanglomerate. Geologic Hazards: None noted.

Table 3-5 (continued)
Inventory of Geologic Resources

Link No.	Length	Mileposts		Description
		Begin	End	
6 cont.		31.2	42.0	Sediment/ Rock Types: Undifferentiated alluvium and older alluvium with some isolated areas of intrusive igneous bedrock.
				Geologic Hazards: Debris flow hazard between Mileposts 35.5 and 39.0.
		42.0	50.0	Sediment/ Rock Types: Alluvium and several small areas of metamorphic or intrusive igneous bedrock.
				Geologic Hazards: Shallow groundwater near Silver Lake; slope instability (near Milepost 48.8); debris flow potential (Mileposts 42.0 to 44.5, 47.5 to 50.0); flooding.
		50.0	63.8	Sediment/ Rock Types: Alluvium with some areas of undifferentiated older alluvium.
				Geologic Hazards: Gullies between Mileposts 56.0 and 59.0 susceptible to flash floods.
		63.8	73.5	Sediment/ Rock Types: Alluvium with intrusive igneous bedrock (Mileposts 63.8 to 64.6) and undifferentiated areas of older alluvium.
				Geologic Hazards: Possible Quaternary fault crossing (Milepost 66.0). High potential for soil erosion in drainages near Mileposts 68.8, 71.7 and 72.0.

Table 3-5 (continued)
Inventory of Geologic Resources

Link No.	Length	Mileposts		Description
		Begin	End	
6 cont.		73.5	83.3	Sediment/ Rock Types: Highly variable, includes alluvium, dissected older alluvium, undifferentiated Tertiary sedimentary rocks and isolated dune sand deposits (Mileposts 82.0 to 83.0). Geologic Hazards: Potential flash floods in gullies, especially in dissected areas southeast of Alford Mountain (Mileposts 79.0 to 83.5); potential debris flow hazard (Mileposts 78.0 to 82.0); active sand dunes.
7	13.5	0.0	13.5	Sediment/ Rock Types: Alluvium, older alluvium, Tertiary sedimentary rocks (Mileposts 12.2 to 13.3) and Mesozoic igneous rocks (Mileposts 8.7 to 9.6). Geologic Hazards: Flooding in gullies/drainages following intense storms; potential debris flows (Mileposts 0.0 to 4.5, 7.4 to 8.8).
8	12.9	0.0	12.9	Sediment/ Rock Types: Alluvium, older alluvium, older lake deposits. Geologic Hazards: Flash flood potential, debris flows (Mileposts 0.0 to 3.5).
9	25.0	0.0	7.0	Sediment/ Rock Types: Alluvium and Mesozoic granitic rocks. Geologic Hazards: Strong potential for flash flooding in Halloran Wash (Mileposts 1.8 to 7.0).

Table 3-5 (continued)
Inventory of Geologic Resources

Link No.	Length	Mileposts		Description
		Begin	End	
9 cont.		7.0	25.0	Sediment/ Rock Types: Alluvium, Tertiary sedimentary rocks (Mileposts 23.5 to 25.0). Geologic Hazards: Possible shallow groundwater near Baker (Mileposts 18.5 to 20.0).
10	19.9	0.0	19.9	Sediment/ Rock Types: Mesozoic granitic rocks (Mileposts 0.4 to 7.7), alluvium. Geologic Hazards: None noted.
11	20.5	0.0	7.0	Sediment/ Rock Types: Alluvium, playa deposits of Soda Lake (Mileposts 1.8 to 3.6). Geologic Hazards: Inundation of playa following infrequent intense storms; shallow groundwater with associated liquefaction potential (Mileposts 0.0 to 6.0).
		7.0	20.5	Sediment/ Rock Types: Mesozoic volcanic and granitic rocks, Tertiary volcanic rocks, alluvium, young alluvium. Geologic Hazards: Slope stability in bedrock areas with moderate to high topographic relief; flash flooding potential (Mileposts 9.5 to 11.5); debris flow hazard (Mileposts 13.2 to 16.4). High potential for soil erosion in young alluvium between Mileposts 16.4 and 20.5.

Table 3-5 (continued)
Inventory of Geologic Resources

Link No.	Length	Mileposts		Description
		Begin	End	
12	4.9	0.0	4.9	Sediment/ Rock Types: Alluvium, playa deposits of Soda Lake (Mileposts 1.1 to 2.3), Tertiary sedimentary rocks. Geologic Hazards: Inundation of playa surface following infrequent intense storms; debris flows (Mileposts 4.0 to 4.9).
13	10.6	0.0	10.6	Sediment/ Rock Types: Alluvium, young alluvium, older alluvium, Tertiary sedimentary and volcanic rocks. Geologic Hazards: Slope instability, minor flash flood potential, debris flow potential (Mileposts 0.0 to 1.2, 6.0 to 7.4).
14	4.9	0.0	4.9	Sediment/ Rock Types: Young alluvium. Geologic Hazards: Possible shallow groundwater with associated liquefaction potential. High potential for soil erosion in young alluvium between Mileposts 1.8 and 4.8.
15	5.0	0.0	5.0	Sediment/ Rock Types: Alluvium, dune sand deposits, older alluvium, granitic bedrock. Geologic Hazards: Active sand dunes; slope instability in bedrock exposures with high topographic relief.

Table 3-5 (continued)
Inventory of Geologic Resources

Link No.	Length	Mileposts		Description
		Begin	End	
16	8.6	0.0	8.6	Sediment/ Rock Types: Alluvium, young alluvium, older alluvium, granitic bedrock. Geologic Hazards: Flash flood potential (at Milepost 4.0); slope instability in areas of high relief.
17	5.3	0.0	5.3	Sediment/ Rock Types: Alluvium, young alluvium, older alluvium, granitic bedrock. Geologic Hazards: Flash flood potential (Mileposts 1.0 to 2.7).
18	3.7	0.0	3.7	Sediment/ Rock Types: Older alluvium, includes young alluvium in drainages, as well as fossiliferous lake deposits from ancient Lake Manix. Geologic Hazards: Flash flood potential in drainages.
19	7.9	0.0	7.9	Sediment/ Rock Types: Alluvium, older alluvium, dune sands. Geologic Hazards: Active sand dunes. High potential for soil erosion in young alluvium between Mileposts 2.0 and 3.0, and in drainages between Mileposts 15.5 and 16.2.

Table 3-5 (continued)
Inventory of Geologic Resources

Link No.	Length	Mileposts		Description
		Begin	End	
20a	21.0	0.0	11.4	Sediment/ Rock Types: Alluvium, young alluvium, dune sands (between Mileposts 4.0 to 7.0), and playa deposits of Coyote Lake (Mileposts 8.4 to 10.3). Geologic Hazards: Active sand dunes; innudation of playa surface may occur following intense storms; shallow ground-water.
		11.4	21.0	Sediment/ Rock Types: Alluvium with some small areas of older alluvium (near Milepost 13.0); Paleozoic quartzite and hornfelsic metamorphic rocks; possible flash flooding in narrow drainages; bedrock outcrops may potentially have some mineral worth; debris flow potential (Mileposts 19.0 to 21.0).
20b	17.9	0.0	9.5	Sediment/ Rock Types: Alluvium, older alluvium, tertiary volcanic and sedimentary rocks, and pre-Tertiary intrusive igneous rocks. Geologic Hazards: Possible slope instability in bed-rock areas.
		9.5	17.9	Sediment/ Rock Types: Alluvial deposits. Geologic Hazards: Route crosses Harper Fault (Milepost 32.1).

Table 3-5 (continued)
Inventory of Geologic Resources

Link No.	Length	Mileposts		Description
		Begin	End	
21a	6.0	0.0	6.0	Sediment/ Rock Types: Alluvium, minor areas of older alluvium and Tertiary sedimentary rocks. Geologic Hazards: None noted.
21b	8.0	0.0	8.0	Sediment/ Rock Types: Alluvium, minor area of older alluvium and Tertiary sedimentary rocks. Geologic Hazards: Route crosses Calico Fault (Milepost 8.0) and Manix Fault (Milepost 6.1). High potential for soil erosion in young alluvium between Mileposts 13.1 and 13.6.
22	21.1	0.0	21.1	Sediment/ Rock Types: Older alluvium, alluvium, and younger alluvium including active channel of Mojave River. Geologic Hazards: Route coincides roughly with Manix Fault zone between Mileposts 8.8 and 13.5, and crosses Calico Fault at Milepost 21.1 flooding in channel and floodplain of Mojave River; liquefaction associated with strong ground motion and shallow groundwater. High potential for soil erosion in young alluvium at Milepost 8.2 and between Mileposts 19.7 and 21.2.

Table 3-5 (continued)
Inventory of Geologic Resources

Link No.	Length	Mileposts		Description
		Begin	End	
23	13.8	0.0	2.8	Sediment/ Rock Types: Young/active alluvium of Mojave River channel. Geologic Hazards: Flooding, erosion due to active stream channel, liquefaction; route crosses Calico Fault at milepost 0.0.
		2.8	13.8	Sediment/ Rock Types: Alluvium, older alluvium, Tertiary volcanic and sedimentary rocks. Geologic Hazards: Route crosses concealed trace of Camprock Fault (milepost 9.8); possible flash flooding (mileposts 10.0 to 13.8); slope instability in areas of moderate to high topographic relief (mileposts 12.0 to 13.8). High potential for soil erosion in young alluvium between mileposts 0.0 and 3.0.
24a	27.3	0.0	18.7	Sediment/ Rock Types: Alluvium, older alluvium, younger alluvium; Tertiary volcanics, granitic bedrock (mileposts 6.6 to 6.9). Geologic Hazards: Proposed route; two branches of Lenwood Fault cross corridor at Mileposts 0.6 and 1.1; flash flood and debris flows are potential hazards in drainages between Mileposts 10.0 and 18.7. High potential for soil erosion in drainages between Mileposts 5.5 and 7.5, and in young alluvium at Mileposts 10.8, 12.0, 14.5, and between 16.0 and 23.0.

Table 3-5 (continued)
Inventory of Geologic Resources

Link No.	Length	Mileposts		Description
		Begin	End	
24a cont.	13.5	18.7	22.6	Sediment/ Rock Types: Young (active) alluvium of Mojave River channel.
				Geologic Hazards: Flooding, active erosion and deposition in Mojave River channel; liquefaction potential associated with strong ground motion; route crosses Helendale Fault at Milepost 21.5.
		22.6	27.3	Sediment/ Rock Types: Alluvium, older alluvium.
				Geologic Hazards: Flooding, debris flows in Buckhorn Wash (Mileposts 27.2 to 27.3).
24b	6.1	0.0	6.1	Sediment/ Rock Types: Young alluvium, alluvium, older alluvium.
				Geologic Hazards: Flooding, debris flows in drainage channels.
25	11.7	0.0	11.7	Sediment/ Rock Types: Alluvium, Tertiary volcanic and sedimentary rocks, granitic bedrock.
				Geologic Hazards: Route crosses two branches of Lenwood Fault (at Mileposts 0.5 and 0.7); potential for flash flooding in Stoddard Valley.

Table 3-5 (continued)
Inventory of Geologic Resources

<u>Link No.</u>	<u>Length</u>	<u>Mileposts Begin</u>	<u>End</u>	<u>Description</u>
26	0.6	0.0	0.6	Sediment/ Rock Types: Alluvium, intrusive igneous bed-rock. Geologic Hazards: None noted.
27	13.4	0.0	13.4	Sediment/ Rock Types: Alluvium, paleozoic/mesozoic volcanic and intrusive igneous rocks, older alluvium, younger alluvium. Geologic Hazards: Route crosses Helendale Fault at Milepost 5.8; flooding, active erosion and deposition in Mojave River channel (Mileposts 11.8 to 13.2); liquefaction potential near Mojave River. High potential for soil erosion in drainage between Mileposts 6.0 and 10.0.
28	3.2	0.0	3.2	Sediment/ Rock Types: Older alluvium, alluvium. Geologic Hazards: Flooding, debris flows in drainage channels.
29	2.1	0.0	2.1	Sediment/ Rock Types: Young alluvium, alluvium, older alluvium. Geologic Hazards: Flooding and debris flows in Fremont Wash and associated branch drainages.

Table 3-5 (continued)
Inventory of Geologic Resources

Link No.	Length	Mileposts		Description
		Begin	End	
30	2.1	0.0	2.1	Sediment/ Rock Types: Alluvium, older alluvium. Geologic Hazards: Flooding and debris flows in incised drainage.
31	5.7	0.0	5.7	Sediment/ Rock Types: Alluvium. Geologic Hazards: Flooding, debris flows (Mileposts 0.0 to 2.0).
32	16.4	0.0	16.4	Sediment/ Rock Types: Alluvium, pre-Tertiary volcanic and intrusive igneous rocks, older alluvium, and young (active) alluvium of Mojave River channel (Mileposts 14.6 to 15.2). Geologic Hazards: Flooding, debris flows in minor drainages and near active channel of Mojave River (Mileposts 14.6 to 15.2); route crosses Helendale Fault at Milepost 4.8.
33	15.5	0.0	15.5	Sediment/ Rock Types: Alluvium, pre-Tertiary volcanic rocks, older alluvium. Geologic Hazards: Flash flooding, debris flow potential near active channel of Mojave River (Mileposts 14.8 to 14.9); route crosses Helendale Fault at Milepost 5.0.

Table 3-5 (continued)
Inventory of Geologic Resources

Link No.	Length	Mileposts		Description
		Begin	End	
34	1.4	0.0	1.4	Sediment/ Rock Types: Older alluvium. Geologic Hazards: Flash flooding, debris flows in channel at Milepost 0.8.
35	4.7	0.0	4.7	Sediment/ Rock Types: Alluvium, older alluvium. Geologic Hazards: Flash flooding, debris flows (between Mileposts 0.0 and 1.0).
36	5.8	0.0	5.8	Sediment/ Rock Types: Alluvium. Geologic Hazards: Minor flash flooding potential.
37	7.3	0.0	7.3	Sediment/ Rock Types: Alluvium, young and old alluvium, intrusive igneous bedrock, Tertiary volcanic and sedimentary rocks. Geologic Hazards: Flash flooding potential in gullies/narrow drainages; crosses South Lockhart Fault at Mile- post 0.2; debris flows between Mileposts 6.3 and 6.8.
38	10.3	0.0	4.7	Sediment/ Rock Types: Alluvium, older alluvium, younger alluvium. Geologic Hazards: Route crosses South Lockhart Fault (Milepost 0.3). High potential for soil erosion in young alluvium between Mile- posts 0.0 and 2.0.

Table 3-5 (continued)
Inventory of Geologic Resources

Link No.	Length	Mileposts		Description
		Begin	End	
38 cont.		4.7	10.8	Sediment/ Rock Types: Alluvium and older alluvium. Geologic Hazards: Possible flash flooding between Mileposts 3.5 and 10.8.
39	6.2	0.0	6.2	Sediment/ Rock Types: Alluvium, older alluvium, Tertiary sedimentary and volcanic rocks. Geologic Hazards: Route crosses Spring Fault at Milepost 2.8.
40a	6.2	0.0	6.2	Sediment/ Rock Types: Alluvium, older alluvium. Geologic Hazards: Flash floods and debris flows between Mileposts 0.0 and 3.0.
40b	4.9	0.0	4.9	Sediment/ Rock Types: Alluvium and older alluvium. Geologic Hazards: None noted.
41	15.5	0.0	15.5	Sediment/ Rock Types: Alluvium, older alluvium, young alluvium in drainages. Geologic Hazards: Flash flooding in drainages/ gullies following infrequent intense storms; route crosses Spring Fault at Milepost 0.2.

Table 3-5 (continued)
Inventory of Geologic Resources

Link No.	Length	Mileposts		Description
		Begin	End	
42	21.9	0.0	21.9	Sediment/ Rock Types: Alluvium, older alluvium, Tertiary sedimentary rocks (Milepost 17.3). Geologic Hazards: Flash flooding potential in channelized drainages; route crosses possible unnamed Quaternary faults (at Mileposts 0.2, 16.5 and 16.9), and Spring Fault at Milepost 19.8.
43	3.9	0.0	3.9	Sediment/ Rock Types: Alluvium and older alluvium. Geologic Hazards: None noted.
44	16.2	0.0	7.0	Sediment/ Rock Types: Alluvium. Geologic Hazards: None noted.
		7.0	16.2	Sediment/ Rock Types: Alluvium, pre-Tertiary igneous and metamorphic rocks.
45	5.7	0.0	5.7	Sediment/ Rock Types: Alluvium, playa deposits of Ivanpah Lake. Geologic Hazards: Inundation of playa surface following intense storms.

TABLE 3-6
INVENTORY OF PALEONTOLOGICAL RESOURCES

Link No.	Link	Length (Mi.)	Paleontologic Resources Description	Self-Inspection Potential
6	6	13.3	Miocene-age mammalian fossils	Moderate-Low
9	9	16.8	Permian-age invertebrate fossils including corals, fusulinids, brachiopods	Low-Moderate
18	18	10.5	Pleistocene-age mammalian fossils	Moderate-Low
19	19	8.3	Pleistocene-age mammalian fossils	Moderate-Low
20b	20b	7.1	Miocene-age mammalian fossils; also fossil palm fragments	Moderate-Low
22	22	8.4	Pleistocene-age mammalian fossils	Low-Moderate
23	23	1.2	Miocene-age mammalian fossils	Low-Moderate
25	25	1.9	Miocene-age mammalian fossils	Moderate-Low
24	24	0.2		Moderate-Low
25	25	8.8		Moderate-Low
26	26	12.3		Low-Moderate
27	27	1.2		Low-Moderate
28	28	8.9		Low-Moderate
29	29	10.1		Low-Moderate
30	30	10.1		Low-Moderate
31	31	10.1		Low-Moderate
32	32	10.1		Low-Moderate
33	33	10.1		Low-Moderate
34	34	10.1		Low-Moderate
35	35	10.1		Low-Moderate
36	36	10.1		Low-Moderate
37	37	10.1		Low-Moderate
38	38	10.1		Low-Moderate
39	39	10.1		Low-Moderate
40	40	10.1		Low-Moderate
41	41	10.1		Low-Moderate
42	42	10.1		Low-Moderate
43	43	10.1		Low-Moderate
44	44	10.1		Low-Moderate
45	45	10.1		Low-Moderate
46	46	10.1		Low-Moderate
47	47	10.1		Low-Moderate
48	48	10.1		Low-Moderate
49	49	10.1		Low-Moderate
50	50	10.1		Low-Moderate
51	51	10.1		Low-Moderate
52	52	10.1		Low-Moderate
53	53	10.1		Low-Moderate
54	54	10.1		Low-Moderate
55	55	10.1		Low-Moderate
56	56	10.1		Low-Moderate
57	57	10.1		Low-Moderate
58	58	10.1		Low-Moderate
59	59	10.1		Low-Moderate
60	60	10.1		Low-Moderate
61	61	10.1		Low-Moderate
62	62	10.1		Low-Moderate
63	63	10.1		Low-Moderate
64	64	10.1		Low-Moderate
65	65	10.1		Low-Moderate
66	66	10.1		Low-Moderate
67	67	10.1		Low-Moderate
68	68	10.1		Low-Moderate
69	69	10.1		Low-Moderate
70	70	10.1		Low-Moderate
71	71	10.1		Low-Moderate
72	72	10.1		Low-Moderate
73	73	10.1		Low-Moderate
74	74	10.1		Low-Moderate
75	75	10.1		Low-Moderate
76	76	10.1		Low-Moderate
77	77	10.1		Low-Moderate
78	78	10.1		Low-Moderate
79	79	10.1		Low-Moderate
80	80	10.1		Low-Moderate
81	81	10.1		Low-Moderate
82	82	10.1		Low-Moderate
83	83	10.1		Low-Moderate
84	84	10.1		Low-Moderate
85	85	10.1		Low-Moderate
86	86	10.1		Low-Moderate
87	87	10.1		Low-Moderate
88	88	10.1		Low-Moderate
89	89	10.1		Low-Moderate
90	90	10.1		Low-Moderate
91	91	10.1		Low-Moderate
92	92	10.1		Low-Moderate
93	93	10.1		Low-Moderate
94	94	10.1		Low-Moderate
95	95	10.1		Low-Moderate
96	96	10.1		Low-Moderate
97	97	10.1		Low-Moderate
98	98	10.1		Low-Moderate
99	99	10.1		Low-Moderate
100	100	10.1		Low-Moderate

TABLE 3-7
SOILS INVENTORY TABLE

Link No.	Length (Miles)	Mileposts		Soil Units	Soil Erosion Potential*
		Begin	End		
0	13.5	0.0	13.5	2, 4	Moderate-Low
1	16.6	0.0	7.0	4	Moderate-Low
		7.0	10.6	5	Low-Nonexistent
		10.6	16.6	4	Moderate-Low
2	6.5	0.0	1.8	4	Moderate-Low
		1.8	2.3	5	Low-nonexistent
		2.3	2.6	4	Moderate-Low
		2.6	2.7	5	Low-Nonexistent
		2.7	6.5	4, 2	Moderate-Low
3	8.4	0.0	0.2	5	Low-nonexistent
		0.2	4.5	4, 2	Moderate-Low
		4.5	8.4	7	Low-Nonexistent
4	1.9	0.0	1.7	2	Moderate-Low
		1.7	1.9	7	Low-Nonexistent
5a	0.2	0.0	0.2	4	Moderate-Low
5b	4.8	0.0	4.0	4	Moderate-Low
		4.0	4.8	7	Low-Nonexistent
5c	12.3	0.0	1.2	7	Low-Nonexistent
		1.2	6.9	4	Low-Moderate
		6.9	10.1	5	Low-Nonexistent
		10.1	12.3	6	Low-Moderate
6	83.3	0.0	0.3	7	
		0.3	1.4	2, 4	
		1.4	2.0	5	
		2.0	10.2	2, 4	
		10.2	12.0	5	
		12.0	29.4	6, 4, 2	
		29.4	31.5	5	
		31.5	33.6	4	
		33.6	34.2	5	
		34.2	38.8	4	
		38.8	40.3	5	
		40.3	41.8	4	

*Soil erosion potential is based on short-term, unmitigated impact.

Table 3-7 (continued)
Soils Inventory Table

<u>Link No.</u>	<u>Length (Miles)</u>	<u>Mileposts</u>		<u>Soil Units</u>	<u>Soil Erosion Potential</u>
		<u>Begin</u>	<u>End</u>		
6 (cont.)	32.5	41.8	42.5	5	Low-Nonexistent
		42.5	46.6	4	
		46.6	46.9	5	
		46.9	47.5	4	
		47.5	47.7	7	
		47.7	48.6	4	
		48.6	48.9	5	
		48.9	49.5	4	
		49.5	50.1	5	
		50.1	54.0	4	
		54.0	54.5	5	
		54.5	64.0	4	
		64.0	64.8	5	
		64.8	70.6	4	
		70.6	71.4	5	
		71.4	80.1	4, 6	
		80.1	80.3	5	
		80.3	80.6	3	
		80.6	80.8	4	
		80.8	82.0	5	
		82.0	82.7	3	
		82.7	83.3	4, 6	
7	13.6	0.0	7.5	6, 4	
		7.5	9.6	5	
		9.6	12.1	4	
		12.1	13.3	5	
		13.3	13.6	4	
8	12.9	0.0	6.0	6, 4	
		6.0	7.4	7	
		7.0	12.9	4	
9	25.0	0.0	0.5	4	
		0.5	1.8	5	
		1.8	24.3	4	
		24.3	25.0	5	
10	19.7	0.0	0.3	4	
		0.3	7.7	5	
		7.7	19.7	4	

Table 3-7 (continued)
Soils Inventory Table

<u>Link No.</u>	<u>Length (Miles)</u>	<u>Mileposts</u>		<u>Soil Units</u>	<u>Soil Erosion Potential</u>
		<u>Begin</u>	<u>End</u>		
11	20.5	0.0	1.8	4	
		1.8	3.6	7	
		3.6	6.8	4	
		6.8	9.1	5	
		9.1	12.4	4	
		12.4	13.2	5	
		13.2	16.4	6	
		16.4	20.5	1	
12	5.0	0.0	1.3	4	
		1.3	2.3	7	
		2.3	4.3	4	
		4.3	5.0	5	
13	10.6	0.0	1.2	5	
		1.2	7.5	4	
		7.5	8.3	5	
		8.3	10.6	6, 4, 2	
14	4.9	0.0	1.7	2	
		1.7	4.9	1	
15	5.0	0.0	0.8	1	
		0.8	1.7	5	
		1.7	2.1	3	
		2.1	2.5	5	
		2.5	2.8	3	
		2.8	3.5	5	
		3.5	4.3	6	
		4.3	4.6	5	
16	8.6	4.6	5.0	6	
		0.0	7.5	2, 4	
		7.5	7.7	5	
		7.7	8.1	4	
		8.1	8.6	5	

Table 3-7 (continued)
Soils Inventory Table

<u>Link No.</u>	<u>Length (Miles)</u>	<u>Mileposts</u>		<u>Soil Units</u>	<u>Soil Erosion Potential</u>
		<u>Begin</u>	<u>End</u>		
17	5.3	0.0	0.9	1	
		0.9	2.7	4	
		2.7	3.4	5	
		3.4	4.8	6	
		4.8	4.9	5	
		4.9	5.3	6	
18	3.7	0.0	3.7	6	
19	7.9	3.7	4.8	6, 4	
		4.8	5.0	3	
		5.0	5.3	4	
		5.3	5.5	3	
		5.5	7.9	4, 6	
20a	21.0	0.0	4.5	4	
		4.5	5.6	3	
		5.6	5.9	4	
		5.9	7.6	3	
		7.6	9.0	2	
		9.0	10.9	7	
		10.9	21.0	2, 6, 4	
20b	17.9	0.0	3.8	4	
		3.8	4.1	5	
		4.1	8.4	4, 6	
		8.4	10.3	5	
		10.3	10.8	3	
		10.8	17.9	4	
21a	6.0	0.0	6.0	6, 4, 2	
21b	8.0	0.0	8.0	6, 4, 2	
22	21.1	0.0	18.1	6, 4	
		18.1	21.1	1	
23	13.8	0.0	2.7	1	
		2.7	10.6	2, 6	
		10.6	11.1	5	
		11.1	12.4	4, 6	
		12.4	13.8	5	

Table 3-7 (continued)
Soils Inventory Table

<u>Link No.</u>	<u>Length (Miles)</u>	<u>Mileposts</u>		<u>Soil Units</u>	<u>Soil Erosion Potential</u>
		<u>Begin</u>	<u>End</u>		
24a	27.3	0.0	0.5	5	
		0.5	0.9	4	
		0.9	1.1	5	
		1.1	6.6	4	
		6.6	6.9	5	
		6.9	11.8	4, 6	
		11.8	12.1	1	
		12.1	12.4	6	
		12.4	14.2	5	
		14.2	14.7	1	
		14.7	16.2	6	
		16.2	17.0	1	
		17.0	?	4	
		?	27.1	6	
		27.1	27.3	1	
24b	6.1	0.0	0.1	1	
		0.1	5.8	6, 2	
		5.8	6.1	1	
25	11.7	0.0	0.5	5	
		0.5	0.7	4	
		0.7	1.8	5	
		1.8	11.4	4, 2	
		11.4	11.7	5	
26	0.6	0.0	0.4	5	
		0.4	0.6	4	
27	13.4	0.0	0.8	4	
		0.8	1.8	5	
		1.8	5.7	4	
		5.7	7.2	5	
		7.2	11.6	6, 4	
		11.6	13.1	1	
		13.1	13.4	6	
28	3.2	0.0	3.2	6, 4	
29	2.1	0.0	0.8	1	
		0.8	2.1	4, 6	

Table 3-7 (continued)
Soils Inventory Table

<u>Link No.</u>	<u>Length (Miles)</u>	<u>Mileposts</u>		<u>Soil Units</u>	<u>Soil Erosion Potential</u>
		<u>Begin</u>	<u>End</u>		
30	2.1	0.0	2.1	6, 4	
31	5.7	0.0	5.7	4	
32	16.0	0.0	4.7	4	
		4.7	5.1	5	
		5.1	6.4	4	
		6.4	6.6	5	
		6.6	11.3	4, 6	
		11.3	11.6	5	
		11.6	12.2	4	
		12.2	14.6	5	
		14.6	15.1	1	
		15.1	16.0	6	
33	15.3	0.0	0.4	5	
		0.4	5.0	4	
		5.0	5.3	5	
		5.3	14.0	4, 6	
		14.0	14.7	5	
		14.7	14.9	1	
		14.9	15.3	6	
34	1.4	0.0	1.4	6	
35	5.1	0.0	5.1	6, 4	
36	5.9	0.0	0.1	1	
		0.1	5.9	4	
37	7.3	0.0	0.5	1	
		0.5	1.3	4, 6	
		1.3	3.2	5	
		3.2	4.6	4, 6	
		4.6	4.8	5	
		4.8	6.3	4	
		6.3	6.8	5	
		6.8	7.3	4, 6	
38	10.3	0.0	0.7	1	
		0.7	10.3	6, 2, 4	

Table 3-7 (continued)
Soils Inventory Table

<u>Link No.</u>	<u>Length (Miles)</u>	<u>Mileposts</u>		<u>Soil Units</u>	<u>Soil Erosion Potential</u>
		<u>Begin</u>	<u>End</u>		
39	6.2	0.0	0.3	6	
		0.3	0.7	5	
		0.7	3.0	4, 6	
		3.0	3.2	5	
		3.2	6.2	4, 2	
40	11.1	0.0	11.1	2, 4, 6	
41	15.5	0.0	10.9	4, 6	
		10.9	11.5	1	
		11.5	13.8	6	
		13.8	14.1	1	
		14.1	15.5	6	
42	21.9	0.0	17.1	4, 6	
		17.1	17.3	5	
		17.3	21.9	6, 2	
43	3.9	0.0	0.2	1	
		0.2	3.9	2, 4	
44	16.2	0.0	7.0	2, 4	Low-Moderate Low-Moderate, Low- Nonexistent
		7.0	16.2	4, 5	
45	5.7	0.0	5.7	2, 4	Low-Moderate

TABLE 3-8
SOIL EROSION IMPACT ASSESSMENT MODEL

<u>Sensitivity (Soil Erosion Potential)</u>	<u>Quality and Quantity (Access Level)</u>	<u>Impact Potential</u>	
		<u>Long-Term</u>	<u>Short-Term</u>
High	(1) Sufficient	Low	Low
	(2) Upgrade	Low	Moderate
	(3) New	Low	High
Moderate to Low	(1) Sufficient	Low	Low
	(2) Upgrade	Low	Low
	(3) New	Moderate to Low	Moderate to Low
Low	(1) Sufficient	Negligible	Negligible
	(2) Upgrade	Low to Negligible	Low to Negligible
	(3) New	Low	Low

**TABLE 3-9
PALEONTOLOGICAL RESOURCES IMPACT ASSESSMENT MODEL**

<u>Sensitivity</u>	<u>Quality and Quantity (Access Level)</u>	<u>Impact Potential (Long-Term)</u>
High to Moderate	(1) Sufficient (2) Upgrade (3) New	Low to Negligible Moderate to Low High to Moderate
Moderate to Low	(1) Sufficient (2) Upgrade (3) New	Negligible Low Moderate to Low

TABLE 3-10
IMPACT ASSESSMENT/MITIGATION PLANNING CHART
SOIL EROSION

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
0	13.5	0.0-13.5	I	L (.5)	-	Sufficient Access/ Soil Types 2 and 4	-	-	-	-
1	16.6	0.0- 7.3	I	L (.5)	-	Sufficient Access/ Soil Types 2 and 4	-	-	-	-
		7.3-10.5	I	L/N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		10.5-16.6	I	L (.5)	-	Sufficient Access/ Soil Type 4	-	-	-	-
2	6.5	0.0- 6.5	I	L (.5)	-	Sufficient Access/ Soil Types 2 and 4 (Type 5 from Mileposts 1.8-2.3 and 2.5-2.7)	-	-	-	-
3	8.4	0.0- 4.5	I	L (.5)	-	Sufficient Access/ Soil Types 2 and 4	-	-	-	-
		4.5- 6.4	I	N (0)	-	Sufficient Access/ Soil Type 7	-	-	-	-
		6.4- 8.4	I	L (.5)	-	New Access/ Soil Type 7	-	-	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
4	1.9	0.0- 1.9	I	L (.5)	-	Sufficient Access/ Soil Type 2	-	-	-	-
5a	0.2	0.0- 0.2	I	L (.5)	-	Sufficient Access/ Soil Type 2	-	-	-	-
5b	4.8	0.0- 4.0	I	L (.5)	-	Sufficient Access/ Soil Type 2	-	-	-	-
		4.0- 4.8	I	N (0)	-	Sufficient Access/ Soil Type 7	-	-	-	-
5c	12.3	0.0- 1.2	I	N (0)	-	Sufficient Access/ Soil Type 7	-	-	-	-
		1.2- 6.9	I	L (.5)	-	Sufficient Access/ Soil Types 2 and 4	-	-	-	-
		6.9- 7.3	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		7.3- 9.0	I	N (0)	-	New Access/Soil Type 5	-	-	-	-
		9.0-10.1	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Impact Description</u>
5c (cont.)		10.1-12.3	I	L (.5)	-	Sufficient Access/ Soil Type 6	-	-	-	-
6	83.3	0.0- 1.4	I	L (.5)	-	Sufficient Access/ Soil Types 2 and 4	-	-	-	-
		1.4- 2.0	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		2.0-10.1	I	L (.5)	-	Sufficient Access/ Soil Types 2 and 4	-	-	-	-
		10.1-12.0	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		12.0-29.4	I	L (.5)	-	Sufficient Access/ Soil Types 2, 4 and 6	-	-	-	-
		29.4-31.2	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		31.2-42.0	I	L (.5)	-	Sufficient Access/ Soil Type 4 (Type 5 from Milepost 33.6- 34.2)	-	-	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
6 (cont.)		42.0-42.4	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		42.4-48.5	I	L (.5)	-	Sufficient Access/ Soil Type 4 (Type 5 from Milepost 47.1- 47.3)	-	-	-	-
		48.5-49.0	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		49.0-49.5	I	L (.5)	-	Sufficient Access/ Soil Type 4	-	-	-	-
		49.5-50.0	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		50.0-63.8	I	L (.5)	-	Sufficient Access/ Soil Type 4 (Type 5 at Mileposts 59.5 and 60.0)	-	-	-	-
		63.8-64.5	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

Link Number	Length	Mileposts	Impact Type	Initial Long- Term Impact Level	Sig- nifi- cant?	Impact Description	Recom- mended Miti- gation	Residual Long- Term Impact Level	Sig- nifi- cant?	Significant Unavoidable Adverse Impact Description
6 (cont.)	64.5-79.8		I	L (.5)	-	Sufficient Access/ Soil Types 4 and 6	-	-	-	-
	79.8-81.8		I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
	81.8-83.5		I	L (.5)	-	Sufficient Access/ Soil Type 4	-	-	-	-
7	0.0- 7.4		I	M/L (5)	Yes	New Access/ Soil Types 4 and 6	12	L (I)	-	-
	7.4- 9.6		I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
	9.6-12.1		I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (I)	-	-
10	12.1-13.3		I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
	13.3-13.5		I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (I)	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
8	12.9	0.0- 5.9	I	L (.5)	-	Sufficient Access/ Soil Types 4 and 6	-	-	-	-
						Sufficient Access/ Soil Type 7				
						Sufficient Access/ Soil Type 4				
						Upgrade Access/ Soil Type 4				
9	25.0	0.0- 0.3	I	L (.5)	-	Upgrade Access/ Soil Type 5	-	-	-	-
						Upgrade Access/ Soil Type 4				
						Upgrade Access/ Soil Type 5				
						Upgrade Access/ Soil Type 4				
10	19.9	0.0- 0.3	I	M/L (.5)	Yes	New Access/ Soil Type 4	12	L (I)	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
10 (cont.)		0.3- 7.8	I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
		7.8- 9.0	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (1)	-	-
		9.0-19.9	I	L (.5)	-	Upgrade Access/ Soil Type 4	-	-	-	-
11	20.5	0.0- 6.8	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (1)	-	-
		6.8- 9.1	I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
		9.1-12.5	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (1)	-	-
		12.5-13.2	I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
		13.2-16.3	I	M/L (5)	Yes	New Access/ Soil Type 6	12	L (1)	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
11 (cont.)		16.3-20.5	I	L (.5)	-	New Access/ Soil Type 1 (high short-term impact)	-	-	-	-
12	4.9	0.0- 4.3	I	M/L (.5)	Yes	New Access/ Soil Type 4	12	L (I)	-	-
		4.3- 4.9	I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
13	10.6	0.0- 1.2	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		1.2- 7.0	I	L (.5)	-	Sufficient Access/ Soil Type 4	-	-	-	-
		7.0- 7.5	I	L (.5)	-	Upgrade Access/ Soil Type 4	-	-	-	-
		7.5- 8.3	I	L/N (0)	-	Upgrade Access/ Soil Type 5	-	-	-	-
		8.3-10.6	I	L (.5)	-	Upgrade Access/ Soil Types 4 and 6	-	-	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
14	4.9	0.0- 4.9	I	L (.5)	-	New Access/ Soil Type 1 (high short-term impact)	-	-	-	-
15	5.0	0.0- 0.8	I	L (.5)	-	New Access/ Soil Type 1 (high short-term impact)	-	-	-	-
		0.8- 2.4	I	L (.5)	-	New Access/ Soil Type 5 (Type 3 at Mileposts 1.6 and 2.0)	-	-	-	-
		2.4- 2.8	I	L (.5)	-	New Access/ Soil Type 3 (high short-term impact)	-	-	-	-
		2.8- 3.4	I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
		3.4- 5.0	I	M/L (5)	Yes	New Access/ Soil Types 4 and 6	12	L (I)	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
16	8.6	0.0- 8.6	I	L (.5)	-	Upgrade Access/ Soil Types 2, 4 and 6 (Type 5 at Mileposts 6.9 and 7.5)	-	-	-	-
17	5.3	0.0- 0.9	I	L (.5)	-	New Access/ Soil Type 1 (high short-term impact)	-	-	-	-
		0.9- 2.7	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (I)	-	-
		2.7- 3.3	I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
		3.3- 5.3	I	M/L (5)	Yes	New Access/ Soil Types 4 and 6 (high short-term impact)	12	L (I)	-	-
18	3.7	0.0- 3.7	I	L (.5)	-	Upgrade Access/ Soil Type 6	-	-	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

Link Number	Length	Mileposts	Impact Type	Initial Long-Term Impact Level	Sig-nificant?	Impact Description	Recom-mended Mitigation	Residual Long-Term Impact Level	Sig-nificant?	Significant Unavoidable Adverse Impact Description
19	7.9	0.0- 4.8	I	M/L (5)	Yes	New Access/ Soil Types 4 and 6	12	L (I)	-	-
		4.8- 5.0	I	L (.5)	-	New Access/ Soil Type 3 (high short-term impact)	-	-	-	-
		5.0- 5.3	I	M/L (5)	Yes	New Access/ Soil Types 4 and 6	12	L (I)	-	-
		5.3- 5.5	I	L (.5)	-	New Access/ Soil Type 3 (high short-term impact)	-	-	-	-
		5.5- 7.9	I	M/L (5)	Yes	New Access/ Soil Types 4 and 6	12	L (I)	-	-
20a	21.0	0.0- 4.5	I	M/L (5)	Yes	New Access/ Soil Types 4 and 6	12	L+ (I)	-	-
		4.5- 5.6	I	L (.5)	-	New Access/ Soil Type 3	-	-	-	-
		5.6- 5.9	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L+ (I)	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

Link Number	Length	Mileposts	Impact Type	Initial Long-Term Impact Level	Sig-nifi-cant?	Impact Description	Recom-mended Mitigation	Residual Long-Term Impact Level	Sig-nifi-cant?	Significant Unavoidable Adverse Impact Description
20a (cont.)		5.9- 7.6	I	L (.5)	-	New Access/ Soil Type 3	-	-	-	-
		7.6- 9.0	I	M/L (5)	Yes	New Access/ Soil Type 2	12	L+ (1)	-	-
		9.0-10.9	I	L (.5)	-	New Access/ Soil Type 7	-	-	-	-
		10.9-21.0	I	M/L (5)	Yes	New Access/ Soil Types 2, 4 and 6	12	L+ (1)	-	-
20b	17.9	0.0- 2.7	I	L (.5)	-	Upgrade Access/ Soil Type 4	-	-	-	-
		2.7- 3.8	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L+ (1)	-	-
		3.8- 4.1	I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
		4.1- 7.8	I	M/L (5)	Yes	New Access/ Soil Type 4 and 6	12	L+ (1)	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
20b (cont.)		7.8- 8.4	I	L (.5)	-	Upgrade/Soil Types 4 and 6	-	-	-	-
		8.4-10.3	I	L (.5)	-	Upgrade Access/ Soil Type 5	-	-	-	-
		10.3-10.8	I	L (.5)	-	Upgrade Access/Soil Type 3 (high ST impacts)	-	-	-	-
		10.8-17.9	I	L (.5)	-	Upgrade Access/ Soil Type 4	-	-	-	-
21a	6.0	0.0- 6.0	I	L (.5)	-	Sufficient Access/ Soil Types 2, 4 and 6	-	-	-	-
21b	8.0	0.0- 7.5	I	L (.5)	-	Sufficient Access/ Soil Types 2, 4 and 6	-	-	-	-
		7.5- 8.0	I	L (.5)	-	Sufficient Access/ Soil Type 1 (high short-term impact)	-	-	-	-
22	21.1	0.0-19.7	I	L (.5)	-	Upgrade Access/ Soil Types 4 and 6	-	-	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
22 (cont.)		19.7-21.1	I	L (.5)	-	Upgrade Access/ Soil Type 1 (moderate short-term impact)	-	-	-	-
23	13.8	0.0- 2.8	I	L (.5)	-	Sufficient Access/ Soil Type 1	-	-	-	-
		2.8-10.7	I	L (.5)	-	Sufficient Access/ Soil Types 2, 4 and 6	-	-	-	-
		10.7-11.1	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		11.1-12.5	I	L (.5)	-	Sufficient Access/ Soil Types 4 and 6	-	-	-	-
		12.5-13.8	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
24a	27.3	0.0- 0.6	I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
		0.6- 0.9	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (1)	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Mitig- ation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
24a (cont.)		0.9 - 1.9	I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
		1.9 - 6.6	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (1)	-	-
		6.6 - 6.9	I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
		6.9 - 11.9	I	M/L (5)	Yes	New Access/ Soil Types 4 and 6	12	L (1)	-	-
		11.9 - 12.1	I	L (.5)	-	New Access/ Soil Type 1 (high short-term impact)	-	-	-	-
		12.1 - 12.4	I	M/L (5)	Yes	New Access/ Soil Type 6	12	L (1)	-	-
		12.4 - 14.3	I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
		14.3 - 14.7	I	L (.5)	-	New Access/ Soil Type 1 (high short-term impact)	-	-	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
24a (cont.)										
		14.7-16.7	I	M/L (5)	Yes	New Access/ Soil Type 6	12	L (I)	-	-
		16.7-17.0	I	L (.5)	-	New Access/ Soil Type I (high short-term impact)	-	-	-	-
		17.0-17.6	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (I)	-	-
		17.6-17.7	I	L (.5)	-	New Access/ Soil Type I (high short-term impact)	-	-	-	-
		17.7-18.7	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (I)	-	-
		18.7-23.0	I	L (.5)	-	New Access/ Soil Type I (high short-term impact)	-	-	-	-
		23.0-27.1	I	M/L (5)	Yes	New Access/ Soil Type 6	12	L (I)	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
24a (cont.)		27.1-27.3	I	L (.5)	-	New Access/ Soil Type I (high short-term impact)	12	-	-	-
24b			I	L (.5)	-	New Access/ Soil Type I (high short-term impact)	-	-	-	-
			I	M/L (5)	Yes	New Access/ Soil Types 4 and 6	12	L (I)	-	-
			I	L (.5)	-	New Access/ Soil Type I (high short-term impact)	-	-	-	-
25	11.7	0.0- 0.4	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		0.4- 0.6	I	L (.5)	-	Sufficient Access/ Soil Type 4	-	-	-	-
		0.6- 1.8	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
25 (cont.)		1.8-11.3	I	L (.5)	-	Sufficient Access/ Soil Types 2 and 4	-	-	-	-
		11.3-11.7	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
26	0.6	0.0- 0.4	I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
		0.4- 0.6	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (1)	-	-
27	13.4	0.0- 0.8	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (1)	-	-
		0.8- 1.8	I	L/N (.1)	-	New Access/ Soil Type 5	-	L	-	-
		1.8- 5.8	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (1)	-	-
		5.8- 7.2	I	L/N (.1)	-	New Access Soil Type 5	-	-	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
27 (cont.)		7.2-10.0	I	M/L (5)	Yes	New Access/ Soil Types 4 and 6	12	L (I)	-	-
		10.0-11.6	I	L (.5)	-	Upgrade Access/ Soil Types 4 and 6	-	-	-	-
		11.6-13.2	I	L (.5)	-	Upgrade Access/ Soil Type 1	-	-	-	-
		13.2-13.4	I	L (.5)	-	Upgrade Access/ Soil Type 6 (moderate short-term impact)	-	-	-	-
28	3.2	0.0- 2.1	I	M/L (5)	Yes	New Access/ Soil Types 4 and 6	12	L (I)	-	-
		2.1- 3.2	I	L (.5)	-	Sufficient Access/ Soil Types 4 and 6	-	-	-	-
29	2.1	0.0- 0.8	I	L (.5)	-	New Access/ Soil Type 1 (high short-term impact)	-	-	-	-
		0.8- 2.1	I	M/L (5)	Yes	New Access/ Soil Types 4 and 6	12	L (I)	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
30	2.1	0.0- 2.1	I	L (.5)	-	Sufficient Access/ Soil Types 4 and 6	-	-	-	-
31	5.7	0.0- 5.7	I	L (.5)	-	Sufficient Access/ Soil Type 4	-	-	-	-
32	16.4	0.0- 4.8	I	L (.5)	-	Sufficient Access/ Soil Type 4	-	-	-	-
		4.8- 5.0	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		5.0-11.2	I	L (.5)	-	Sufficient Access/ Soil Type 4 (Type 5 at Milepost 6.5)	-	-	-	-
		11.2-11.6	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		11.6-12.2	I	L (.5)	-	Sufficient Access/ Soil Type 4	-	-	-	-
		12.2-14.4	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Mitiga- tion</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
32 (cont.)		14.4-15.3	I	L (.5)	-	Sufficient Access/ Soil Type 1	-	-	-	-
		15.3-16.4	I	L (.5)	-	Sufficient Access/ Soil Type 6	-	-	-	-
33	15.5	0.0- 0.4	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		0.4- 5.0	I	L (.5)	-	Sufficient Access/ Soil Type 4	-	-	-	-
		5.0- 5.3	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		5.3- 6.5	I	L (.5)	-	Sufficient Access/ Soil Type 4	-	-	-	-
		6.5-11.0	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (1)	-	-
		11.0-14.0	I	L (.5)	-	Sufficient Access/ Soil Types 4 and 6	-	-	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
33 (cont.)		14.0-14.7	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		14.7-14.9	I	L (.5)	-	Sufficient Access/ Soil Type 1	-	-	-	-
		14.9-15.5	I	L (.5)	-	Sufficient Access/ Soil Type 6	-	-	-	-
34	1.4	0.0- 1.4	I	L (.5)	-	Sufficient Access/ Soil Type 6	-	-	-	-
35	4.7	0.0- 4.7	I	L (.5)	-	Sufficient Access/ Soil Types 4 and 6	-	-	-	-
36	5.8	0.0- 2.5	I	M/L (5)	Yes	New Access/ Soil Types 4 and 6	12	L (I)	-	-
		2.5- 4.3	I	L (.5)	-	Upgrade Access/ Soil Type 4	-	-	-	-
		4.3- 5.8	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (I)	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
37	7.3	0.0- 0.4	I	L (.5)	-	New Access/ Soil Type 1 (high short-term impact)	-	-	-	-
		0.4- 1.3	I	M/L (5)	Yes	New Access/ Soil Types 4 and 6	12	L (I)	-	-
		1.3- 3.2	I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
		3.2- 4.6	I	M/L (5)	Yes	New Access/ Soil Types 4 and 6	12	L (I)	-	-
		4.6- 4.8	I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
		4.8- 6.2	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (I)	-	-
		6.2- 6.6	I	L (.5)	-	New Access/ Soil Type 5	-	-	-	-
		6.6- 7.3	I	M/L (5)	Yes	New Access/ Soil Types 4 and 6	12	L (I)	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
38	10.3	0.0- 0.8	I	L (.5)	-	Sufficient Access/ Soil Type 1	-	-	-	-
		0.8-10.3	I	L (.5)	-	Sufficient Access/ Soil Types 2, 4 and 6	-	-	-	-
39	6.2	0.0- 0.3	I	L (.5)	-	Sufficient Access/ Soil Type 6	-	-	-	-
		0.3- 0.6	I	N (0)	-	Sufficient Access/ Soil Type 5	-	-	-	-
		0.6- 6.2	I	L (.5)	-	Sufficient Access/ Soil Types 2, 4 and 6	-	-	-	-
40a	6.2	0.0- 6.2	I	L (.5)	-	Sufficient Access/ Soil Types 2, 4 and 6	-	-	-	-
40b	4.9	0.0- 4.9	I	L (.5)	-	Sufficient Access/ Soil Types 4 and 6	-	-	-	-
41	15.5	0.0-10.9	I	L (.5)	-	Sufficient Access/ Soil Types 4 and 6	-	-	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
41 (cont.)		10.9-11.5	I	L (.5)	-	Sufficient Access/ Soil Type 1	-	-	-	-
		11.5-13.8	I	L (.5)	-	Sufficient Access/ Soil Type 6	-	-	-	-
		13.8-14.1	I	L (.5)	-	Sufficient Access/ Soil Type 1	-	-	-	-
		14.1-15.5	I	L (.5)	-	Sufficient Access/ Soil Type 6	-	-	-	-
42	21.9	0.0- 2.4	I	M/L (5)	Yes	New Access/ Soil Type 4	12	L (1)	-	-
		2.4-17.2	I	L (.5)	-	Upgrade Access/ Soil Types 4 and 6	-	-	-	-
		17.2-17.4	I	L/N (0)	-	Upgrade Access/ Soil Type 5	-	-	-	-
		17.4-21.9	I	L (.5)	-	Upgrade Access/ Soil Types 2 and 6	-	-	-	-

Table 3-10 (continued)
Impact Assessment/Mitigation Planning Chart
Soil Erosion

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>	<u>Impact Type</u>	<u>Initial Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Impact Description</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Long- Term Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Significant Unavoidable Adverse Impact Description</u>
43	3.9	0.0- 0.2	I	L (.5)	-	New Access/ Soil Type 1 (high short-term impact)	-	-	-	-
		0.2- 3.9	I	M/L (5)	Yes	New Access/ Soil Types 2, 4 and 6	12	L (1)	-	-
44	16.2	0.0- 3.4	I	L (.5)	-	Upgrade Access/ Soil Type 4	-	-	-	-
		3.4- 7.0	I	M/L (5)	Yes	New Access/Soil Type 4	12	L (1)	-	-
		7.0-16.2	I	M/L (5)	Yes	New Access/Soil Types 4 and 5	12	L (1)	-	-
45	5.7	0.0- 5.7	I	M/L (5)	Yes	New Access/Soil Types 2 and 4	12	L (1)	-	-

TABLE 3-11
IMPACT ASSESSMENT/MITIGATION PLANNING CHART
PALEONTOLOGICAL RESOURCES

Link Number	Length	Mileposts	Impact Type	Initial Long-Term Impact Level	Sig-nifi-cant?	Impact Description	Recom-mended Mitigation	Residual Long-Term Impact Level	Sig-nifi-cant?	Significant Unavoidable Adverse Impact Description
6	83.3	79.7-82.0	2	N (1)	-	Sufficient Access/ possible fossil-bearing strata	-	-	-	-
18	3.7	0.5- 3.7	2	M/L (5)	Yes	Upgrade Access/ possible late Pleisto- cene vertebrate fossils in lake deposits	12A	L (1)	No	-
19	7.9	0.0- 0.8	2	H/M (11)	Yes	New Access/ possible late Pleisto- cene vertebrate fossils in lake deposits	12A	M/L (5)	No	-
20b	17.9	2.0- 2.7	2	L (5)	-	Upgrade Access/ possible fossil-bearing strata	-	-	-	-
22	21.1	0.0- 2.0	2	M/L (5)	Yes	Upgrade Access/ possible late Pleisto- cene vertebrate fossils in lake deposits	12A	L (1)	No	-

TABLE OF CONTENTS

	PAGE
CHAPTER 4 - BIOLOGICAL RESOURCES	4-1
Introduction	4-1
Study Approach and Methods	4-1
Study Personnel	4-2
Study Area Overview	4-2
Vegetation Types	4-2
Wildlife	4-5
Legally Protected Species	4-12
Species of Special Interest	4-19
Areas of Special Biological Interest	4-25
Wetlands	4-33
Biological Resources Within and Adjacent to the Alternative Corridors	4-33
Vegetation Types	4-34
Wildlife	4-35
Legally Protected Species	4-36
Species of Special Interest	4-39
Areas of Special Biological Interest	4-41
Agency Contacts	4-41
Impact Assessment and Mitigation Planning	4-41
Introduction	4-41
Impact Assessment Model	4-43
Mitigation Planning	4-43
Residual Residual Impact Levels	4-44
Conclusions	4-47
Agency Contacts (Balance page)	4-47
References	4-47

CHAPTER 4 BIOLOGICAL RESOURCES

TABLE OF CONTENTS

	PAGE
CHAPTER 4 - BIOLOGICAL RESOURCES	4- 1
Introduction	4- 1
Study Approach and Methods	4- 1
Study Personnel	4- 2
Study Area Overview	4- 3
Vegetation Types	4- 3
Wildlife	4- 8
Legally Protected Species	4-12
Species of Special Interest	4-19
Areas of Special Biological Interest	4-28
Wetlands	4-33
Biological Resources Within and Adjacent to the Alternative Corridors	4-33
Vegetation Types	4-34
Wildlife	4-35
Legally Protected Species	4-36
Species of Special Interest	4-38
Areas of Special Biological Interest	4-41
Agency Contacts	4-44
Impact Assessment and Mitigation Planning	4-45
Introduction	4-45
Impact Assessment Model	4-45
Mitigation Planning	4-53
Results: Residual Impact Levels	4-56
Conclusions	4-57
Agency Contacts (follows page)	4-57
References	4-58

LIST OF TABLES

TABLE NUMBER	TITLE	FOLLOWS PAGE
4- 1	Common and Characteristic Plant Species in the Study Area	4-63
4- 2	Common and Characteristic Amphibians and Reptiles of the Study Area	4-63
4- 3	Common and Characteristic Birds of the Study Area	4-63
4- 4	Common and Characteristic Mammals of the Study Area	4-63
4- 5	Plant Species of Special Interest in the Study Area	4-63
4- 6	Vegetation Types Traversed by the Alternative Corridors	4-63
4- 7	Desert Tortoise Habitat Traversed by the Alternative Corridors	4-63
4- 8	Mohave Ground Squirrel Habitat Traversed by the Alternative Corridors	4-63
4- 9	Potential Habitat of Special Interest Plant Species Traversed by the Alternative Corridors	4-63
4-10	Potential and Known Golden Eagle Nesting Habitat Near the Alternative Corridors	4-63
4-11	Desert Bighorn Sheep Habitat Traversed by the Alternative Corridors	4-63
4-12	Sensitivity of Biological Resources Potentially Affected by the Proposed Project	4-63
4-13	Impact Level Model	4-63
4-14	Matrix of Recommended Selective Mitigation Measures and Affected Resources	4-63
4-15	Changes in Impact Level Due to Application of Recommended Selective Mitigation Measures	4-63
4-16	Impact Assessment/Mitigation Planning Chart, Biological Resources	4-63

List of Tables (continued)

TABLE NUMBER	TITLE	FOLLOWS PAGE
-----------------	-------	-----------------

APPENDIX A TABLES

A- 1	Plant Species of Potential Special Interest Reviewed and Removed from Further Consideration	4-63
------	---	------

APPENDIX B TABLES

B- 1	Inventory of Biological Resources Within and Adjacent to Corridors	4-63
------	--	------

APPENDIX C TABLES

C- 1	Plant Species Observed During Field Surveys	4-63
------	---	------

CHAPTER 4

BIOLOGICAL RESOURCES

INTRODUCTION

The Los Angeles Department of Water and Power (DWP), a member of the Southern California Public Power Authority (SCPPA), together with other participants, proposes to construct, operate and maintain a 500kV transmission line from the Mead or McCullough Switching Station near Boulder City, Nevada, to the Victorville or Adelanto Switching Station in California. This report presents results of technical studies on the biological resources that could potentially be affected by the proposed project.

STUDY APPROACH AND METHODS

The proposed project extends across more than 150 miles of biological habitats in the Mojave Desert of California and Nevada. The large areal extent of the proposed project required that numerous biological resources be examined. These resources varied greatly in type, location, abundance and sensitivity due in part to the wide geographic range of the project. As a result, a two-level method of analysis was utilized to summarize and describe biological resources that could be affected by the project. A regional-level analysis was conducted by identifying important biological resources near the alternative corridors. A regional study area was defined using an approximate 10-mile-wide zone surrounding all alternative corridors. Biological resources within this study area were identified and briefly described in tables, texts and figures, based on available data in scientific literature, environmental documents, personal communications from knowledgeable persons, and files from the California and Nevada Bureau of Land Management (BLM). Regional information developed for the study area provided a background and context in which to interpret and evaluate the occurrence and importance of biological resources within the alternative corridors.

Once the regional-level study was completed, a site-specific analysis of biological resources within and adjacent to the alternative corridors was conducted. A two-mile-wide study corridor centered on the proposed routes was examined using recent aerial photography and field reconnaissance of all corridors. Site-specific information was collected, mapped at a large scale, tabulated by milepost, and summarized in this report.

Biological resources examined in this study include the following: (1) native vegetation types; (2) common and characteristic reptiles, amphibians, birds and mammals; (3) legally protected species designated rare, threatened or endangered by the Federal or state governments; (4) species of special interest which have limited legal status, but are of concern to agencies and the interested public; and (5) areas of special biological interest.

The proposed transmission line follows several major existing corridors which have undergone previous environmental review, including the Interstate 15

(I-15) corridor (Woodward-Clyde 1979; BLM 1980a; Jones & Stokes 1980) and the Intermountain Power Project (IPP) corridor (BLM 1979). In addition, biological resources of the California Mojave Desert were studied by BLM to develop the California Desert Conservation Area (CDCA) Plan (BLM 1980b). Detailed background data to develop the CDCA Plan are on file at the California Desert District Office in Riverside (BLM 1984). Under the direction of Mr. Larry Foreman, Desert District Biologist of BLM, these files were used as major data sources for this technical study. As a result of the previous studies and existing BLM data, there is extensive information on biological resources in the study area and, in certain cases, along the specific alternative transmission line corridors.

Field surveys of the proposed transmission line corridors were conducted between November 30 and December 8, 1984 by two 2-man teams of biologists. Each team was equipped with a four-wheel drive vehicle and was able to access all but the most remote, high elevation corridors due to the availability of existing access roads and nearby highways. During the field surveys, all vegetation types within one mile of the centerline were mapped. All important observations of plant and animal life, particularly of legally protected and special interest species, were recorded. All field data were recorded on U.S Geological Survey (USGS) 7.5- and 15-minute quad maps; photographic slides were taken periodically.

Due to the season of the year, biological activity of most animals in the Mojave Desert was low. In addition, spring and summer flowering herbs and grasses were not identifiable. Hence, observation of many species in the field such as desert tortoise, winter bird visitors, most reptiles, and many rare plants were limited. However, the abundance of previous investigations have provided important seasonal data that are mostly sufficient for purposes of an impact assessment. Where additional data are required to complete the impact assessment or refine the mitigation planning, limited field surveys in the appropriate season will be recommended.

STUDY PERSONNEL

This study was directed by Dr. John T. Gray, a Project Ecologist with Dames & Moore, Santa Barbara. Dr. Gray has over five years experience conducting and directing biological inventory, impact assessment, and mitigation planning studies for environmental documents in California and Nevada. Dr. Gray was responsible for the study design, supervising the literature review, developing rare plant information, directing the field survey efforts, summarizing data for the report, and the final compilations, editing and production of the report.

The literature review, field surveys, and initial data summary were conducted by the biological consulting firm Consultants in Wildlife and Environmental Services Agency (CWESA) of Fresno, under subcontract and supervision of Dames & Moore. CWESA staff members involved in the study include Mr. Bill Clark (M.A., biology; wildlife and botany), Mr. Curt Uptain (M.A., biology; wildlife and vegetation), and Mr. Stephen Juarez (B.A., biology; wildlife and

avifauna). CWESA staff has extensive experience in conducting biological field surveys in southern California, including the Mojave Desert. CWESA is currently involved in mitigation surveys along the IPP corridor in Utah and Nevada for rare plants and desert tortoise.

STUDY AREA OVERVIEW

The biological resources of the study area are described in the following sections. The study area includes all alternative routes with an area of approximately 10 miles on either side of each route. Detailed maps of biological resources in the study area are available for review at the Dames & Moore offices in Phoenix, Arizona. Generalized vegetation and wildlife data are shown on Figures 4-2 and 4-3, respectively, in the accompanying Map Volume.

Vegetation Types

The vegetation of the study area can be classified into eight different types, primarily based on the physical characteristics of the vegetation and the dominant plant species. The classification, regional distribution, and dominant species of each vegetation type in the study area are briefly described below. The vegetation types are defined following the classification systems of Cheatham and Haller (1975), Munz (1974), and Thorne (1976) for California; and Beatley (1976), Brown (1982), and Bensen and Darrow (1983) for Nevada.

Data on the species composition and regional distribution of each vegetation type were developed from: (1) a review of scientific literature and desert floras (Munz 1974; Thorne 1976; Beatley 1976; Vasek and Barbour 1977; Brown 1982; Thorne et al. 1981; DeDecker 1984); (2) previous botanical field surveys along the I-15 corridor (Woodward-Clyde 1979; BLM 1980a), IPP corridor (BLM 1979; Schultz 1982, 1984), Ivanpah generating station site (Southern California Edison (SCE) 1981), Barstow-to-Las Vegas motorcycle race course (BLM 1974), and the BLM CDCA Plan (BLM 1980b) and supporting in-house documents (BLM 1984); (3) field reconnaissance surveys of the alternative corridors described previously; and (4) a review of 1981-1984 black-and-white aerial photography (1:10,000 scale) of the alternative corridors.

Common and characteristic plant species of each vegetation type are listed in Table 4-1. The distribution of each vegetation type in the study area was mapped, based on regional vegetation maps from Kuchler (1977) and Brown (1978). The distribution of vegetation types along each corridor is presented later in this report under Biological Resources Within and Adjacent to Alternative Corridors.

Vegetation types consist of associations of plant species that vary with environmental conditions such as soil type, slope, exposure, moisture availability, and disturbance. The study area includes a wide variety of soils, parent material, and topographic features. Although most of the study area is

dominated by a single vegetation type (creosote bush scrub), there are many local vegetation patterns that reflect the complex environmental factors. In most instances, these factors result in a change in species composition, with little or no change in the physiognomy and appearance of the vegetation which is primarily scrub.

Much of the vegetation in the study area has not been disturbed, particularly at higher elevations. However, manmade disturbances occur near Victorville, Barstow, and other built-up areas due to manmade structures, roads, and agricultural activities; the latter is primarily restricted to irrigated farmlands in the Mojave River Valley. Cattle grazing occurs over a large portion of the study area. Off-road vehicle use is regulated on BLM land and is generally restricted to specified roads and recreation areas; however, road scars are scattered throughout the study area, excluding the higher mountain ranges.

Creosote Bush Scrub

Creosote bush scrub occurs on well drained, nonalkaline soils of sandy flats, bajadas, alluvial fans, and upland slopes throughout most of the Mojave Desert. It is absent from dry lake bottoms or valley floors with alkaline or dense soils. It is poorly developed on rocky outcrops, steep slopes, or in washes. Creosote bush scrub generally occurs below 4,000 feet elevation, above which it is replaced by Joshua tree woodland or blackbush scrub.

Creosote bush scrub consists of widely spaced, even-sized creosote bush (Larrea tridentata) shrubs with various smaller shrubs distributed in the open spaces. Burrobush (Ambrosia dumosa) is a small shrub that is a common codominant of creosote bush scrub throughout most of the Mojave Desert. A large number of low-growing winter annual herbs may appear in the spring, depending on the timing and amount of rainfall.

The density and species composition of creosote bush scrub varies considerably throughout the Mojave Desert due to the influence of topography, climate and soils. For example, on lower bajadas with finer soil textures and greater temperature extremes, Larrea may be the only conspicuous plant. On higher bajadas with better cold air drainage and coarser soils with more microsites, a phase of creosote bush scrub occurs that is rich in shrub, cacti, and annual species.

In the central Mojave Desert, creosote bush scrub is mostly codominated by Larrea and burrobush. Other subordinate species include species of saltbush (Atriplex spp.), cheeseweed (Hymenoclea salsola), boxthorn (Lycium andersonii), Mormon tea (Ephedra spp.), cassia (Cassia armata), brittlebush (Encelia farinosa), pencil cholla (Opuntia ramosissima), and Mojave yucca (Yucca schidigera).

In the study area, creosote bush scrub is the dominant vegetation type, occurring on most flats, bajadas, and gentle slopes.

Saltbush Scrub

This vegetation type is often referred to as alkali sink scrub and consists of a low scattered scrub occurring in valley floors, basins, sinks and playas below 4,000 feet elevation. It is dominated by low-growing, scattered halophytic shrubs from the Chenopodiaceae. This vegetation type occupies habitats of extreme aridity, either due to physical factors or soil chemical properties. Two phases of saltbush develop under these conditions. The first is called the xerophytic phase. It occurs on lower bajadas, in basins, or around margins of playas and is characterized by extremely dry soils. It is dominated by shrubs with limited salt tolerance, including all-scale (Atriplex polycarpa), shadscale (A. confertifolia), four-wing saltbush (A. canescens), and desert holly (A. hymenelytra). The halophytic phase is found on saline or alkali soils of playas, in sinks, or near seeps where available surface or ground water is high in mineral content. Important halophytic species in this phase include pickleweed (Allenrolfea occidentalis), glasswort (Salicornia spp.), seep weed (Suaeda spp.), and greasewood (Sarcobatus vermiculatus). Gradients between the halophyte and xerophytic scrub often occur around dry lakes with halophytic species in the playa being replaced by xerophytic species around the margins. Saltbush scrub is generally replaced at higher elevations by creosote bush scrub throughout its range.

The xerophytic phase of saltbush scrub is widely distributed over the Mojave Desert, often forming extensive stands dominated by all-scale, particularly in the western Mojave Desert. Halophytic saltbush scrub is also widely distributed, but restricted to special habitats determined by local topographic conditions.

In the study area, several major basins with dry or existing lakes support both saltbush scrub phases, including Jean Dry Lake, Eldorado Valley, and Hidden Valley in Nevada; and Ivanpah Valley (includes Roach and Ivanpah dry lakes), Soda and Silver lakes and surrounding terrain, Red Pass Lake, East and West Cronese dry lakes, Coyote Lake, Troy Lake, and Harper Dry Lake in California. Extensive areas with scattered stands of xerophytic saltbush scrub dominated by all-scale is present in the western portions of the study area near Kramer Junction.

Blackbush Scrub

Blackbush scrub is a low dark monotonous scrub vegetation dominated by blackbush (Coleogyne ramosissima), sometimes in pure stands. It occurs on shallow, rocky or gravelly, usually calcareous soils of flats, mesas, or relatively steep upper bajadas. Blackbush scrub is found above Joshua tree woodland and creosote bush scrub of California and sagebrush scrub in Nevada. It occurs at elevations of 3,500 to 5,000 feet and is excluded from basins.

Although the species richness of blackbush scrub is lower than creosote bush scrub, it contains many subordinate species whose occurrences vary greatly depending on local environmental conditions. In southern Nevada, common

associated species include Spanish bayonet (Yucca baccata), turpentine bush (Thamnosma montana), hopsage (Grayia spinosa), winter fat (Eurotia lanata), bud sagebrush (Artemisia spinescens), Utah agave (Agave utahensis), and species of Ephedra, Dalea, Atriplex and Tetradymia. In this area, blackbush scrub may contain floristic elements of the Great Basin Desert such as Artemisia spp. and as such is considered a transitional community.

Blackbush scrub occurs in small localized areas at higher elevations in the study area. It is particularly well developed above 7,000 feet in the Clark, McCullough and Spring mountains near the California-Nevada border, and in small areas in the Avawatz, Ord and Granite mountains in the western portion of the study area.

Joshua Tree Woodland

Joshua tree woodland is a scrub vegetation consisting of a Joshua tree (Yucca brevifolia) overstory and an understory of various shrubs and perennial herbs. It is not a true woodland because the trees are widely spaced and contribute only a small percentage of the community biomass. However, the unusual shape and size of Joshua trees makes them a conspicuous element of the vegetation.

Joshua tree woodland occurs on well drained, loose, sandy or gravelly soils of mesas and gentle slopes at 2,500 to 5,000 feet elevation. It typically occurs between creosote bush scrub and pinyon-juniper woodland, grading into each adjacent vegetation type. Joshua tree woodland is found in the same elevational zone as blackbrush scrub; however, the latter usually occurs on rocky, steep slopes with heavier soils. Joshua tree woodland outlines the geographic extent of the Mojave Desert in California and Nevada, occurring along the lower slopes of the bordering mountains, particularly along the western margins and the base of the San Bernardino Mountains. It also occurs in smaller stands at higher elevation slopes within the central portions of the Mojave Desert.

A wide variety of species occurs as an understory to Joshua tree woodland throughout its range, including creosote bush, burrobush, Mormon tea, little-leaf ratany (Krameria parvifolia), cheeseweed, hopsage, boxthorn, blackbush, spinn menodora (Menodora spinescens), Mojave yucca, various cacti such as silver cholla (Opuntia echinocarpa), pencil cholla, and barrel cactus (Ferocactus acanthoides), and several grass species such as galleta grass (Hilaria rigida) and bush muhly (Muhlenbergia porteri).

In the study area, Joshua tree woodlands are well developed between 3,500 and 4,500 feet on the Clark Mountains, portions of Shadow Valley, near Halloran Springs, and in localized areas near the McCullough Mountains. Scattered Joshua trees also occur in the Shadow Mountains, Kramer Hills, and near Adelanto in the western portion of the study area.

Pinyon-Juniper Woodland

Pinyon-juniper woodland is dominated by two small coniferous trees - pinyon pine (Pinus monophylla) and Utah juniper (Juniperus osteosperma) - interspersed with a large number of smaller shrubs, cacti, and perennial herbs. It typically is found above 4,500 feet elevation on rocky or gravelly, well drained soils. It is the dominant high elevation vegetation of the major mountain areas in the Mojave Desert of California and Nevada.

The flora of pinyon-juniper woodland is relatively rich in species compared to lower elevation desert scrub vegetation types. Common species include Great Basin sagebrush (Artemisia tridentata), blackbush, Mormon tea, apache plume (Fallugia paradoxa), matchweed (Gutierrezia sarothrae), boxthorn, grizzly bear cactus (Opuntia erinacea), turpentine bush, and Spanish bayonet.

In the study area, pinyon-juniper woodland is present between 4,800 and 5,200 feet on Clark Mountain, and at similar elevations in the Ivanpah and McCullough mountains.

Desert Wash Scrub and Woodland

This azonal vegetation type consists of the aggregation of plants found in the sandy bottoms of wide canyons, in incised arroyos of upper bajadas, and in the sandy, shallow washes of lower bajadas. It is closely associated with creosote bush scrub and occurs mostly below 4,500 feet elevation. Numerous minor desert washes occur throughout the lower and middle elevations of the Mojave Desert. Most are dominated by cheeseweed, although many other species may also be important, including matchweed, Mormon tea, bladder sage (Salazaria mexicana), desert almond (Prunus fasciculata), and little-leaf ratany. Cheeseweed is the most common and characteristic dominant species of this vegetation type. The occurrence of desert wash scrub in the study area is too localized and widespread to map at a regional scale.

Desert wash woodland (or microphyllous woodland) is poorly developed in the Mojave Desert compared to the Colorado Desert. It consists of dense to sparse thickets along washes with available underlying ground water. The woodland is dominated by one or more of the following species: cat claw (Acacia greggii), desert willow (Chilopsis linearis), palo verde (Cercidium floridum), smoke tree (Dalea spinosa), mesquite (Prosopis glandulosa), and the introduced tamarisk (Tamarix spp.).

Desert willow and mesquite stands in the study area are present at East and West Cronese lakes, near Soda Lake and the Mojave River Wash at Soda Springs, and along selected portions of the Mojave River from near Helendale to Soda Springs.

Riparian Woodland

Riparian woodland consists of a dense thicket of shrubs and trees along a major river or stream channel with significant surface or subsurface flow to support woody vegetation. This vegetation type is only present in the study area along the upper reaches of the Mojave River from south of Victorville to near Helendale, and at Afton Canyon. Dominant plants include willow (Salix spp.), cottonwoods (Populus fremontii), and arrowweed (Pluchia sericea). There is surface water in these portions of the river during much of the year. Other portions of the Mojave River consist of broad sandy wash areas as described above.

Mojave Montane Forest

This vegetation type consists of small, highly restricted groves of Rocky Mountain white fir (Abies concolor ssp. concolor) mixed with Utah juniper and pinyon pine. These open woodlands occur on the steep north-facing slopes and narrow canyons of the higher ranges of the eastern Mojave Desert in California and southern Nevada. In the study area, it occurs in protected areas (65 hectares total) near the crest of Clark Mountain between 5,700 and 7,100 feet elevation. Species richness in this high elevation vegetation type is relatively poor; however, many unusual and highly restricted shrubs and perennial herbs are found in the Mojave montane forest.

Wildlife

Four wildlife habitats can be identified in the study area based primarily on the structure and species composition of the vegetation type which provides food and shelter for resident and migratory wildlife. Each habitat supports a fairly well-defined assemblage of amphibians, reptiles, birds and mammals. However, due to the overlapping distribution of habitat types and the highly mobile nature of birds and mammals, there is considerable overlap in the distribution of wildlife species between each habitat.

In the following sections, the common and characteristic wildlife species in each habitat type are briefly described and are also listed in Table 4-2 (amphibians and reptiles), Table 4-3 (birds), and Table 4-4 (mammals). This information was developed primarily from scientific literature (BLM 1978; Karl 1984), natural history guides (Ingles 1965; Stebbins 1966; Garrett and Dunn 1981), extensive sighting records from the California and Nevada BLM files (BLM 1984), experience of the study team biologists, and observations during the field surveys.

Scrub Habitat

The habitat consists of relatively homogenous, low-growing shrubs with or without interspersed herbs and grasses. It includes the vegetation types of

creosote bush, saltbush, and blackbush scrub. As a result, it occurs over a wide elevational range and in a variety of topographic situations throughout the study area. The habitat is relatively simple, with one or two dominant plant species, low cover, and scarce water resources. It is the most common wildlife habitat in the study area.

Amphibians and Reptiles

Amphibians are relatively scarce in the scrub habitat due to the absence of water. Couch's spadefoot toad may be found in some areas. It is a nocturnal species that is active after rains. During dry periods, it buries itself in loose soil or in rodent burrows.

Reptiles are more common in the scrub habitat, particularly lizards. Typical species include the western whiptail, side-blotched lizard, long-nosed leopard lizard, desert iguana, and desert horned lizard. Lizards in rocky areas of scrub habitat include the chuckwalla, collared lizard, and banded gecko. The desert night lizard is likely to occur under Joshua trees.

Common snakes of the scrub habitat include the glossy snake, gopher snake, coachwhip or red racer, and shovelnose snake. Less common but characteristic snakes of the scrub habitat are the rosy boa, patchnose snake, and night snake.

A distinctive snake of the scrub habitat is the sidewinder. This rattlesnake species is venomous; however, it is not as toxic as its close relative, the Mojave green rattlesnake. The latter species, which also occurs in scrub habitat, is larger than the sidewinder and is capable of inflicting a bite that causes more respiratory distress than any other North American pit viper.

Birds

In general, the abundance and diversity of the scrub habitat avifauna are low. Birds are more common where the vegetation is profuse and in desert wash woodlands. Roadrunners are a common resident bird in the scrub habitat that hunt reptiles in the early morning. The most conspicuous bird is the common raven which nests in cliffs and hunts for prey and carrion over the broad desert expanses. Common resident songbirds include the horned lark, black-throated sparrow, and cactus wren. Costa's hummingbirds are a relatively common nesting summer visitor. Common birds of prey include the red-tailed hawk, golden eagle, and American kestrel.

Mammals

Most mammals of the scrub habitat are nocturnal as a means of water conservation. Two notable exceptions are the common dwarf white-tailed antelope squirrel and black-tailed jackrabbit. Typical nocturnal mammals

include the Merriam's kangaroo rat, little pocket mouse, desert woodrat, and southern grasshopper mouse.

Woodland Habitat

This habitat consists of widely scattered trees with a sparse understory of shrubs, grasses and herbs. It includes the vegetation types of Joshua tree woodland and pinyon-juniper woodland. The former occurs throughout the study area, including scattered pockets of the western study area, in Shadow Valley, on Clark Mountain, and in the McCullough Mountains. Pinyon-juniper woodland is primarily restricted to the higher elevations of the Clark, Ivanpah, and south McCullough mountains. The woodland habitat provides greater food and structural complexity than the scrub habitat.

Amphibians and Reptiles

As in the scrub habitat, amphibians are relatively scarce due to the lack of significant surface water resources. Typical lizard species include the desert spiny lizard, desert night lizard, and Gilbert's skink. Common snakes are the desert night snake, regal ringneck snake, and Mojave green rattlesnake.

Birds

The woodland habitat supports a relatively abundant and varied avifauna due to the presence of trees and a diverse understory. Joshua trees are used for nests by Scott's orioles, cactus wrens, ash-throated flycatchers, and ladder-back woodpeckers. Other common birds of the woodland habitat include the scrub jay, red-shafted flicker, common poor-will, black-tailed gnatcatcher, western bluebird, pinyon jay, and mountain quail. Resident raptors nest in cliffs in or near the woodland habitat, including the raven, red-tailed hawk, golden eagle, and prairie falcon.

Mammals

The Yuma myotis and big brown bat are common bat species that utilize caves and rocky terrain in the woodland habitat. The Panamint kangaroo rat, canyon mouse, and pinyon mouse are common, mostly in pinyon-juniper woodlands. Common carnivores include the coyote, gray fox, and bobcat. Other important mammals include the ubiquitous mule deer and desert cottontail, as well as the less common desert bighorn sheep.

Riparian and Wash Habitats

This habitat consists of sparse to densely vegetated thickets and woodlands along stream channels. Well-developed riparian woodland is present along the

upper Mojave River from near Victorville to Helendale. This habitat is dominated by cottonwoods, willows and tamarisks. Along the lower Mojave River and in other areas of the Mojave Desert with seasonally standing or flowing water are desert wash woodland habitats. These are small localized areas dominated by mesquite, cat-claw, or desert willow. Both riparian and desert wash woodland habitats provide important habitat for a diverse and abundant herptofauna and avifauna due to its structural complexity, abundant cover and food, and availability of water.

Amphibians and Reptiles

Common amphibians along riparian and desert wash woodland areas include the leopard frog, red-spotted toad, and bullfrog. The desert spiny lizard commonly takes refuge in riparian habitats of the Mojave Desert. Zebra-tailed lizards and the Mojave fringe-toad lizards are also common residents of these habitats, particularly along sandy washes. Common snake species in these habitats include the rosy boa and western blind snake.

Birds

The riparian woodland habitat along the Mojave River and at Afton Canyon, as well as the scattered mesquite and desert willow thickets in the study area, support an abundant and diverse avifauna. A variety of resident, migratory, and winter visitor bird species occur in these habitats, such as the belted kingfisher, downy woodpecker, northern flicker, western flycatcher, black phoebe, western kingbird, bushtit, orange-crowned warbler, yellow-rumped warbler, western tanager, and lesser goldfinch, among others. Low-lying areas with seasonal standing water, such as at Harper Dry Lake, East Cronese Lake, and Soda Lake, provide a temporary stopover habitat for migratory waterfowl such as the black-bellied plover, willet, and western sandpiper.

Mammals

A variety of bats may be found in the riparian woodland habitat, including the fringe-tailed myotis. Virginia opossums, striped skunks, and the western harvest mouse are common mammals of these habitats.

Montane Forest Habitat

This habitat consists of a relatively small, isolated coniferous forest near the crest of Clark Mountain where pinyon-juniper woodland is mixed with a residual stand of Rocky Mountain white fir. This high elevation forest habitat provides year-round and seasonal habitat for a variety of wildlife. However, the small areal extent of this habitat limits the abundance of wildlife species.

Amphibians and Reptiles

Characteristic species of this habitat include Gilbert's skink and the western skink. The pine-gopher snake is the most common snake in this habitat.

Birds

Common birds of the Montane forest habitat include the western wood peewee, plain titmouse, bushtit, and scrub jay. Cooper's hawk is a common summer resident in this habitat. Other bird species commonly observed are the white-throated swift, mountain quail, and turkey vulture.

Mammals

Deer mice occur in this habitat as well as in lower elevation habitats. Other common mammals include the Panamint chipmunk, pinyon mouse, and mule deer. Several bats reside in this habitat, including the fringed myotis, big brown bat, and spotted bat.

Legally Protected Species

Plants

Legally protected plants include species officially listed as rare and endangered by the California state government (California Department of Fish and Game (CDFG) 1984), critically endangered by the Nevada state government (Nevada Division of Forestry (NDF) 1984), or threatened and endangered by the Federal government (U.S. Fish and Wildlife Service (USFWS) 1984). The potential occurrence of these species in the study area was evaluated by a review of their regional distribution using regional floras (Munz 1974; Beatley 1976; Thorne et al. 1981; DeDecker 1984); California BLM data in the CDCA Plan (BLM 1980b) and supporting in-house data (BLM 1984; Albiston 1984); Nevada BLM data (Fisher 1980; Driver 1984); data from the California Native Plant Society (CNPS) and California Natural Diversity Data Base (CNDDB) 1984; data from the NDF (Murphy 1984); data from the Northern Nevada Native Plant Society (NNPS) (Mozingo and Williams 1980; Pinzl 1984); and previous botanical investigations along the I-15 and IPP transmission line corridors (BLM 1979; Woodward-Clyde 1979; BLM 1980a; Jones & Stokes 1980; Schultz 1982, 1984).

Based on these investigations, no legally protected plant species are known or likely to occur within the study area.

Fish and Wildlife

Legally protected fish and wildlife include species listed as rare or endangered by the California and Nevada state governments (CDFG 1980, 1984; Nevada Department of Wildlife (NDW) 1984), or threatened and endangered by the Federal government (USFWS 1984).

Several wildlife species described below are listed as rare or endangered by the California or Nevada state governments, but do not have Federal status as a legally protected species. However, these species may be listed as candidates for Federal listing. Federal candidate species are assigned to one of two categories depending on the current state of knowledge of the species and its biological appropriateness for listing (USFWS 1980, 1983). Category 1 includes species for which the USFWS currently has on file substantial information on biological vulnerability and threats to support the appropriateness of proposing to list as endangered or threatened species. Category 2 includes species for which sufficient information is available to indicate possible listing, but for which additional data are required on vulnerability and threats. Species in these categories are usually given protection by BLM under the Endangered Species Act, pending the resolution of their official status (BLM 1980b).

Mohave Tui Chub

The Mohave tui chub (Gila bicolor mohavensis) is designated an endangered species by the California state government (CDFG 1980) and Federal government (USFWS 1984). This species represents the only native fish of the Mojave River in California. It originated from a large Pleistocene lake and river system along the Mojave River drainage, becoming restricted to the upper Mojave River and nearby tributaries as these water systems receded. In the 1930s, arroyo chubs (Gila orcutti) from coastal California were introduced into the headwaters of the Mojave River and hybridized with the Mohave tui chub (Hubbs and Miller 1943). By 1970, genetically pure Mohave tui chubs had been eliminated from the Mojave River by hybridization and subsequent introgression. However, a small population of Mohave tui chub has persisted in isolated ponds at Soda Springs. Mohave tui chub have subsequently been introduced into seeps and marshlands on the China Lake Naval Weapons Center and into a small pond near the town of Hinkley.

Gila Monster

The Gila monster (Heloderma suspectum) is designated a rare species by the Nevada state government (NDW 1984) and a candidate (Category 2) for Federal listing (USFWS 1982). It is a relatively uncommon large venomous lizard that occurs primarily in Arizona and northern Mexico and in portions of Nevada, California, Utah, and New Mexico (Stebbins 1966). It is diurnal in the moderate temperatures of spring and early summer, nocturnal in midsummer, and dormant in the winter. Gila monsters have an unusual and irregular color

pattern for concealment. They feed on bird and reptile eggs and small rodents. Gila monsters utilize the burrows of other animals, dig their own, or retreat to natural cavities. Gila monsters are most common on the lower slopes of desert mountains below 4,000 feet elevation, ranging into the outwash plains (Larson 1977). Preferred local habitats consist of rocky or sandy washes, canyons, and stream beds with seasonal surface water (Cross 1978).

The Gila monster is only known from two single sighting records in California: on the western slopes of the Providence Mountains and seven miles southwest of the California-Nevada border in the Clark Mountains (McGurty 1977). The latter location occurs in the study area. Suitable habitat appears to occur in the Clark Mountains such as in the lower slopes, bajadas, canyons and passes. However, surveys of suitable habitat on Clark Mountain (including the Keany Pass area) by Mitchell (1978), and the eastern Mojave Desert by McGurty (1977) failed to discover any Gila monsters. Hence, the occurrence of Gila monsters in the Clark Mountains in the study area is expected to be extremely rare. McGurty (1977) has suggested that Gila monsters in California may represent either remnant relictual populations, recent invaders, or released captives.

In southern Nevada, Gila monsters are more common, particularly along the Colorado River and in the Las Vegas, Mojave and Virgin river valleys (Bradley and Deacon 1965). Although there are no published records of Gila monsters in the Nevada portions of the study area, suitable habitat occurs in the washes of the McCullough Mountains and surrounding smaller ranges (Burr 1981; Price 1984).

Desert Tortoise

The desert tortoise (*Gopherus agassizii*) is designated a rare species by the Nevada state government (NDW 1984), a candidate (Category 2) for Federal listing (USFWS 1982), a California BLM sensitive species (BLM 1980), and has been considered for listing as a state rare species in the last several years (Brode 1984). The CDFG Code (Section 5000) prohibits the taking or sale of the desert tortoise in California.

The desert tortoise occurs in the deserts of California, southern Nevada, southwestern Utah, western Arizona, and northern Mexico. The greatest abundance of desert tortoise is in the Mojave Desert of California (Lukenbach 1982). It now occupies a smaller range than previously and population densities apparently have been reduced within its current range due to habitat loss, human activities, and illegal collecting.

The desert tortoise is primarily active during the early spring and fall, less active in the summer, and dormant in the winter. In March or April, tortoise emerge from burrows and are active most of the day. During the summer, tortoise have a bimodal activity pattern to escape the hottest time of day or they may remain dormant in their burrows to reduce heat stress and water

loss. Peak activity occurs in the spring when temperatures are mild and food is abundant. In the fall, males may be active, particularly with courtship and mating behavior. From October to February, tortoises are dormant in their burrows. Peak breeding activity occurs in the spring (March to June). Egg laying occurs in spring and summer; hatchlings emerge in the fall but leave burrows very infrequently.

The desert tortoise is herbivorous and feeds primarily on the foliage and flowers of spring annuals. Common plant genera include locoweeds (*Astragalus* spp.), Gilas (*Gilia* spp.), blazing stars (*Mentzelia* spp.), desert dandelions (*Malacothrix* spp.), and phacelias (*Phacelia* spp.). In the dry springs and during the summer, desert tortoise rely more on perennial grasses and dried herbs. Infrequent summer rains provide a second blossom of annual herbs for the fall activity period. Tortoises subsist on their stored fat when foods are not available as well as during their summer aestivation and winter dormancy periods.

Burrows are important to desert tortoises because most of their time is spent underground to regulate heat and water loss, and to escape predation. The depth and type of burrow used varies depending on seasonal temperatures and soil type. In the Mojave Desert of California and southern Nevada, the deep permanent burrows used for periods of dormancy are 1 to 3 meters deep at an angle of 10 to 30°. During the spring and summer, as well as in the fall activity period, tortoise use shallower burrows situated on flats, interwash areas, and often under shrubs. Desert tortoise utilize several different shallow burrows during the summer. They may also seek cover in a small depression (pallet) under a shrub.

The home range of the desert tortoise varies according to the geographic location, food supply, and population density. It varies in size from 10 to 100 acres in Utah to 96 to 190 acres in the western Mojave Desert of California (Woodbury and Hardy 1948; Berry 1974). In southern Nevada, Burge (1977) found the mean home range size of males and females to be 48 to 49 and 28 to 67 acres, respectively. Activities of hatchlings and juveniles are restricted closely to the burrows.

Desert tortoise occur in three vegetation types in the Mojave Desert of California and Nevada, including creosote bush scrub, Joshua tree woodland and, to a lesser extent, saltbush scrub. Desert tortoise are usually excluded from dry lake beds and unvegetated alkaline basins. Although no single soil type is preferred, it must be friable enough for digging yet firm enough to support the burrow. Soil types in desert tortoise habitat of western Mojave Desert range from gravelly flats in Stoddard Valley to sandy soils in Fremont Valley and windblown sands near Hinkley (Lukenbach 1982). Desert tortoise are generally uncommon above 3,500 feet elevation and in rocky areas, such as on upper bajadas and rugged mountain slopes.

Data on the distribution and estimated densities of desert tortoise in the study area were primarily gathered from California and Nevada BLM maps and files and a review by Lukenbach (1982) of the desert tortoise in California. In

general, desert tortoise are distributed in suitable habitat throughout most of the study area. Populations are scattered and vary in size and number based on local habitat conditions. Relatively high densities (100 to 250 or more individuals per square mile) of tortoise occur in the western portion of the study area, particularly in and near Stoddard Valley, along Highway 395, west and north of Harper Dry Lake, and west of the Calico Mountains. These areas represent some of the most important remaining desert tortoise habitat in California (BLM 1980).

Desert tortoise abundance is lower in the central and eastern portion of the study area due, in part, to greater elevations and topographic features. Areas of moderate density (40 to 155 individuals per square mile) occur in the alluvial plains north and south of Baker, in the Shadow Valley, surrounding Ivanpah Dry Lake, and in the northern Ivanpah Valley of Nevada. Relatively high density (more than 155 individuals per square mile) populations occur directly south of Baker and in the southern Ivanpah Valley. The latter area also represents one of the most important desert tortoise habitats in the eastern Mojave Desert (Lukenbach 1982).

Swainson's Hawk

The Swainson's hawk (Buteo swainsoni) is designated a rare species by the California state government (CDFG 1983), a protected species by the Nevada state government (NDW 1984), and a candidate (Category 2) for Federal listing (USFWS 1982). This species formerly occurred throughout most of California and Nevada, excluding the high mountain ranges. However, since the 1940s, there has been a severe decline in its population in both states. It is a summer resident breeder that generally nests in large trees in open plains, grasslands, and flat terrain. Cottonwood, sycamore, or willow trees are the most frequently used nesting sites. Currently, Swainson's hawk breeds in the high desert regions of northeastern California and northern Nevada, sparingly in the Central Valley, and south in the coastal woodlands from Marin County to Santa Barbara County (Bloom 1979).

Swainson's hawk is a scarce breeder in the Mojave Desert due to the lack of well developed habitat (Grinnell and Miller 1944). Nests of Swainson's hawk in the desert are generally restricted to Joshua trees and/or riparian woodlands dominated by cottonwoods. Historical nesting records in the study area occur near Adelanto and Victorville. The most recent nesting records near the study area were in 1978 at Cima Dome and Lanfair Valley (Bloom 1979). Suitable habitat appears to occur in the study area along the Mojave River Valley from Victorville to Helendale where large willows and cottonwoods occur, and in areas of Joshua tree woodland near Adelanto, Shadow Valley, Clark Mountains, and Beer Bottle Pass. However, the occurrence of nesting Swainson's hawks in the study area is expected to be rare to infrequent. Migrating Swainson's hawks are likely to occur for brief periods of time throughout much of the study area each year.

California Yellow-billed Cuckoo

The California yellow-billed cuckoo (Coccyzus americanus occidentalis) is designated a rare species by the California state government (CDFG 1980) and a candidate (Category 2) for Federal listing (USFWS 1982). It is a summer resident breeder in California that historically nested along rivers and streams in the coastal valleys from Mexico to San Francisco, in the Central Valley, along the Colorado River, and at small isolated locations elsewhere. Based on a comprehensive statewide survey conducted by Gaines (1977) and Gaines and Laymon (1984), the nesting range of the yellow-billed cuckoo has been reduced, due to the loss of riparian habitat throughout California, to Sacramento Valley, Santa Maria River, Owens Valley, Amargosa River, and the Colorado River. Yellow-billed cuckoos nest in dense riparian thickets dominated by willows and other woody riparian species such as cottonwoods. Other habitat features include dense understory, high humidity, surface water, at least 120 meters of riparian foraging habitat, with at least 10 hectares of continuous vegetation for nesting.

Yellow-billed cuckoos have never been numerous in the Mojave Desert (with the exception of the Colorado River populations) due to the lack of suitable habitat. Only two sightings have occurred in the study area: at Yermo in 1910 and in the upper Mojave River Narrows in 1978 (Mulder 1978; Gaines and Laymon 1984). Neither sighting was of a confirmed nesting individual. There appears to be only two suitable habitat areas in the study area. The upper and lower Mojave River Narrows near Victorville contain extensive willow and cottonwood woodlands and may support occasional nesting birds, although no birds have ever been recorded at this location (Mulder 1984). Riparian habitat north of Victorville is of poor quality due to the smaller and scattered stands of willows and lack of standing water. Afton Canyon may also provide suitable habitat for infrequent breeding yellow-billed cuckoos, although no birds have ever been recorded at this location (Gaines and Laymon 1984).

Migrants and transients may occur along the Mojave River in riparian thickets of dense willow, mesquite, and/or cottonwood from near Victorville to Soda Springs. However, their occurrence would be rare and highly transient.

Least Bell's Vireo

The least bell's vireo (Vireo bellii pusillus) is designated an endangered species by the California state government (CDFG 1980) and a candidate (Category 1) for Federal listing (USFWS 1982). It is a summer resident breeder in California from March to August that nests in riparian habitats. Historically, this species was very common in the Central Valley, the interior valleys of the central and southern coast, and in desert oasis and riparian woodlands of the Owen Valley, Death Valley, and Mojave River Valley (Goldwasser 1978; Goldwasser et al. 1980). Least bell's vireo nest in low dense riparian thickets, usually dominated by willows with a well developed understory.

The least bell's vireo is probably extirpated as a breeding species in the Central Valley, Owens Valley, and coastal valleys north of Santa Barbara County although some suitable habitat remains unused in these areas. Major reasons for the statewide decline in distribution and abundance include loss of riparian habitat, predation, and nest parasitism by the brown-headed cowbird. Recent statewide surveys by Goldwasser (1978) and Goldwasser et al. (1980) indicate that least bell's vireos are not known to nest in the study area, nor have there been any sightings in the study area since at least 1970. The nearest recent records are of nesting populations in Little Morongo Canyon and Big Morongo Wildlife Reserve in southeastern San Bernardino County (Goldwasser 1978). Hence, it is unlikely that least bell's vireo now nest in the study area, although the study area is within its historic range and contains suitable habitat. Occurrence of the least bell's vireo in the study area would be expected to be very rare and of migrant individuals only.

Mohave Ground Squirrel

The Mohave ground squirrel (Spermophilus mojavensis) is designated a rare species by the California state government (CDFG 1980). It has no Federal status. This species occurs only on a restricted portion of the Mojave Desert, from Olancho in Inyo County south to Victorville, and from the Tehachapi Mountains in Kern County east to the Granite Mountains in San Bernardino County (Hall and Kelson 1959; Wessman 1977). A recent survey by Wessman (1977) indicated that the Mohave ground squirrel occurs 40 miles east of the former known eastern margin of its range in the vicinity of Fort Irwin and the Avawatz Mountains. Although this range extension added about 1,800 square miles to the known Mohave ground squirrel habitat, Wessman (1977) found evidence to suggest the loss and/or displacement of this species in the Lucerne Valley.

The Mohave ground squirrel occurs on a wide variety of habitat types and has been trapped in saltbush scrub, creosote bush scrub, shadscale scrub, sandy creosote bush scrub, and Joshua tree woodland. Within its known range, this species is widespread, but relatively uncommon. It is only absent from rocky, steep slopes and dry lake playas. Areas of high concentration appear to occur in portions of Inyo County and in the Superior Valley/Lane Mountain area of San Bernardino County (BLM 1979). Mohave ground squirrel habitat mostly occurs between 1,500 and 5,000 feet (Wessman 1977).

The Mohave ground squirrel shares much of its range with the white-tailed antelope squirrel (Ammospermophilus leucurus), and Beechy ground squirrel (Spermophilus beecheyi) near agricultural and urbanized areas. The Mohave ground squirrel is closely related to the round-tailed ground squirrel (Spermophilus tereticaudus) which occurs in low elevation habitats with fine windblown sand. The entire eastern margins of the Mohave ground squirrel range is bordered by the range of the round-tailed ground squirrel. Wessman (1977) found overlap in the range of these two species at the Tiefert Mountains, Coyote Cry Lake, and Helendale. Hybridization may have occurred in the Helendale area.

The Mohave ground squirrel is diurnal and active from March through July (Bartholemew and Hudson 1961; Zembel and Gall 1980). During the rest of the year it remains in underground burrows in a state of torpor. The Mohave ground squirrel eats on a variety of green vegetation.

The Mohave ground squirrel occurs in the western portion of the study area, west of the Mojave River from Victorville to Helendale, then north and east to Lane Mountain and the Tiefort Mountains. North of Helendale, the Mohave ground squirrel does not live on, or cross, the Mojave River and its broad wash areas (Wessman 1977). It also appears that it is no longer found east of the river between Victorville and Helendale in its historic range (BLM 1980). In the westernmost portions of the study area, it is expected to be widespread but infrequently encountered in all but the most rocky, upland habitats.

Species of Special Interest

Species of special interest include plants and animals that are of concern to agency personnel, conservationists and biologists because they: (1) appear to be of limited distribution; (2) are experiencing population declines; (3) are vulnerable to potential threats to their existence, or (4) are of unusual scientific, educational or recreational value. These species usually do not have the same rarity or vulnerability as legally protected species; hence, many do not have the same legal protection afforded rare, endangered or threatened species. However, most are protected under various Federal or state regulations. In addition, it is possible that if present trends continue or worsen, many of these species may be added to official Federal and state rare species in the future.

There are two general categories of special interest species: (1) species that are candidates for official Federal listing as threatened or endangered (USFWS 1980, 1982, 1983); and (2) species designated as sensitive species by BLM. Many species which have Federal candidacy status also have been identified as species of special interest by private conservation groups such as the Audubon Society, CNPS (Smith and York 1984) and NNPS (Mozingo and Williams 1980; NNPS 1984).

Federal candidate species are assigned to one of two categories depending on the current state of knowledge of the species and its biological appropriateness for listing (USFWS 1980, 1983). Category 1 represents species for which sufficient information is available to support the biological appropriateness for official listing as threatened or endangered. Category 2 includes species for which sufficient information is available to indicate possible listing, but for which additional data are required on vulnerability and threats. Species in these categories are usually given protection by BLM under the Endangered Species Act, pending the resolution of their official status (BLM 1980b).

BLM in California has developed sensitive species lists for plants and animals (BLM 1980b, 1984). These species are all Federal candidate species (Categories 1 and 2) and are managed accordingly. Nevada BLM has not developed a

sensitive species list but utilizes, at a minimum, the species on the Federal candidate list (Categories 1 and 2) (Driver 1984; Price 1984).

Several private conservation organizations have developed unofficial lists of special interest species. These categories are briefly described below.

Plants

Plants of special interest in the study area are defined to include (1) Federal candidate species, Categories 1 and 2; and (2) California BLM sensitive species. The Federal candidate list for plants was initially published in 1980 (USFWS 1980). In 1983, a revised list was published based on new information on the rarity and vulnerability of the 1980 candidate species (USFWS 1983). As a result, many species were placed in new categories, including a large number that were removed from candidate status (i.e., placed in Category 3C).

California BLM has developed a list of sensitive species (BLM 1984) that are afforded special protection and management on BLM lands. These species are all Category 1 or 2 Federal candidate species; however, some Federal candidate species which occur on BLM land have not been added to the sensitive species list. Both BLM sensitive species and Federal candidate species are given protection under the Endangered Species Act with regard to actions that may adversely affect these species (BLM 1980b). Plants on the California BLM sensitive species list, as well as Category 1 and 2 Federal candidates in California, were reviewed for their potential occurrence in the study area.

The CNPS has produced several rare plant inventories since 1974. This organization of private botanists and conservationists has developed an extensive data base on rare plants in California in cooperation with state agencies. The 1980 inventory (Smith et al. 1980) was the first comprehensive report; two addendums were issued since 1980 (Smith et al. 1981, 1982). In 1984, a new CNPS inventory (Smith and York 1984) was prepared with new categories (lists). Many species were moved to different lists based on new information on their distribution and vulnerability. The 1980 Lists 1 and 2 are now designated List 1A (presumed extinct) and List 1B (rare or rare and endangered) in the 1984 inventory. The CNPS status of Federal candidates and/or BLM sensitive species in the study area were reviewed for this project.

Nevada BLM has not developed a sensitive species list; however, it recognizes all Category 1 and 2 Federal candidate species as species requiring protection and special management (Driver 1984). In addition, Nevada BLM may use the recommendations of the NNPS concerning the priority given to candidate species on BLM lands.

NNPS is a private conservation organization similar to CNPS. A comprehensive inventory of species of special interest in Nevada was produced by NNPS in 1980 (Mozingo and Williams 1980). This inventory is updated each year at NNPS workshops attended by private botanists and agency personnel.

The 1984 lists of special interest species (NNPS 1984) include three categories in decreasing order of importance: (1) sensitive species, (2) watch list, and (3) other rare species. Sensitive species include Federally listed plants, many Category 1 and 2 candidate plants, and any other species that have similar rarity or vulnerability as determined by NNPS. The watch list contains some candidate species as well as many ex-candidate species; these species are not considered to face immediate or serious threats. The other rare list contains remaining ex-candidate species. The NNPS status of Federal candidate species in the study area was reviewed for this project.

A preliminary list of special interest plants that may occur in the central portions of Mojave Desert of California and southern Nevada was developed by a review of several regional floras (Munz 1984; Beatley 1976; Thorne et al. 1981; DeDecker 1984). This list was refined to include only species that could potentially occur in the study area, either within or adjacent to the proposed corridors, using California BLM data in the CDCA Plan (BLM 1980b) and supporting in-house data (BLM 1984; Albiston 1984), Nevada BLM data (Fisher 1980; Driver 1984), data from the CNDDDB and CNPS (CNDDDB 1984), data from NNPS (Mozingo and Williams 1980; Pinzl 1984), and previous botanical investigations along the I-15 and IPP transmission line corridors (BLM 1979; Woodward-Clyde 1979; BLM 1980a; Jones & Stokes 1980; Schultz 1982, 1984).

Based on these investigations, it was established that 10 species of special interest are known to occur in the study area and could occur near or within alternative corridors. The regional distribution and occurrence of these species in the study area are summarized on Table 4-5 and briefly described below.

Species that were deleted from the preliminary list of potential special-interest species are presented in Table A-1 in Appendix A. These species were not considered further in the site-specific analyses for this project for one or more of the following reasons: (1) they have been recently (1983) removed from the Federal candidate list (assigned to Category 3C); (2) they are not known to occur in the study area; and/or (3) they may occur in the study area but are not expected to occur within or adjacent to the alternative corridors.

Astragalus jaegerinus (Jaeger milkvetch)

This species is a perennial herb restricted to a 10-square-mile area on, or partially on, Coolgardie Mesa between Barstow and Goldstone in the south-central portion of the study area. It occurs in creosote bush scrub and is commonly associated with Joshua trees. This species is found on dry stony hillsides and desert mesas in granite sand and gravel at elevations of 300 to 3,000 feet.

Chorizanthe spinosa (Mojave spineflower)

This species is an annual herb that occurs in scattered but widespread areas of the western Mojave Desert in Kern and San Bernardino counties. It is found on sandy washes and flats of creosote bush scrub and Joshua tree woodland at elevations of 2,500 to 3,500 feet. Like most desert winter annuals, it does not appear conspicuously each year. Potential habitat for these species occurs throughout the western end of the study area where populations have been found near Kramer Junction, Harper Dry Lake, and Adelanto.

Coryphantha vivipara var. alversonii (Foxtail cactus)

This cactus species occurs over a wide area of the Colorado Desert and western and southern Mojave Desert, ranging from the Little San Bernardino Mountains to the Chuckwalla Mountains. It occurs on steep stony slopes of creosote bush scrub and Joshua tree woodland at elevations of 2,000 to 5,000 feet. It is expected to be very rare in the study area. The only known record of this species in the study area is at Clark Mountain Station.

Cymopterus deserticola (Desert cymopterus)

This species is a perennial herb that appears to occur very infrequently in the western Mojave Desert near Victorville, Edwards Air Force Base, and Kramer Junction. It occurs as small isolated populations on loose sandy soil of flats in old dune areas. It is found in creosote bush scrub or Joshua tree woodland at elevations of 2,300 to 4,300 feet. Only two populations have been recorded in the study area: near Kramer Junction and in Apple Valley. Other small, localized populations could potentially occur in the western portions of the study area.

Eriophyllum mohavensis (Barstow woolly sunflower)

This species is only known from four locations, all of which occur in the study area: (1) Iverson Ranch near Barstow, (2) adjacent to Mojave River across from the railroad station in Barstow, (3) scattered populations five to eight miles northeast of Kramer Junction, and (4) an unconfirmed location midway between Barstow and Kramer Junction. It is a small woolly annual herb that occurs in fine sandy or gravelly areas of varying slope at 2,000 to 3,000 feet elevation. The populations northeast of Kramer Junction are the largest known to date. A one-half section of land containing this species in this area has been declared an Area of Critical Environmental Concern (ACEC) by BLM. It is possible that additional small isolated populations could occur in the western portion of the study area.

Penstemon bicolor ssp. bicolor and roseus (Bicolored penstemon)

These subspecies are tall perennial herbs that have different flower colors. Both are restricted to mountain ranges in southern Nevada and co-occur throughout much of their respective ranges. Subspecies bicolor is only known from the east slope of the Spring Mountains in the eastern portion of the study area. It occurs in creosote bush scrub or washes, roadsides, disturbed sites, and limestone bedrock exposures at elevations of 2,000 to 5,000 feet. Subspecies roseus is more widely distributed in the Eldorado, McCullough, and Arrow Canyon mountains in similar habitats and elevations.

Phacelia anelsonii (Nelson's phacelia)

This annual species is widely distributed throughout portions of California (northeastern Mojave), Nevada (southern areas), and Utah. It is commonly found in disturbed and rocky soils between 4,000 and 5,500 feet with blackbush scrub and creosote bush scrub. It is also found in washes, on scree slopes, and talus areas. Populations are usually very small and very rare. It is known to occur in at least one site in the McCullough Mountains, and may occur in the Clark Mountains.

Puccinellia parishii (Parish's alkali grass)

This annual grass is very rare in California, being restricted to alkali seeps. It was originally collected in the Lucerne Valley. A recent collection of this species, or species closely resembling it, was made near Kramer Junction. Its occurrence in the study area is likely to be extremely rare, in highly localized sites, and only in the western portions of the study area.

Sclerocactus polyancistrus (Mojave fishhook cactus)

This short globular-shaped cactus is widely distributed from California to Nevada (Nye County only) and Arizona. In California, it occurs in small, isolated populations with widely scattered individuals from Red Rock Canyon to Inyo County. It occurs from 2,500 to 6,500 feet elevation. It is often collected illegally. There are several known populations of this species in the western portion of the study area.

Sphaeraleca rusbyi ssp. erimicola (Rusby's desert mallow)

This is a very rare perennial herb that mostly occurs in the Panamint Mountains of Inyo County in creosote bush scrub and blackbush scrub at elevations of 3,000 to 4,500 feet. It also has been recorded in at least one location in the Clark Mountains in the study area.

Wildlife

Wildlife species of special interest in the study area are defined as species which are not officially listed as rare, threatened or endangered, but have one or more of the following designations: (1) Federal candidate species, Categories 1 and 2 (USFWS 1982); (2) California BLM sensitive species (BLM 1980); and/or (3) protected species according to the administrative codes of the CDFG (1984), or NDW (1984).

Of the following three special interest species, only one is a Federal candidate species (ferruginous hawk), and only one is a BLM sensitive species (desert bighorn sheep).

Golden Eagle

The golden eagle (Aquila chrysaetos) is a fully protected bird according to the CDFG (Section 3511) and NDW (1984) codes which prohibit the hunting and harassment of this species. Golden eagles are also on the third priority list developed by Remsen (1978) for CDFG, describing bird species of special concern in California. This list includes species without legal status that appear to be suffering population declines and habitat loss, or are particularly vulnerable to known or future threats. Species on the third priority list are not in any present danger of extirpation and their populations, including those of the golden eagle, are not declining; however, by virtue of their small populations in California, they are potentially vulnerable to future threats (Remsen 1978).

The golden eagle is a resident of the mountainous regions of the western United States. In the Mojave Desert of California and southern Nevada, it is a relatively uncommon resident inhabiting numerous isolated mountain ranges with suitable nesting habitat. The rocky terrain of these desert mountains provides appropriate nesting substrates on cliffs overlooking broad desert valleys and plains. In a survey of nesting sites in California, Thelander (1974) located 25 pairs of nesting eagles in the Mojave and Colorado deserts. Although golden eagles nest throughout California and Nevada, they are mostly located in undeveloped, remote mountainous areas.

Golden eagles are most commonly observed in broad open habitats where their dominant prey item, rabbits, are readily found. They hunt over a broad elevation range in many vegetation types, with the exception of dense forests and scrub vegetation. Home range size varies from 20 to 50 square miles. Nests are usually located on remote, steep cliffs overlooking open foraging habitat. Nests may occasionally be established in large transmission line towers and trees. Nesting sites are generally isolated from human activities and disturbances. Although breeding territories are established, nest sites may be located within several miles of one another. Golden eagles often must compete with other raptors such as prairie falcons and American kestrels for nesting sites. Nests are built and eggs laid in February and March. The incubation period is 5-6 weeks, followed by a nestling period of 8 to 10 weeks.

Golden eagle nesting sites have been recorded in the following mountain ranges of the study area based on data from BLM (1984) - Fairview and Sidewinder Mountains, Granite Mountains, Ord Mountains, Stoddard Ridge, Newberry Mountains, Calico Mountains, Clark Mountain Range, and McCullough Mountains. In addition, suitable nesting habitat appears to be available in many of the other isolated mountain ranges of the study area. It is likely that at least several golden eagle pairs nest in the mountains of the study area. These residents, as well as occasional migrants from the north, are likely to forage throughout all but the developed portions of the study area.

Ferruginous Hawk and Other Raptors

The ferruginous hawk (Buteo regalis) is a Category 2 candidate for Federal listing (USFWS 1982), a protected bird in Nevada (NDW 1984), and is on the Audubon Society's 1982 Blue List (Tate and Tate 1982). The latter list includes bird species whose populations throughout the nation appear to be declining or are of sufficiently small numbers to warrant concern and increased observations.

Ferruginous hawks breed in central Canada, south to eastern Oregon, Nevada, Arizona, and New Mexico. During the winter, they extend their range into California and Mexico. These raptors occur in open, semi-arid plains and nest in a variety of situations including low hillsides, small cliffs, trees along creeks, and on transmission line towers. Ferruginous hawks begin nesting in March and April, and move to southern wintering areas in September. They are fairly common winter visitors in certain inland valleys of southern California, including Owens Valley (Inyo County), Carrizo Plains (San Luis Obispo County), and the Lucerne Valley (San Bernardino County) (Garrett and Dunn 1982). The latter is adjacent to the western portion of the study area.

Ferruginous hawks are likely to be fairly common winter visitors (September to April) in the study area, foraging over open and lowland creosote bush scrub, grassy areas, and croplands (in the western portion of the study area).

There are no other raptors that reside or make seasonal use of the study area that are candidates for Federal listing (USFWS 1982) or BLM sensitive species. However, there are several raptors on the list of California bird species of special concern developed by Remsen (1978) for the CDFG that are known to occur in the study area. Although these species do not have the status afforded to candidate and/or BLM sensitive species, they are of concern to conservationists and resource agency personnel due to their rarity, vulnerability, and/or unusual scientific interest. These species include the merlin, northern harrier, Cooper's hawk, sharp-shinned hawk, and prairie falcon. Only the latter is a resident breeder in the Mojave Desert of California and Nevada. The other species are rare to uncommon winter visitors and migrants to mostly selected, localized portions of the Mojave Desert; hence, their occurrence in the study area is very seasonal, restricted in location, and rare to uncommon.

The prairie falcon (Falco mexicanus) is a resident breeder of the inland arid western United States. During the late winter and early spring breeding period, prairie falcons establish nests on cliffs and escarpments in mountainous terrain. Nesting sites are usually associated with nearby broad, open habitat for hunting. During the winter, they extend their range to agricultural areas and to the coast. Nesting habitat requirements closely resembles that described for the golden eagle. In the study area, prairie falcon nesting sites have been recorded in most major mountain ranges in the western portion of the study area based on data from the BLM (1984) including, Bell Mountain, Fairview Mountain, Stoddard Ridge, Newberry Mountains, Calico Mountains, and Cady Mountains. In addition, suitable nesting habitat appears to exist in most other mountain ranges of the study area. Prairie falcons are expected to be occasional to fairly common residents of the study area, nesting in similar habitats as the golden eagle and foraging throughout most of the study area.

Desert Bighorn Sheep

There are three races of bighorn sheep in California: Nelson's (desert) bighorn sheep (Ovis canadensis nelsoni) which is found in the desert mountain ranges; Peninsular bighorn sheep (O. c. cremnobates) which is distributed along the western margin of the Colorado Desert from Palm Springs south into Mexico; and the California bighorn (O. c. californiana) which is found in the Sierra Nevada. The latter two are listed as rare by the California state government (CDFG 1980), but have no Federal status. The desert bighorn sheep is a BLM sensitive species (BLM 1980), a fully protected mammal according to the CDFG Code (Section 4700), and a regulated game animal in Nevada (NDW 1984).

In California and Nevada, desert bighorn sheep occur in more than 50 and 24 mountain ranges, respectively (Weaver 1982; McQuivey 1978). Most of the interior mountain ranges of the Mojave Desert support small populations of less than 20 sheep. The distribution and abundance of sheep have been severely reduced since the nineteenth century due to effects of human activities; as a result, many desert mountain ranges no longer support sheep populations.

Desert bighorn sheep occur in relatively inaccessible mountainous areas characterized by rocky, rugged and steep terrain (Monson and Sumner 1980). Rolling hills and gentle slopes of alluvial fans are only used briefly and infrequently. Habitat requirements include adequate food, water, escape routes, and shelter. Bighorn sheep feed on a wide variety of plants. Research in Nevada indicates that their diet consists primarily of forage grasses (65 to 75 percent), shrubs (15 to 30 percent), and herbs (5 to 10 percent) (McQuivey 1978). Important plants include needlegrass (Stipa), galleta grass, Mormon tea, winter fat (Erotia lanata), saltbush, filaree (Erodium cicutarium), and buckwheat (Eriogonum spp.).

Bighorn sheep are social animals and form different groups during the year. Most herds are groups of ewes with lambs and juveniles. During the fall

rutting period (July to November), rams join the maternal groups; at other times, rams occur singly or in small bachelor groups. Most observations of sheep consist of single individuals or groups of two to three; groups of more than 10 to 12 are unusual but occur occasionally in larger mountain ranges. Lambing occurs in February to March in special habitats which are scattered through the range and characterized by precipitous terrain, numerous escape routes, and shelter (caves). Lambing habitat is usually not readily identifiable due to its scattered occurrence, inaccessibility, and lack of dependence on water sources or food (McQuivey 1978).

Water is the most important factor limiting the distribution and abundance of bighorn sheep. Sheep acquire a small amount of water in plant food; this is an important source in the winter and may provide sufficient water such that sheep do not require free water for one to two weeks, and possibly longer periods. During the summer period, water demand is greater and sheep usually visit a permanent water source daily, but may do without water for three to five days.

Bighorn sheep occupy permanent home ranges that vary in size depending on the availability of water and food. In the winter, sheep are widely scattered and are known to travel 10 to 20 miles away from water sources. During the summer, sheep occupy a restricted area near a permanent water source, rarely traveling more than one to two miles away. Sheep feed at lower elevations in the spring, moving upslope following the progression of plant growth as summer develops. Bighorn sheep are not migratory, but infrequent dispersals may occur between mountain ranges due to long-term population expansions or seasonal disposals in search of food.

The desert bighorn sheep occur in several mountain ranges of the study area. These occurrences include permanent habitat (maximum year-round area) of known populations, restricted (summer) habitat surrounding water sources, and transient habitats or movement corridors (winter habitats).

In the Nevada portion of the study area, bighorn sheep are known to occur in the following mountain ranges: South Springs, McCullough, Roach, Sheep, Highland, and Eldorado (McQuivey 1976, 1978). Permanent populations are present only in the South Springs, South McCullough, Highland, and Eldorado mountains. Within the study area, the McCullough Mountains provide the largest continuous habitat and permanent population - approximately 151 square miles and 158 sheep, respectively, in 1976 (McQuivey 1978). The South Springs and Highland ranges support significantly fewer sheep. Population surveys conducted in 1984 indicate that approximately 81 sheep reside in the McCullough Mountains. The reason for the decline in numbers is unclear (Delaney 1984).

There are 18 water sources in the McCullough Mountains south of the pass, of which only two are year-round (McQuivey 1976). There is only one seasonal water source in the northern McCullough Mountains. As a result, during the summer critical period, sheep are restricted to the southern areas near the water sources. There is past evidence of limited seasonal dispersal and

possible migration between the south McCullough Mountains and Roach and Sheep mountains; however, the latter areas do not have sufficient water and habitat features to support permanent, and possibly regular seasonal, populations. Compared to other mountain ranges in southern Nevada, the McCullough Mountains provide relatively poor sheep habitat due to limited water sources; hence, the population densities for this range are comparatively low (McQuivey 1978).

A permanent small population occurs in the Clark Mountain range near the state border. Sheep in this range apparently move freely between the Clark and South Spring mountains. These movements appear to be normal dispersal and not migrations (Weaver and Hall 1971). The population in the Clark Mountains was estimated at 40 individuals (Weaver and Hall 1971; Weaver 1982), and the population appears to be stable. A 1984 survey of the Clark Mountains showed there were 114 sheep, indicating the population has increased recently (Bickel 1984). Several permanent springs occur on Clark Mountain itself, although most are used by cattle and humans. Pachalka Spring appears to be the only one used regularly by sheep. Although there is high quality habitat between Keany Pass and Stateline, there is no permanent water source in this area.

The nearby Mescal Range and Ivanpah Mountains do not support permanent populations of sheep (Weaver et al. 1969). These mountains are only expected to be used infrequently for travel or transient feeding.

Shadow Mountain apparently supported sheep populations in the past, but none have been recorded in the past several decades (Weaver 1982). Similarly, the Ord and Rodman mountains and Alvord Mountain in the western portion of the study area represent historic habitat for desert bighorn sheep, but none have been sighted since the 1950s and 1960s. In 1984, a survey of the Newberry Mountains (BLM 1984) revealed bighorn sheep at several areas, all near Kane Spring. The discovery of sheep in these mountains suggests a possible reinvasion by sheep of previously abandoned ranges.

Permanent populations of desert bighorn sheep are present on Avawatz Mountain and the Cady Mountains. In the former location, where the population is considered stable, sheep are known to make infrequent movements to portions of the Soda Mountains for brief periods of time (Weaver and Mensch 1970). A permanent population of 25 sheep occupy the Cady Mountains, including rugged bluffs above Afton Canyon (Weaver et al. 1969). Sheep populations in the Cady Mountains appear to be declining (Weaver 1982).

Areas of Special Biological Interest

Areas of special biological interest represent localized areas that contain biological resources that: (1) are of limited distribution; (2) are of unusual scientific or educational value; (3) represent prime examples of native biotic habitats; and/or (4) contain unusual assemblages of species. They are usually

small in area, localized, and broadly defined by topographic, hydrologic and geologic features such as basins, river channels, and mountain slopes.

Many of the following areas of special biological interest on public land in California are actively managed and/or protected by California BLM as described in the CDCA Plan (BLM 1980b). Management tools for these areas include designation of ACECs, Habitat Management Plans (HMP), and Unusual Plant Assemblages (UPA), in decreasing order of priority and protection status. ACECs are developed to protect significant areas or resources from adverse impacts of other uses. HMPs are developed to actively manage and maintain wildlife resources of areas of special interest, with limited multiple use allowed. UPAs are protected and/or managed according to the value and vulnerability of the botanical resources involved.

There are 16 areas of special biological interest in the study area. These areas are briefly described below.

South McCullough Mountains

This relatively narrow and elongated mountain range in the eastern portion of the study area is of special biological interest for several reasons. One, it is of sufficient elevation to support pinyon-juniper woodland on its higher slopes and ridges; this vegetation type is absent, or only occurs sparingly, on adjacent mountain ranges in the southern Nevada portion of the study area. Two, this range provides habitat for an important desert bighorn sheep population due to the presence of numerous seasonal and several permanent water sources. Third, these mountains also provide habitat for nesting raptors (golden eagles, prairie falcons, red-tailed hawks, ravens) and Gila monsters (in at least one or two desert wash areas).

Ivanpah Valley

Ivanpah Valley is located at the California/Nevada border. It supports relatively high densities of desert tortoise in the creosote bush scrub surrounding the dry lake. Tortoise densities of greater than 200 individuals per square mile have been recorded in the southern portion of the valley (Lukenbach 1982). Ivanpah Valley has been designated a crucial habitat for desert tortoise in the CDCA Plan (BLM 1980b). No HMP has been developed by California BLM for this area to date.

Clark Mountain Range

These mountains near the California-Nevada border represent an unusual biological feature in the study area for both botanical and wildlife resources. Clark Mountain reaches an elevation of 7,929 feet and supports a small relictual stand of Rocky Mountain white fir (Abies concolor var. concolor) near the crest. At lower elevations, pinyon-juniper woodland grades into Joshua

tree woodland and blackbush scrub, providing a striking example of a desert and montane vegetation gradient. Several regionally rare plant species occur at higher elevations in the range; in addition, many Great Basin plant species reach their western distribution limits on Clark Mountain at mid-elevations. There is also an unusually rich assemblage of cacti species.

A permanent population of bighorn sheep occupies the range, as well as several pairs of nesting golden eagles. Clark Mountain represents one of the two locations in California where Gila monsters have been recorded. Clark Mountain has been designated an ACEC by California BLM (BLM 1980b).

Shadow Valley

Shadow Valley is located west of the Clark Mountain Range and contains scattered areas of undisturbed Joshua tree woodland. It also supports relatively high densities of desert tortoise (BLM 1980b). Lukenbach (1982) estimates tortoise density of 40 to 155 individuals per square mile in the south central portion of the valley. No HMP has been developed by California BLM to date for Shadow Valley, nor has this area been proposed for ACEC designation (BLM 1980b).

Soda Springs

The Soda Springs area in the central portion of the study area is of special biological interest because of important botanical and wildlife resources. Along the western edge of Soda Dry Lake, there are numerous seeps and springs which have produced ponds, brackish meadows, and freshwater marshes. Hence, there are many aquatic and semi-aquatic plants present including salt grass (Distichlis spicata), rush (Juncus copperi), alkali weed (Nitrophila occidentalis), ditch grass (Ruppia cirrhosa), and bulrush (Scirpus spp.). The Federally endangered Mohave tui chub also occurs in three small ponds at Soda Springs; these ponds represent the Critical Habitat for this species and are carefully managed by the BLM and USFWS. Soda Springs has been designated an ACEC by California BLM (BLM 1980b).

East Cronese Lake and Mojave River Wash

East Cronese Lake is a dry lake bed in the central portion of the study area that is occasionally inundated during wet winters or due to flood diversions from the Mojave River Basin. Under such conditions, it provides seasonal habitat and/or a stopover site for migrating or overwintering waterfowl. Desert wash vegetation is well-developed south of East Cronese Lake and in the Mojave River Wash. In these areas, there are scattered hummocks dominated by desert willow (Chilopsis linearis) and honey mesquite (Prosopis glandulosa). These wash woodland/scrub areas represent some of the largest in the central Mojave Desert. No HMP has been developed to date for this area

by California BLM, nor has this area been proposed for an ACEC designation (BLM 1980b).

Crucifixion Thorn Stand

The crucifixion thorn (Castela emoryi) is a plant species more common in the Colorado Desert of California and Mexico. Its northernmost population occurs as an isolated stand northwest of Afton Canyon. Due to the highly disjunct nature of this stand, it is of special botanical interest. This area has been designated a UPA by California BLM (BLM 1980b).

Afton Canyon

The Mojave River surfaces in Afton Canyon during most of the year, supporting riparian woodlands dominated by willows (Salix spp.), cottonwood (Populus fremontii), mesquite (Prosopis glandulosa), tamarisk (Tamarix spp.), and scattered marsh plants such as cattails (Typha spp.). These riparian woodlands and ponds provide habitat for the regionally rare western pond turtle, a diverse and abundant avifauna, and several less common amphibians and reptiles of the Mojave Desert. In addition, bighorn sheep are known to water along portions of the river in Afton Canyon. Afton Canyon has been designated an ACEC by California BLM (BLM 1980b).

Cady Mountains

These mountains, located in the central portion of the study area, provide habitat for a permanent population of bighorn sheep. In addition, golden eagle and prairie falcons are known to nest in these mountains (BLM 1980b). This area was originally proposed for ACEC status; no HMP has been developed to date by California BLM (BLM 1980b).

Newberry Mountains

These mountains in the south-central portion of the study area, along with the adjacent Granite, Rodman and Ord mountains, provide nesting habitat for many raptors, particularly golden eagles and prairie falcons (BLM 1980b). Due to the abundant, continuous nesting habitat available, these mountains are of special biological interest. They were originally proposed for ACEC status; no HMP has been developed to date by California BLM (BLM 1980b).

Harper Dry Lake

Harper Dry Lake, northeast of Kramer Junction, is seasonally inundated lake surrounded by irrigated agricultural fields. Runoff from these fields are the major surface water source. Marsh vegetation dominated by cattails (Typha

spp.) and sedges (Carex spp.) occur along the western edge of the lake. Harper Dry Lake supports a very diverse resident and seasonal avifauna, particularly of wetland bird species and migratory waterfowl. California BLM has established an ACEC at Harper Dry Lake and a HMP has recently been completed.

Eriophyllum Mohavensis Preserve

This 32-acre parcel located northeast of Kramer Junction contains one of the largest known populations of Barstow woolly sunflower (Eriophyllum mohavensis), a BLM sensitive species and candidate (Category 2) for Federal listing. Other smaller populations of this species occur in the western portion of the study area. A HMP has been developed for this area which is also designated an ACEC.

Sclerocactus Preserve

Two half-sections of land on the bluffs overlooking Helendale have been designated an ACEC by California BLM. This area contains a large population of Mojave fishhook cactus (Sclerocactus polyancistrus), a BLM sensitive plant species and a candidate (Category 2) for Federal listing. Other smaller populations of this species are known from nearby areas. No HMP has been developed for this area to date by California BLM.

Western Mojave Saltbush Community

This area encompasses a significant portion of the western Mojave Desert where at least four species of Atriplex dominate (in various combinations) the low-lying alluvial plains and basins. This vegetation type has been gradually displaced by agricultural activities and grazing. Because of the widespread and often disjunct distribution of this vegetation type and due to the large portion of its occurrence on non-public land, it has not been designated an ACEC. However, it is designated a UPA by California BLM (BLM 1980b). In the study area, it occurs in scattered low-lying areas near Kramer Junction.

Fremont/Stoddard Valleys

This area of special biological interest encompasses the broad alluvial plains from Fremont Valley (outside the study area), southeast to Stoddard Valley. This includes most of the western portion of the study area. This area supports very high densities of desert tortoise, including localized areas with greater than 200 individuals per square mile. It has been designated a crucial habitat for desert tortoise by California BLM (BLM 1980b). It was originally proposed for ACEC status; no HMP has been developed for this area.

Mojave River

The Mojave River represents an important wetland habitat in the study area, as well as the Mojave Desert. Dense riparian woodland dominated by willows (Salix spp.), cottonwoods (Populus fremontii), and other wetland shrubs and trees occur from south of Victorville to near Helendale. Cottonwoods become less frequent along the lower reaches of the river, as it broadens into a wide sandy desert wash with dune areas and scattered desert wash woodlands dominated by mesquite (Prosopis glandulosa) and cat claw (Acacia greggii). The riparian woodlands and areas of surface water (Victorville to Helendale and at Afton Canyon) provide habitat for a diverse avifauna and various amphibians and fish, respectively.

Wetlands

Pursuant to Executive Order 11990 (Protection of Wetlands, May 24, 1977), wetland resources in the study area have been identified and include the following habitats:

- Desert wash (scrub and woodland) - along various seasonal streams and broad drainages throughout the study area.
- Riparian woodland - along the perennially flowing portions of the Mojave River.
- Lakes - perennial and seasonally inundated (dry lakes) bodies of water throughout the study area.

These habitats, to various degrees, support wetland vegetation and a comparatively abundant and varied fauna (particularly with regard to migrant birds and/or waterfowl). These wetland habitats are described in the previous sections of this report.

BIOLOGICAL RESOURCES WITHIN AND ADJACENT TO THE ALTERNATIVE CORRIDORS

The biological resources within one mile of the centerline of each alternative corridor (link) are described in the following sections. In addition, important biological resources adjacent to these corridors are also described. This information was developed based on an extensive literature review, contacts with knowledgeable persons, and field reconnaissance surveys.

The locations of important biological features, field observations, and major transitions between vegetation/habitat types were carefully noted, described and mapped for each corridor. In order to summarize the extensive results of these detailed, site-specific analyses for over 550 miles of alternative corridors, comprehensive inventory figures and tables were developed. Important biological resources within and adjacent to the two-mile-wide study

corridors are shown in Figures 4-2 and 4-3 (Map Volume). A more detailed description of biological resources is presented in Table B-1 (Appendix B) by milepost for each link. Information from Figures 4-2 and 4-3 and Table B-1 are summarized briefly in the following sections and associated summary tables.

Vegetation Types

Vegetation types traversed by the proposed corridors are shown on Figure 4-2 and described in detail on Table B-1. A summary and tabulation of miles of vegetation type traversed by each alternative corridor are presented in Table 4-6.

Creosote bush scrub is the dominant vegetation type traversed by the proposed corridors. This vegetation type often intergrades with saltbush scrub and Joshua tree woodland; in these transition areas, Larrea tridentata is the dominant plant species, but widely scattered Atriplex spp. or Yucca brevifolia are present due to local environmental or topographic conditions. Creosote bush scrub occurs along all alternative corridors, over a wide range of elevations (Table 4-6).

Saltbush scrub is the second most common vegetation type traversed by the alternative corridors. It primarily occurs in low-lying basins and along edges of dry lakes and the Mojave River. Saltbush scrub is absent from 19 of the alternative corridors, most of which are in mountainous areas or on alluvial fans and bajadas. The greatest continuous areas of saltbush scrub occur along Link 3 (Ivanpah Valley), 8 (Shadow Valley), 11 (Soda Springs area), 20a (Coyote Dry Lake), and 38-42 (near Kramer Junction) as shown on Figure 4-2.

Blackbush scrub traversed by the proposed corridors is restricted to the higher slopes of the Clark Mountain Range (Links 5c-8) and the western portions of Shadow Valley (Links 7-10) as shown on Figure 4-2 and summarized in Table 4-6.

Joshua tree woodland is also a highly restricted vegetation type that is traversed by several alternative corridors. Widely scattered Joshua trees occur in creosote bush scrub and blackbush scrub vegetation types throughout the mid- and upper-elevation corridors (Links 1-10) in the McCullough and Clark mountains and in Shadow Valley, although the density of trees is too low to be classified as a Joshua tree woodland. Well-developed Joshua tree woodland occurs in small, scattered patches in the western portion of the study area, including portions of Links 30, 31 and 36-39 near Kramer Junction and Adelanto.

Desert wash vegetation occurs along all alternative corridors; however, only wash scrub with dense vegetation and well-defined channels or wash woodland with Acacia greggii, Chilopsis linearis, and/or Prosopis glandulosa were identified and mapped as shown in Figure 4-2 and summarized in Table 4-6. Major desert washes are traversed by Links 1 (Eldorado Valley), 11 (Mojave

River Wash), 14 (East Cronese Lake and Mojave River Wash), 16 (East Cronese Lake), 23 (Mojave River), 24 (Mojave River), and 27 (Mojave River).

Riparian woodland is only traversed by Links 27, 32 and 33 at the upper Mojave River where Salix spp. and Populus fremontii are present in well-defined willow, mesquite, smoketree, and the introduced tamarisk. Major desert wash areas with well-developed woodland vegetation occur in the Eldorado Valley (Link 1), Mojave River Wash (Link 11), East Cronese Lake (Links 14 and 16), and along the lower Mojave River (Links 23, 24 and 27).

Riparian woodland consists of a dense thicket of shrubs and trees along a river or stream channel with significant surface flow. Dominant plants include willow and cottonwood trees. This vegetation type is present in the study area only along the upper reaches of the Mojave River (south of Victorville to near Helendale) and at Afton Canyon. The upper Mojave River is traversed by three corridors (Links 27, 32 and 33).

No Mojave montane forest or pinyon-juniper woodland are traversed by the alternative corridors, although both are present at upper elevations on Clark Mountain. Pinyon-juniper woodland also occurs in the McCullough Mountains.

Wildlife

Four wildlife habitats can be identified in the study area, based primarily on the structure and species composition of the vegetation type which provides food and shelter: (1) scrub habitat (including desert wash habitat), (2) woodland habitat, (3) riparian woodland habitat, and (4) montane forest habitat. Each habitat supports a fairly well-defined assemblage of amphibians, reptiles, birds and mammals.

Wildlife species associated with scrub habitats are expected to be most prevalent along the alternative corridors because creosote bush scrub and saltbush scrub are the most common vegetation types traversed by the corridors. The abundance and diversity of this species assemblage are expected to vary according to local conditions, including availability of food, water sources, and disturbances. With the exception of built-up areas in the Mojave River Valley area from Barstow to Victorville and in the immediate areas surrounding Baker, Soda Springs, Yermo, and Kramer Junction, a full complement of scrub habitat species is expected to occur during all or part of the year in the study area.

Desert wash habitats occur in scattered locations along many alternative corridors as shown on Table 4-6. However, the abundance and diversity of wildlife species are not expected to increase in these localized habitats unless there is a concentration of wash trees such as mesquite and catclaw.

Woodland habitat occurs primarily in the Joshua tree woodland areas in the Clark and McCullough mountains (Links 1-10), in Shadow Valley (Links 7-10), and near Kramer Junction and Adelanto (Links 30, 31 and 36-39). Wildlife

species expected to occur in this habitat type were described previously in this report.

Riparian woodland habitat is traversed by Links 27, 32 and 33 along the upper Mojave River where numerous resident, migrant, and seasonal visitor bird species are likely to occur (Figure 4-2).

Montane forest habitat is not traversed by the alternative corridors.

Due to the season of the year when field surveys were conducted (November and December), only a limited number of indirect and direct observations were made of wildlife along the alternative corridors. In general, most amphibians, reptiles and mammals reduce their activities during the early winter when temperatures are dropping and food is relatively scarce. In addition, few winter bird visitors have arrived in the Mojave Desert and most fall migrants have already passed through. Hence, the abundance and diversity of the avifauna along the corridors were relatively low during the field surveys. However, despite these limitations on observations of wildlife, direct sighting and indirect evidence of common wildlife along the alternative corridors were noted and are listed in Table B-1.

Legally Protected Species

Legally protected species include plants or animals designated rare, threatened or endangered by the California or Nevada state governments or the Federal government, as described previously.

Plants

No legally protected plant species are known or likely to occur within or adjacent to the alternative corridors.

Fish and Wildlife

Several legally protected fish and wildlife species are known or could potentially occur in the study area. Their occurrence within or adjacent to the alternative corridors is briefly described below.

Mohave Tui Club

The alternative corridors do not directly traverse the habitat of this legally protected fish which is restricted (in the study area) to ponds at Hinkley and Soda Springs. Links 11 and 42 pass within several miles of these ponds.

Gila Monster

The legally protected Gila monster is an uncommon resident of sandy washes in southern Nevada and an extremely rare resident (or possibly foreign introduction) from the eastern Clark Mountains. Corridors in southern Nevada pass by or within wash habitats likely to support Gila monsters. Link 0 passes within one mile of a potential Gila monster habitat; Link 1 passes through a potential Gila monster habitat in the McCullough Mountains, as well as within several miles of another potential habitat; and Link 5c traverses potential Gila monster habitat on the eastern slopes of the Clark Mountain Ranges (Figure 4-3). The actual occurrence of Gila monsters in the habitats within the alternative corridors is expected to be rare to uncommon.

Desert Tortoise

The desert tortoise occurs throughout most lowland scrub habitats in the study area. Relatively high populations are known to occur in the Ivanpah Valley, Shadow Valley, and in the western portion of the study area.

Known and potential desert tortoise habitat traversed by the alternative corridors is shown on Figure 4-3, described in detail in Table B-1, and summarized in Table 4-7. Estimated density values have been classified into five levels as described in Table 4-7: (1) not present, (2) low to rare or possibly absent, (3) low to moderate, (4) moderate to high, and (5) very high. The length of tortoise habitat (according to density class) traversed by each alternative corridor is presented in the table.

For at least a portion of their length, most corridors traverse potential desert tortoise habitat. Areas unlikely to provide suitable tortoise habitat such as rocky areas and barren dry lakes were readily identifiable along most corridors. The remaining portions of the corridors provide at least the lowest density class habitat for tortoise (0 to 50 individuals per square mile) as shown in Table 4-7. Localized areas of low to moderate densities (50 to 150 individuals per square mile) occurred along numerous corridors throughout the study area. Areas of moderate to high densities (100 to 250 individuals per square mile) are traversed by Link 10 in the Shadow Valley, Link 20b in the central portion of the study area, and by Links 24a-25, 27-29, 30, 32, 33, 36, 37 and 39-43 in the western portion of the study area (Figure 4-3). Very high density areas (greater than 250 individuals per square mile) are only traversed by Links 24a (1.8 miles) and 27 (6.5 miles).

Swainson's Hawk

This legally protected bird species is a summer breeder in the Mojave Desert, but nesting records have not been recorded recently in the study area. Suitable nesting habitat is potentially available along the Mojave River from Victorville to Helendale, and in areas of Joshua tree woodland near Adelanto, Shadow Valley, Clark Mountain, and Beer Bottle Pass. However, the future

occurrence of nesting Swainson's Hawks within or near any of the alternative corridors is expected to be rare; in contrast, migrating Swainson's Hawks are likely to temporarily utilize most open habitats traversed by the alternative corridors throughout the study area.

California Yellow-billed Cuckoo

Suitable nesting for this legally protected species in the study area is restricted to the dense riparian woodlands of the upper Mojave River. Although the occurrence of nesting individuals in this area is expected to be very rare, Links 32 and 33 may traverse potential habitat used by migrant yellow-billed cuckoos.

Least Bell's Vireo

This legally protected species which is restricted to riparian habitats has not been sighted in the study area since 1970. Suitable habitat appears to occur along portions of the Mojave River, including the areas crossed by Links 27, 32 and 33. However, the occurrence of the least bell's vireo in the study area would be expected to be very rare and of migrant individuals only.

Mohave Ground Squirrel

All or part of the alternative corridors in the western portion of the study area traverse the known range of the legally protected Mohave ground squirrel, as shown on Figure 4-3 and summarized on Table 4-8. No data are available on the relative density of ground squirrels in the traversed habitats.

Species of Special Interest

Species of special interest in the study area are defined previously in this report, where a brief description of their general distribution in the study area is provided.

Plants

There are 10 plants of special interest that are known to occur in the study area, as discussed previously and summarized in Table 4-5. Of these species, two are very unlikely to occur near or within the alternative corridors - Coryphantha vivipara var. alversonii and Puccinellia parishii. The other eight species are known or likely to occur within portions of the corridors based on the proximity of known populations to the corridors and/or the presence of suitable habitat within or near the corridors.

Potential habitat and locations of known populations of special interest species within and adjacent to the alternative corridors are shown on Figure 4-2. In addition, miles of potential habitat traversed by alternative corridors for each species are summarized and tabulated in Table 4-9.

Astragalus jaegerianus (Jaeger's milkvetch)

Potential habitat for this species is located along 13.5 miles of Links 20a and 20b between Lane Mountain and the Calico Mountains. The nearest known populations are two miles northwest of Link 20a as shown on Figure 4-2.

Chorizanthe spinosa (Mojave spineflower)

This species appears to be widely scattered in the northwest portion of the study area. Potential habitat for Mojave spineflower is traversed by most corridors in the western study area, as shown on Figure 4-2 and summarized in Table 4-9.

Coryphantha vivipara var. alversonii (Foxtail cactus)

This species is known from only one site in the study area in the Clark Mountains. As shown on Figure 4-2, this site is one mile south of Link 8. This species is not expected to occur within or directly adjacent to the alternative corridors. Its occurrence in the study area is expected to be very rare.

Cymopterus deserticola (Desert cymopterus)

This species is known from populations near Kramer Junction and Apple Valley and is expected to be widely distributed in scattered locations in the western portion of the study area. Potential habitat occurs along numerous corridors in the western study area as shown on Figure 4-2 and summarized in Table 4-9.

Eriophyllum mohavensis (Barstow woolly sunflower)

This species is known from current populations near Kramer Junction and from a historic record near Barstow. Suitable habitat in these areas is traversed by portions of Links 21b-23, 37-39 and 41-42 as shown on Figure 4-2 and summarized in Table 4-9.

Penstemon bicolor ssp. bicolor and roseus (Bicolored perstemons)

These two subspecies are known from various populations in the mountains of southern Nevada, including the McCullough Mountains. Approximately 6.0

miles of Link 1 traverses potential habitat of these species as shown on Figure 4-2.

Phacelia anelsonii (Nelson's phacelia)

In the study area, this species is only known from a single population in the McCullough Mountains. However, potential habitat is traversed at higher elevations of the McCullough and Clark mountains along portions of Links 1 and 5c-8 (Figure 4-2).

Puccinellia parishii (Parish's alkali grass)

This very rare species is restricted to small, localized alkali seeps and has not been positively confirmed in the study area. As a result, its occurrence along the alternative corridors is very unlikely.

Sclerocactus polyancistrus (Mojave fishhook cactus)

Several populations of this cactus are known in the western portion of the study area. Potential habitat for this species is traversed by several corridors as shown on Figure 4-2 and in Table 4-9.

Sphaeralcea rusbyi ssp. eremicola (Rusby's desert mallow)

This species is only known from one population near Keany Pass, Clark Mountains. Potential habitat is traversed by alternative corridors in the Clark Mountain Range, including Links 5c-8 (Figure 4-2, Table 4-9).

Wildlife

Wildlife species of special interest that are known or likely to occur in the study area were described previously. Potential habitat of these species that occurs within or adjacent to the alternative corridors is shown on Figure 4-3, described in detail in Table B-1, and briefly described below.

Golden Eagle

The golden eagle is a resident of the study area. Nesting pairs are expected to occur in at least several of the major mountain ranges traversed by the alternative corridors. In addition, golden eagles are expected to travel and hunt over most of the study area throughout the year.

The alternative corridors traverse and/or pass by several mountain ranges that provide known or likely nesting habitat for the golden eagle as summarized in

Table 4-10. Link 44 is within one mile of known golden eagle nesting habitat in the Calico Mountains. Links 5c, 6, 7 and 8 are no closer than two miles to the nesting sites on Clark Mountain and Link 23 is four to five miles north of known nesting sites in the Newberry Mountains (Figure 4-3). All other corridors listed in Table 4-10 are either adjacent or several miles from potential nesting habitat; no corridors directly traverse potential nesting habitats.

Ferruginous Hawk and Other Raptors

The ferruginous hawk is likely to be a common winter visitor throughout the study area. This species does not nest in the Mojave Desert. It is likely to occur along most of the alternative corridors for brief periods of time during the winter for hunting and traveling events.

The prairie falcon is a relatively common resident breeder in the study area as described previously, utilizing similar nesting and foraging habitats as the golden eagle. Hence, the occurrence of potential prairie falcon nesting habitats along the alternative corridors follows that described previously for the golden eagle.

Desert Bighorn Sheep

Permanent populations of the desert bighorn sheep occur in several of the mountain ranges in the study area. In addition, there are several smaller mountain ranges that provide seasonal (winter) habitat for bighorn sheep as shown on Figure 4-3.

The alternative corridors pass through or near portions of several known, permanent habitat areas, including the McCullough Mountains (Link 1), Clark Mountains (Links 5c-8), and Cave/Cady Mountains (Links 15 and 17). Several corridors pass through or near transient bighorn sheep habitat in the Sheep Mountains (Link 1), Roach Mountains (Link 2), Spring Mountains (Link 6), and Soda Mountains (Link 11), as summarized in Table 4-11. These latter habitats are expected to be utilized very infrequently by bighorn sheep during brief periods of the winter.

Areas of Special Biological Interest

Areas of special biological interest in the study area are defined and described previously. The occurrence of alternative corridors within or adjacent to these areas is discussed below.

South McCullough Mountains

Link 1 traverses the mid-point of the McCullough Mountains, one to two miles north of McCullough Pass. This point of crossing is several miles north of the pinyon-juniper woodland and permanent desert bighorn sheep populations in the southern range.

Ivanpah Valley

Links 5b, 5c and 6 traverse the northern portion of Ivanpah Valley as shown on Figure 4-3. Links 5b and 5c also cross a small portion of the barren Ivanpah Dry Lake. Areas of Ivanpah Valley that support the highest population densities of desert tortoise occur several miles to the south.

Clark Mountain Range

Links 5c and 6 traverse the two major passes on the southern and northern slopes of Clark Mountain, respectively. Both links cross areas of Joshua tree woodland, blackbush scrub, and cacti-rich creosote bush scrub; however, no pinyon-juniper woodland or montane forest vegetation is within or directly adjacent to these corridors. Both links are also along the margins of the permanent range of desert bighorn sheep on Clark Mountain and two miles away from known golden eagle nesting sites. Hence, Links 5c and 6 avoid traversing many of the important biological habitats of Clark Mountains. Both corridors are within the Clark Mountain ACEC.

Links 7 and 8 pass along the southwestern slopes of Clark Mountain near Mohawk Hill. Neither corridor traverses important biological habitats of this area of special interest.

Shadow Valley

Links 6, 7 and 8 traverse Shadow Valley, including areas of blackbush scrub and Joshua tree woodland. Only Links 7 and 8 occur in the areas of moderate density desert tortoise habitat according to the data of Lukenbach (1982).

Soda Springs

Links 11 and 12 traverse the northern portion of Soda Lake; however, this area is barren and devoid of any wetland plant species or habitat. Link 11 passes within one mile of the Soda Springs ACEC and two miles of the ponds containing the Mojave tui chub.

East Cronese Lake and Mojave River Wash

Portions of Links 11, 14, 15, 16 and 17 traverse the desert wash habitat of lower East Cronese Lake and the Mojave River Wash. Only portions of Links 14 and 16 pass through well-developed, but widely distributed hummocks of mesquite (Prosopis glandulosa) and desert willow (Chilopsis linearis).

Crucifixion Thorn Stand

Link 6 at Milepost 75 passes within two miles of the crucifixion thorn stand.

Afton Canyon

Links 15, 17 and 18 are one to two miles north of Afton Canyon, passing through the western portions of Cave Mountain. These links do not traverse the ACEC boundaries or areas of riparian woodland; however, Links 15 and 17 are located in an area used periodically by the permanent bighorn sheep population to the south in the Cady Mountains.

Cady Mountain

No alternative corridor traverses these mountain ranges. However, to the extent the permanent bighorn sheep population in this mountain range travels across Afton Canyon to Cave Mountain, Links 15 and 17 would occur in bighorn sheep habitat.

Newberry Mountains

Link 23 passes within four to five miles of the Newberry Mountains, well away from potential and known raptor nesting sites.

Harper Dry Lake

Link 41 occurs within one mile of Harper Dry Lake. Between Mileposts 0 and 0.9, this corridor traverses a dry low-lying area that is a topographic extension of Harper Dry Lake. This barren area does not support wetland plants nor does it appear to provide seasonal habitat for migrating waterfowl.

Eriophyllum Mohavensis Preserve

Link 41 passes within one and one-half miles of this ACEC. Although this corridor is not near the protected population of Barstow woolly sunflower in the ACEC, it (as well as other nearby links) does traverse potential habitat for this species.

Sclerocactus Preserve

Link 27 passes within one and one-half miles of this ACEC. Although this corridor is not near the population of Mojave fishhook cactus in the ACEC, it (as well as other nearby links) does traverse potential habitat for this species.

Western Mojave Saltbush Community

Several corridors in the western portion of the study area near Kramer Junction traverse scattered, localized areas dominated by the saltbush scrub designated a UPA by BLM. These corridors include Links 36-39, 40a and 41-43.

Fremont/Stoddard Valley

This area of special biological interest represents a broad regional area with relatively high densities of desert tortoise. It occurs in the western portion of the study area and is traversed by most of the corridors west of Barstow. Specific corridors which traverse areas of very high tortoise density include Links 24 and 27.

Mojave River

This area of special biological interest is traversed at locations by Links 23, 24, 27, 32 and 33. Riparian woodland is present at the crossings of Links 27, 32 and 33, while desert wash scrub/woodland is present at the crossings of Links 23 and 24.

AGENCY CONTACTS

Contacts were made with various agency personnel to gather data on biological resources of the study area and to discuss specific concerns with potential impacts of the proposed project. A summary of these contacts is presented at the end of this chapter.

IMPACT ASSESSMENT AND MITIGATION PLANNING


Introduction

The potential impacts of the proposed project on biological resources of the study area are evaluated in the following sections. Impacts to vegetation, wildlife, legally protected species, and special-interest species are addressed along all alternative transmission line corridors (see Figure 5-1 in the Map Volume). The magnitude and significance of adverse impacts due to the construction and operation of the proposed project are assessed below based on the sensitivity of the affected resources, the description of the proposed project as provided by DWP, and an impact assessment model developed by Dames & Moore for this study. This qualitative model was developed in order to produce logical, consistent and repeatable impact level ratings for potential adverse effects along each alternative corridor. These impact levels, which represent composite measures of impact magnitude and significance, are converted into quantitative data for use in comparing potential impacts along alternative links during the multi-disciplinary route selection process described elsewhere. Mitigation measures incorporated into this impact assessment process include "generic" measures proposed by DWP and included in the proposed project, and "selective" measures recommended by Dames & Moore to further reduce significant impact levels.

Impact Assessment Model

A qualitative impact assessment model was developed in order to evaluate the magnitude and significance of potential adverse impacts in a consistent, semi-quantitative manner such that impact levels along different alternative corridors can be compared to one another. This model applies only to construction impacts because they represent the only potentially significant impacts of the proposed project due to the temporary and permanent disturbance of biotic habitats associated with construction activities. In addition, only construction impacts provide a means to differentiate between impact levels of alternative links due to the varying requirements for new or upgraded access roads along each alternative corridor. As described in the following sections, impacts due to the operation and maintenance of the proposed transmission line (using any of the alternative corridors) are not expected to result in significant impacts on biological resources, nor do they differ greatly between alternative links.

The impact assessment process is comprised of seven elements as shown in the following diagram:



These elements, as well as the process by which impact levels are assessed in the model, are described in the following sections.

Impact Types

There are four biological impact types identified as described below:

Impact Type	Description
1	Adverse effects on vegetation.
2	Adverse effects on resident and seasonal wildlife and their habitat..
3	Adverse effects on legally protected species (animals only in this study).
4	Adverse effects on species of special interest (plants and animals).

Each of these impact types can be further categorized into construction- and operations-associated impacts as described in the following section.

Construction Impacts

Construction impacts for the proposed project include direct and indirect adverse effects. Both categories of impacts result from the requirement for access of workers and equipment along the proposed right-of-way to construct the transmission towers and lines. The level of these impacts varies between alternative links because the need to improve or establish new access (i.e., build roads) varies from link to link. This condition, along with the conclusion that only construction activities would result in potentially significant impacts, indicate that an evaluation of construction impacts will provide the best means to evaluate impact levels along any single alternative corridor (individual significance), as well as to compare impact levels between alternative links (relative significance).

Direct Effects

Direct impacts due to construction activities consist of temporary and/or permanent disturbance, displacement, and/or removal of biological resources which include native vegetation, wildlife and their habitat, legally protected species (animals only in this study), and species of special interest (plants and animals).

Areas of Temporary Disturbance (Short-Term Impacts) - Areas of temporary disturbance include tower base construction zone (200 X 200 feet every 3.8 to 4.4 miles); wire-pulling sites (200 X 200 feet every 2 miles); wire-splicing sites (20 X 50 feet every 2 miles); construction yards (400 X 540 feet every 20 to 30 miles); and batch plant yards (one to two acres every 20 to 30 miles). These areas will be cleared of vegetation when grading and recontouring are necessary to accomplish construction activities. In other areas, vegetation will not be removed. Although vegetation in the latter areas will be crushed, revegetation will potentially occur faster than in cleared areas. All equipment will be removed following construction and, if necessary, disturbed land will be recontoured. These disturbed areas will be left to revegetate naturally and to be recolonized by wildlife as vegetation becomes established.

Temporary disturbances represent short-term impacts. The restoration of biotic habitats in the desert typically occurs at a slow rate, often requiring decades to restore vegetative biomass to pre-disturbance levels. Although temporary disturbances are considered short-term impacts, the duration of the impact on biological resources may vary from several years to several decades. However, unlike long-term impacts, these areas of temporary disturbance will eventually be restored to functioning biotic habitats.

Areas of Permanent Disturbance (Long-Term Impacts) - Permanent disturbances represent long-term impacts that would persist for the life of the project. Areas of permanent disturbance and removal include tower bases (35 X 35 feet), access roads, and spur roads. The roads will be unpaved and approximately 14 feet wide, located primarily within the 200-foot-wide right-of-way. They will be used to transport vehicles and equipment during construction and routine maintenance. The use of these roads by the public after completion of construction activities is dependent on the wishes and authority of the landowners and DWP.

There are three types of access roads for the proposed project based on the occurrence of nearby existing access roads. One, new access roads would be built where no roads currently exist. Two, existing access roads of poor quality in the proposed right-of-way would be upgraded and used wherever possible. Three, existing sufficient access roads would be used wherever possible without need for improvement.

Acreages of potential permanent disturbance and removal for both access roads and spur roads combined are calculated as follows. New access roads would require the permanent removal of about 2.0 acres per mile of transmission line. Existing roads requiring upgrading would result in the permanent

loss of 1.5 acres per mile. Sufficient existing roads would require the removal of 0.5 acre per mile for the construction of spur roads. Access requirements for each alternative link are described in the project description. Most alternative corridors have existing sufficient or to-be-upgraded roads that would be utilized.

Indirect Effects

Indirect construction impacts consist of the potential new or increased unauthorized vehicular and human activities along access roads. Such activities could result in new or increased adverse effects on the local biota due to off-road vehicle (ORV) travel; collecting of cacti; incidental damage to native vegetation or special-interest plants; and harassment, disturbance or hunting of legally protected and special-interest wildlife. The magnitude and significance of these impacts would vary depending on the need for new or improved access roads along alternative corridors.

Another indirect construction impact would consist of the potential disturbance of wildlife in adjacent habitats due to the temporary introduction of, or increase in, vehicular travel, equipment noise, and human activity. Although this impact could result in the displacement of wildlife from adjacent habitats, the magnitude and significance of this impact are low because: (1) the disturbance is temporary and localized; (2) many wildlife species may potentially acclimatize to the offsite disturbance because it would not represent a direct threat; and (3) wildlife is expected to recolonize adjacent habitats following the completion of temporary construction activities. This indirect impact would not be potentially significant and would not vary between alternative links. Hence, this impact is not considered further in the assessment model, nor are any selective mitigation measures recommended except in bighorn sheep habitat (see Impact Intensity section).

Operations

The operating transmission line would consist of stationary, unmanned, and nonmechanical structures with negligible maintenance needs. Hence, operation and maintenance of the proposed line are not expected to result in any significant impacts to biological resources for any alternative route chosen. Potential impacts associated with increased human and vehicular activities along new or improved access roads during operations are addressed previously under Construction Impacts.

Transmission line towers and access roads are not expected to present an effective obstacle to wildlife movement. However, avian collisions with conductors resulting in injury and death are likely to occur infrequently and all along the chosen route by a variety of birds, particularly migratory waterfowl. This impact is considered to be of minor magnitude and significance because: (1) the line is not located adjacent to bodies of water where larger numbers of waterfowl would be in low-altitude flights; and (2) the rate of mortality due to

collisions with conductors is expected to be extremely small compared to the number of birds in the study area due to the small size of the obstacles (i.e., conductor wire) relative to flight space. Impacts to wildlife due to electrical and magnetic fields emitted from the operating conductors are expected to be negligible as discussed in a separate report. Electrocution of raptors on transmission towers would not occur due to the great distance between conductors relative to wing spans of golden eagles and other large raptors. In fact, the proposed transmission towers provide new perch and nesting sites for raptors in the study area--a net beneficial impact.

Operation impacts described above are not included in the impact assessment model because potential impacts associated with operation of the proposed project would not result in significant impacts nor would they differ between alternative corridors.

Resource Sensitivity

The sensitivity of the biological resources within and adjacent to the proposed alternative corridors are described below. Sensitivity is based primarily on the characteristics of the resource, including (in no particular order): (1) the legal and/or management status of the resource; (2) the apparent rarity (geographic distribution and general abundance) of the resource at both regional and state-wide levels; (3) the vulnerability and resilience of the resource to significant impacts associated with the construction of the proposed project; (4) the quantity of the resource potentially affected or its proximity to disturbance activities; and (5) the quality of the resource at that specific locale. The latter element is defined as the degree to which human disturbance such as agricultural activities or urban development has affected the resource. In this study, resource quality was not an important element in assigning resource sensitivities because most alternative links traversed habitats with only minor human disturbance.

Four levels of sensitivity are defined ranging from minimal (i.e., areas with nominal biota such as urban development) to maximum (i.e., areas of state-wide concern due to rarity and severe endangerment such as riparian woodlands). The sensitivities of the biological resources encountered along the alternative corridors are presented in Table 4-12. Note that the presence of several resources along a corridor may increase the biological sensitivity of that link beyond the individual sensitivities of the component resources.

Resource Quantity

The resource's quantity (i.e., the amount of the biological resource potentially affected) is defined here to incorporate the areal extent of temporary and permanent physical disturbances (direct impacts) and the potential increase in human access associated with new or improved access roads (indirect impacts). As such, it is a measure of the impact intensity. Areas of temporary disturbance can be described by a constant rate along each alternative

corridor as described in the Construction Impacts section. These areas include the disturbance zone around tower bases, wire-pulling sites, wire-splicing sites, construction yards, and batch plant yards. Impacts associated with access and spur road requirements (i.e., areas of permanent disturbance) vary in intensity or quantity of the resource affected along each alternative link. In general, existing access roads will be used when the proposed right-of-way closely parallels an existing utility corridor or where other existing roads provide sufficient access. Existing narrow or poor-quality access roads will be upgraded as needed rather than build new roads. New roads will be constructed where there is no existing access. The length of spur roads will increase with improved or new access roads.

Based on the above considerations, three levels of access requirements have been identified in which the quantity of the resource affected by direct and indirect construction impacts (i.e., impact intensity) would vary as described below:

<u>Access Requirement</u>	<u>Description</u>
Sufficient	Existing access roads are sufficient and no road building or improvement activities would occur. Intensity of impacts are due to temporary and permanent disturbances from spur roads, tower base and associated construction zone, wire-pulling and wire-splicing sites, construction yards, and batch plant yards. The major physical disturbance (i.e., road) and opportunity for unauthorized human and vehicular access already exist. Hence, this represents the lowest intensity of construction impacts and least amount of resource affected.
Upgraded	Existing roads would require upgrading. Other temporary and permanent disturbances would occur as described above. The intensity of impacts and quantity of resource affected are greater than above due to increased extent of permanent disturbances along the upgraded roads and the increased potential for unauthorized access on improved roads.
New	New access road would be constructed, resulting in new areas of permanent disturbances greater than at the lower two levels. All other areas of temporary and permanent disturbances would occur as described above. This level represents the maximum intensity of impacts and quantity of resource affected because new biotic habitats would be disturbed and new opportunities would arise for unauthorized human and vehicular activities along the right-of-way.

Access road requirements (i.e., impact intensity) for each alternative link are described by milepost in the Results section.

Generic Mitigation Measures

These measures represent elements of the proposed project designed to minimize and avoid, when feasible, significant impacts to biological resources along each alternative corridor. Generic mitigation measures consist of procedures, design features, and restrictions which will be implemented along all corridors. They are fully described in the separate project description. The generic mitigation measures that would apply to biological resources are listed below, including a brief description of their effectiveness in reducing impacts on biological resources.

1. All construction vehicle movement outside the right-of-way will normally be restricted to pre-designated access, contractor-acquired access, or public roads.
2. The areal extent of construction activities will normally be pre-determined with activity restricted to and confined within those limits.

These mitigation measures would require that the areas of temporary construction disturbance due to vehicle movements, tower construction, and other activities be minimized to the extent feasible by restricting the area of construction. These measures would apply to the entire lengths of all corridors and would minimize the total extent of temporary vegetation and wildlife habitat removal.

3. In construction areas where recontouring is not required, vegetation will be left in place wherever possible and the original contour will be maintained to avoid excess root damage and to allow for resprouting.

This mitigation measure would facilitate the restoration of biotic habitats by prohibiting clearing and grading in areas of temporary construction disturbance. Natural, unaided revegetation occurs more quickly in areas where vegetation is only crushed than in areas where it is removed. Hence, this measure would potentially increase the rates of revegetation and wildlife recolonization.

4. In construction areas where recontouring is required, revegetation and/or reseeding will occur as required by the landowner or BLM.

Although artificial revegetation efforts in disturbed desert vegetation have not generally proven to be successful (i.e., faster or more complete than natural, unaided revegetation), this measure could be an important procedure to restore special habitats disturbed by construction activities. It would only be applied where it is likely that revegetation procedures would be successful or necessary as required by the landowner or BLM.

Impact Level

Impact level is a composite index of the magnitude and significance of construction impacts (direct and indirect) associated with the proposed project. It represents the end product of the impact assessment model. Impact levels are assigned by the impact assessment model shown in Table 4-13. This matrix indicates the combinations of proposed access requirements and resource sensitivity which result in different impact levels after the application of generic mitigation measures. This formal method of assigning impact levels provides a consistent and repeatable impact assessment that can be used to evaluate the significance of individual impacts, as well as to compare impact levels between alternative corridors.

There are three basic impact levels used in this study for initial impact levels as shown on Table 4-13, as well as for residual impact levels described below. These levels are: L (low), M (moderate), and H (high). In addition, there are intermediate values within this three-level framework such that a range of initial and residual impacts could occur.

Initial impact levels for each alternative corridor are presented in the Impact Assessment/Mitigation Planning (IA/MP) charts at the end of this chapter. These impact levels are assigned for all portions of each link and are delineated by mileposts. Impact levels vary along each corridor as the access requirements and/or resource sensitivity change. In addition, impact types and access levels are designated along each alternative corridor in the IA/MP charts. A brief summary of the initial impact levels is provided below.

High Impact Level (H)

The maximum or very high impact level represents a significant impact to biological resources and occurs where there is a maximum sensitivity resource and a requirement for new access as shown on Table 4-13. The highest impacts occur at two locations along alternative corridors where very high density desert tortoise habitat (greater than 250 individuals per square mile) would be traversed.

Medium-high impact levels can occur at all levels of access requirements, depending on the sensitivity of the resource affected. These conditions occur where new access roads are required that traverse a desert wash, permanent range of bighorn sheep, or high density (100 to 250 individuals per square mile) desert tortoise habitat (IA/MP charts). High impact levels also occur along links where new roads would affect three or more moderate sensitivity resources such as habitats in the western portion of the study area that support special-interest plants, the Mohave ground squirrel, and moderate density desert tortoise populations (50 to 100 individuals per square mile). This impact level also occurs where an upgraded road would be installed through a riparian woodland.

Moderate Impact Level

Moderate impacts are not considered significant; they occur under a wide variety of combinations of access requirements and resource sensitivities as shown on Table 4-13 and in the IA/MP charts. These impact levels occur at least once along most alternative corridors.

Low Impact Level

Low impact levels are not significant. They are assigned only when there is an existing access road (sufficient) or an existing road that only requires minor improvements (upgraded) as shown on Table 4-13. Because most of the alternative corridors follow existing rights-of-way where there are existing roads, these impact levels are the most frequently assigned in the IA/MP charts.

Negligible Impact Level

This impact level only occurs along corridors that traverse minimal sensitivity resources such as barren lake beds, built-up areas, and croplands. This impact level occurs infrequently and for only short lengths as shown in the IA/MP charts.

Mitigation Planning

Mitigation Goals

Despite the inclusion of generic mitigation measures in the project, there are many moderate to occasional high impact levels along alternative corridors as shown in the IA/MP charts. Selective mitigation measures have been developed by Dames & Moore in consultation with DWP in order to reduce initial impact levels. The goal of these recommended mitigation measures is to eliminate significant construction impacts (very high and high impact levels) and reduce certain other construction impacts (moderate and some low impact levels) to the extent that is reasonable and feasible.

Recommended Selective Mitigation Measures

A number of selective mitigation measures were originally developed by Dames & Moore in consultation with DWP and BLM for the proposed project. Several of these apply directly to the protection of biological resources during construction activities. These can be divided into two general categories: (1) access road restrictions, and (2) avoidance measures. These recommended selective mitigation measures are described below, using the original number designations.

In addition, the application of these mitigation measures to specific resources and access requirements is noted below and summarized on Table 4-14. Selective mitigation measures are applied independently to portions of proposed corridors that have: (1) moderate to very high impact levels, or (2) low impact levels with one or more moderate sensitivity resource.

Access Road Restrictions

1. No new access will be constructed in designated areas; e.g., construction and maintenance will be accomplished without benefit of new access in designated areas. This would minimize ground disturbance, limit new or improved access ability.

This mitigation measure would prohibit the construction of new access roads in areas of high biological sensitivity such as riparian woodlands, desert washes, bighorn sheep habitat, and potential desert tortoise habitat of very high density (greater than 250 individuals per square mile). Access to construction sites would be accomplished by cross-country travel. Hence, there would be no permanent loss of vegetation or wildlife habitat. Residual impacts would consist of disturbance to vegetation by crushing and the temporary displacement of wildlife.

2. No widening or upgrading of existing access roads will be undertaken in the area (same benefits as No. 1 above).

This mitigation measure would be broadly applied to areas of low to high resource sensitivity and moderate or greater impact levels. It only applies to areas where an existing sufficient access road is present, and prohibits any temporary or permanent disturbances due to road improvement activities.

4. All access roads not required for maintenance will be permanently closed using the most effective and least environmentally damaging methods appropriate to that area with concurrence of the landowner. This would limit new or improved accessibility into the area.

This mitigation measure would discourage unauthorized vehicular access and human activity along specified access roads that traverse areas of high biological sensitivity such as riparian woodlands, desert washes, and bighorn sheep habitat. It would only apply to corridors with existing sufficient or upgraded roads. Adverse impacts associated with unauthorized access such as ORV travel and illegal cacti collecting would be reduced by this measure.

Avoidance Measures

8. In designated areas structures will be placed so as to avoid sensitive features and/or to allow conductors to clearly span the features, within limits of standard tower design. This would minimize amount of sensitive feature disturbed and/or reduce visual contrast.

This mitigation measure would be implemented primarily to avoid direct impacts from construction activities on the following maximum and major sensitivity resources: riparian woodlands and desert washes, respectively. These biological resources are usually easily delineated in the field by obvious features and can be avoided during construction activities. This measure would require the minor relocation of tower bases and the spanning of portions of natural waterways.

11. In designated areas construction activities will be modified during breeding season of sensitive, listed, or proposed threatened or endangered species. This would reduce disturbance to sensitive species.

This mitigation measure would consist of two submeasures to avoid direct impacts to desert tortoise and bighorn sheep in specified areas by conducting construction activities during certain seasons. These submeasures are described as follows:

- 11a. Construction activities in known desert bighorn sheep habitat would be limited to the summer and fall months when the sheep are restricted to their small, isolated crucial habitats away from potential disturbances by construction activities.

- 11b. Construction activities in potential desert tortoise habitat with moderate to high densities (100 to 250 individuals per square mile) or very high densities (greater than 250 individuals per square mile) would be restricted, to the extent feasible, to the period of October to March when tortoises are underground and hibernating. Hence, construction activities and vehicular movements on the ground surface would not greatly disturb the tortoise and eliminate mortality by vehicular collisions. If construction activities in these areas occur during portions of the year when tortoises are active, special surveys would be conducted to relocate tortoises from areas of potential disturbance as described in Item 13b.

13. Prior to construction, an ecological field review of tower and access road design will be conducted by an appropriately qualified professional to identify site-specific impacts to threatened, endangered, or otherwise sensitive vegetation and wildlife, and to determine the most effective means to mitigate those impacts. Possible mitigation measures could include minor adjustments in tower and road locations, closing access roads, relocating sensitive species, habitat improvements, etc.

This mitigation measure would be implemented to locate and protect special-interest plants and desert tortoise as described in the two submeasures below:

- 13a. The presence of special-interest plants in areas likely to provide suitable habitat would be evaluated by seasonal surveys prior to construction. Populations in potential disturbance zones would be

identified and the locations of towers, construction yards, and new access roads would be altered to avoid the populations.

- 13b. During construction activities in high and very high density desert tortoise habitat, a qualified biologist would be on-site to locate and collect any hibernating tortoise that would be adversely affected by construction activities. These tortoises would be relocated according to requirements of the BLM and USFWS. Similar, possibly more intensive, surveys would be conducted to identify active burrows and to relocate tortoises from areas of potential disturbance if construction activities occur when tortoises are above ground and active.

Effectiveness of Selective Mitigation Measures

The number and type of selective mitigation measures applied to proposed corridors are dependent on the initial impact level and resources affected. All high initial impact levels are reduced to moderate or low residual impact levels by the application of selective mitigation measures. All moderate impact levels are reduced to low or negligible residual levels, while low impact levels with moderate sensitivity resources are reduced to negligible residual levels. The effectiveness by which selective mitigation measures would reduce the initial impact levels is shown in Table 4-15.

Results: Residual Impact Levels

After the selective application of the recommended mitigation measures following the procedures described above, all initial impact levels have been reduced to moderate or lower residual impact levels. These results are summarized by milepost and link in the IA/MP charts.

All moderate to high initial impact levels due to the potential disturbance of special-interest plants can be mitigated by the pre-construction botanical survey (Measure 13a). All high initial impact levels associated with potential disturbance of riparian woodlands or desert washes can be mitigated by avoidance procedures and restrictions on areas of disturbance (Measures 2 and 8). Mitigation for impacts within bighorn sheep habitat (initial high impact level) include elimination of new access road (Measure 1), closure of existing access roads after construction (Measure 4), and/or a seasonal restriction on construction activities (Measure 11a). Measure 2 is recommended for many corridors in order to reduce moderate to high initial impact levels due to potential disturbance of special-interest plants, desert tortoise habitat, and Mohave ground squirrel habitat. Mitigation 13b is recommended to reduce potentially significant impacts on moderate to very high density desert tortoise habitat. In several special instances, this measure is also recommended for desert tortoise habitat of low to moderate density. In areas with the highest density desert tortoise populations, Measure 1 is also recommended to reduce the high initial impact level.

Conclusions

Based on the above impact assessment the following conclusions are made:

1. Impacts due to operation and maintenance of the proposed project are not expected to result in any significant adverse impacts to biological resources; hence, no specific mitigation measures are recommended. A potential net beneficial impact could occur due to increased perch and nesting sites for raptors on transmission towers.
2. Construction activities represent the only potentially significant impacts to biological resources. The type and intensity of these impacts vary between alternative corridors depending primarily on the sensitivity of the affected resources and the access road requirements.
3. Construction of the proposed project would result in high initial impact levels (i.e., significant impacts) along many alternative corridors, particularly those requiring new access roads.
4. All moderate to very high initial impacts can be feasibly reduced by the application of reasonable and effective selective mitigation measures. Hence, construction of the project along any route is not expected to result in significant impacts to biological resources.

AGENCY CONTACTS

BIOLOGICAL RESOURCES

Agency	Date	Type of Contact	Summary
United States Bureau of Land Management Barstow Resource Area, Barstow, California Daryl Albiston, Natural Resource Specialist	12/5/84	Meeting	Rare plant occurrence in study area and BLM's concerns about project impacts.
California Desert District, Riverside Larry Foreman, District Biologist	10/2/84 to 12/4/84	Meetings/ Telephone	BLM data on biological resources in the study area and BLM's concerns with project impacts.
Kristin Berry, Wildlife Biologist/Desert Tortoise Expert	12/6/84	Telephone	Current data on tortoise distribution in study area.
Las Vegas District, Las Vegas, Nevada Dave Pulliam, Wildlife Biologist	11/27/84	Telephone	Requested data and received referrals to staff members.
Terry Driver, Range Conservationist	12/3/84	Meeting	Rare plant occurrence in the study area and BLM's concerns about project impacts.
Stateline Resource Area, Las Vegas, Nevada Roy Price, Wildlife Biologist	12/3/84	Meeting	BLM data on bighorn sheep, raptors, desert tortoise, and Gila monsters in study area, and BLM's concerns about project impacts.

Agency Contacts (continued)
Biological Resources

Agency	Date	Type of Contact	Summary
United States Department of Fish and Wildlife Service Endangered Species Office Laguna Niguel, California Ray Bransfield, Biologist	12/11/84	Telephone	USFWS's concerns about project impacts.
Karla Kramer, Wildlife Biologist	12/11/84	Telephone	USFWS's data on biological resources in study area and concerns with project impacts; data request.
Sacramento, California Jim Bartel, Botanist	11/27/84	Telephone	USFWS's concern about rare plants in the study area.
Ted Rado, Wildlife Biologist	12/13/84	Letter	Significant biological resources in the study area.
Great Basin Complex, Reno Donald Sada	12/7/84	Letter	Occurrence of endangered species in study area.
Utah, Idaho and Nevada areas, Idaho Bob Parenti, Wildlife Biologist	11/27/84	Telephone	Inform USFWS of project and receive referred to district office.
California Department of Fish and Game, Region 5, Chino Frank Hoover, Fisheries Biologist	12/18/84	Letter	Provided information on desert tortoise and Gila monsters in study area.

Agency Contacts (continued)
Biological Resources

Agency	Date	Type of Contact	Summary
California Department of Fish and Game (continued) Unit 5-7, Victorville Bob Vernoy, Wildlife Biologist	12/1/84	Telephone	CDFG data on biological resources in study area and CDFG's concerns with project impacts.
California Natural Diversity Data Base and California Native Plant Society, Sacramento Rick York, Director, Rare Plants	12/6/84	Telephone	Status of CNPS-listed plants.
Nevada Department of Wildlife, Las Vegas Dan Delaney, Wildlife Biologist/ Bighorn Sheep Biologist	12/3/84	Meeting	Occurrence of bighorn sheep in study area (Nevada) and NDW's concerns about project impacts.
Butch Padilla, Wildlife Biologist	12/3/84	Meeting	NDW data on bighorn sheep, raptors, and desert tortoise in the study area, and NDW's concerns with project impacts.
Ross Haley, Wildlife Biologist	12/27/84	Telephone	Desert tortoise and raptors in the study area.
Nevada Division of Forestry, Carson City Patrick Murphy, Assistant State Forester	11/28/84, 11/29/84	Letter	Occurrence of state-listed plants in the study area.

Agency Contacts (continued)
Biological Resources

Agency	Date	Type of Contact	Summary
Nevada State Museum and Nevada Native Plant Society, Carson City Ann Pinzl, Botanist	11/28/84	Letter	Provided data on rare plants in the study area.
University of California, Davis Department of Wildlife and Fisheries Alice Karl, Wildlife Biologist/Desert Tortoise Biologist	12/4/84		Assisted in field surveys in Nevada, particularly concerning potential desert tortoise habitat.

REFERENCES

- Albiston, D. 1984. Personal communication. Bureau of Land Management, Barstow, California.
- Beatley, J.C. 1976. Vascular Plants of the Nevada Test Site and South-central Nevada. Laboratory of Nuclear Medicine and Radiation Biology, University of California, Los Angeles.
- Benson, L. and R.A. Darrow. 1981. Trees and Shrubs of the Southwestern Deserts. University of Arizona Press, Tucson.
- Berry, K.H. 1974. Desert Tortoise Relocation Project: Status Report for 1982. California Division of Highways Contribution F-9353.
- Bickel, J. 1984. Bighorn Sheep Census in the Clark Mountain Range. Bureau of Land Management. Unpublished report.
- Bloom, P.H. 1979. The Status of the Swainson's Hawk in California, 1979. California Department of Fish and Game. Miscellaneous publication.
- Bradley, W.G. and J.E. Deacon. 1966. Distribution of the Gila Monster in the Northern Mojave Desert.
- Brown, D.E. 1978. Biotic Communities of the Southwest. U.S. Department of Agriculture, Forest Service. General Technical Report RM-78.
- Brown, D.E., editor. 1982. Biotic Communities of the American Southwest-United States and Mexico. Desert Plants 4:1-342.
- Bureau of Land Management. 1984. Miscellaneous in-house files. BLM California Desert District Office, Riverside. Background data to develop CDCA Plan.
- _____. 1980a. Allen-Warner Valley Energy System. Environmental Impact Statement, Final Draft.
- _____. 1980b. The California Desert Conservation Area: Plan Alternatives and Environmental Impact Statement. Draft. U.S. Department of the Interior.
- _____. 1979. IPP. Volume I, Salt Wash Proposal. Draft Environmental Statement, Intermountain Power Project.
- _____. 1974. Proposed Barstow-Las Vegas Motorcycle Race. Final Environmental Impact Statement.
- Burge, B.L. 1977. Movements and Behavior of the Desert Tortoise (Gopherus agassizii). M.S. thesis. University of Nevada, Las Vegas.

References (continued)

- California Department of Fish and Game. 1984. Rare or Endangered Plants. List of July 1984. Sacramento, California.
- _____. 1980. At the Crossroads 1980. A report on California's endangered and rare fish and wildlife. Sacramento, California. Plus 1983 amendments.
- California Natural Diversity Data Base. 1984. Data base reports and maps for California Desert area. In BLM files. California Desert District Office, Riverside, California.
- Cheatham, N.H. and J.R. Haller. 1975. An annotated list of California habitat types. University of California Natural Land and Water Reserves System. Unpublished manuscript. University of California, Santa Barbara.
- Cross, J.K. 1978. Annotated Literature Review on the Ecology of the Gila Monster (Heloderma suspectum). U.S. Bureau of Reclamation Contract 8-07-32-V0040.
- DeDecker, M. 1984. Flora of the Northern Mojave Desert, California. California Native Plant Society, Berkeley, California.
- Delaney, D. 1984. Personal communication. Nevada Department of Wildlife, Las Vegas, Nevada.
- Driver, T. 1984. Personal communication. Bureau of Land Management, Las Vegas, Nevada.
- Faanes, C. 1983. Assessment of Power Line Siting in Relation to Bird Strikes in the Northern Great Plains. U.S. Fish and Wildlife Service, Jamestown, North Dakota.
- Fisher, J. 1980. Field Research of Rare Plants in Las Vegas ES Area, Clark County, Las Vegas, Nevada. Report submitted by WESTEC Services, Inc. to Bureau of Land Management, Las Vegas, Nevada.
- Gaines, D. 1977. Current Status and Habitat Requirements of the Yellow-billed Cuckoo in California. California Department of Fish and Game, Sacramento, California. Miscellaneous report.
- Gaines, D. and S.A. Laymon. 1984. Decline, Status and Preservation of the Yellow-billed Cuckoo in California. *Western Birds* 15:49-80.
- Garrett, K. and J. Dunn. 1981. Birds of Southern California. Status and Distribution. Los Angeles Audubon Society, Los Angeles.

References (continued)

- Goldwasser, S. 1978. Distribution, Reproductive Success, and Impact of Nest Parasitism by Brown-headed Cowbirds on Least Bell's Vireo. California Department of Fish and Game Nongame Wildlife Investigations, Job IV. Final report.
- Goldwasser, S., D. Gaines and S.R. Wilbur. 1980. The Least Bell's Vireo in California - a de facto Endangered Race. *American Birds* 34:742-745.
- Grinnell, J. and A.H. Miller. 1944. The Distribution of the Birds of California. *Pacific Coast Avifauna* 27.
- Hall, R.E. and K.R. Kelson. 1959. *Mammals of North America*. Ronald Press, New York.
- Hubbs, C.L. and R.R. Miller. 1943. Mass Hybridization Between Two Genera of Cyprinid Fishes in the Mohave Desert, California. *Michigan Academy of Science, Arts and Letters* 28:343-378.
- Ingles, L.G. 1965. *Mammals of the Pacific States*. Stanford University Press, Stanford, California.
- Jones and Stokes Associated, Inc. 1980. Assessment of Transmission Line Corridor Impacts on Biological Resources. Report for California Public Utilities Commission.
- Karl, A.E. 1984. Habitat Associations of Vertebrate Wildlife in the Mojave Desert of Southern Nevada. Contract report, University of California, Davis.
- Kuchler, A.W. 1977. Natural Vegetation of California (map). Department of Geography, University of Kansas, Lawrence.
- Luckenback, R.A. 1982. Ecology and Management of the Desert Tortoise (*Gopherus agassizii*) in California. Pp. 1-37.
- McGurty, B.M. 1977. Reptiles and Amphibians of the Eastern Mojave Desert. Contract report to California BLM.
- McQuivey, R.P. 1978. The Desert Bighorn Sheep of Nevada. Nevada Department of Wildlife, Biological Bulletin 6.
- _____. 1976. The Status and Trend of Desert Bighorn Sheep in Nevada. Federal Aid in Wildlife Restoration, Project W-48-7. Special report.
- Mitchell, J.V. 1978. Composition and Abundance of Reptiles and Amphibians of The Clark Mountain Area, San Bernardino, California, May 1978-September 1978. Contract report to California BLM.

References (continued)

- Monson, G. and L. Sumner. 1980. The Desert Bighorn, Its Life History, Ecology and Management. University of Arizona Press, Tucson, Arizona.
- Mozingo, H.N. and M. Williams. 1980. Threatened and Endangered Plants of Nevada. U.S. Department of the Interior. Miscellaneous report.
- Mulder, B. 1978. Yellow-billed Cuckoo (Coccyzus americanus). Bureau of Land Management. Unpublished report.
- Munz, P.A. 1974. A Flora of Southern California. University of California Press, Berkeley, California.
- Nevada Department of Forestry. 1984. Critically Endangered Species List. Nevada Revised Statutes.
- Nevada Department of Wildlife. 1984. List of Rare and Endangered Fish and Game. Nevada Revised Statutes, 501.110.
- NNPS (Northern Nevada Native Plant Society). 1984. Sensitive Plant List of Nevada. Miscellaneous report. Reno, Nevada.
- Pinzl, A. 1984. Letter to J.D. Gray, Dames & Moore, from the Nevada State Museum, Carson City, Nevada.
- Price, R. 1984. Personal communication. Bureau of Land Management, Las Vegas, Nevada.
- Remsen, J.V. 1978. Bird Species of Special Concern in California: An Annotated List of Declining or Vulnerable Bird Species. California Department of Fish and Game, Wildlife Management Branch. Administrative Report 78-1.
- SCE (Southern California Edison). 1981. Ivanpah Generating System. Application for Certification. Biological resources. Report submitted by Bio Systems Analysis, Inc., San Francisco, California.
- Schultz, J.S. 1984. Intermountain Power Project (IPP), $\pm 500\text{kV}$ DC Intermountain-Adelanto Line I, California Portion. Botanical resources inventory and significance evaluation. Final report to Los Angeles Department of Water and Power.
- . 1982. Intermountain Power Project (IPP), $\pm 500\text{kV}$ DC Intermountain-Adelanto Line I (revised), Victorville, California Reroute. Botanical resources inventory and significance evaluation. Final report to Los Angeles Department of Water and Power.

References (continued)

- Smith, J.P. 1982. Inventory of Rare and Endangered Vascular Plants of California, First Supplement. California Native Plant Society.
- . 1981. Inventory of Rare and Endangered Vascular Plants of California, Second Supplement. California Native Plant Society.
- Smith, J.P., R.J. Cole and J.O. Sawyer. 1980. Inventory of Rare and Endangered Vascular Plants of California. California Native Plant Society. Special Publication 1.
- Smith, J.P. and R. York. 1984. Inventory of Rare and Endangered Vascular Plants of California. California Native Plant Society.
- Stebbins, R.C. 1966. A Field Guide to Western Reptiles and Amphibians. Houghton Mifflin Company, Boston, Massachusetts.
- Tate, J. and D.J. Tate. 1982. The Blue List for 1982. American Birds 36:126-135.
- Thelander, C.G. 1974. Bald Eagle Reproduction in California, 1972-1973. California Department of Fish and Game, Wildlife Management Branch, Sacramento, California. Administrative Report 73-5.
- Thorne, R.F. 1976. The Vascular Plant Communities of California. In Plant Communities of Southern California, J. Latting, editor. California Native Plant Society, Berkeley, California. Special Publication 2.
- Thorne, R.F., B.A. Prigge and J. Hendrickson. 1981. A Flora of the Higher Ranges and the Kelso Dunes of the Eastern Mojave Desert in California. Reprinted from: ALISO 10(1), October 1981. Rancho Santa Ana Botanic Garden, Claremont, California.
- U.S. Fish and Wildlife Service. 1984. Endangered and Threatened Wildlife and Plants. Federal Register 48:34182 (1983).
- . 1983. Supplement to Review of Plant Taxa for Listing. Federal Register 48:53640-53670.
- . 1982. Review of Vertebrate Wildlife for Listing as Endangered or Threatened Species. Federal Register 47:58454.
- . 1980. Review of Plant Taxa for Listing as Endangered or Threatened Species. Federal Register 45:82480-82569.
- Vasek, F.C. and M.G. Barbour. 1977. Mojave Desert Scrub Vegetation. In Terrestrial Vegetation of California, M.G. Barbour and J. Major, editors. John Wiley and Sons, New York.

References (continued)

- Weaver, R.A. 1982. Bighorn in California: A Plan to Determine the Current Status and Trend. California Department of Fish and Game, Federal Aid in Wildlife Restoration Project W-51-R. Sacramento, California.
- Weaver, R.A. and J.M. Hall. 1971. Bighorn Sheep in the Clark, Kingston and Nopah Mountain Ranges (San Bernardino and Inyo Counties). California Department of Fish and Game, Wildlife Management Branch. Administrative Report 72-3.
- Weaver, R.A. and J.L. Mensch. 1970. Bighorn Sheep in Northwestern San Bernardino and Southwestern Inyo Counties. California Department of Fish and Game, Wildlife Management Branch. Administrative Report 70-3.
- Weaver, R.A., J.L. Mensch and R.D. Thomas. 1969. A Report on Desert Bighorn Sheep in Northeastern San Bernardino County. California Department of Fish and Game, Federal Aid in Wildlife Restoration Project W-51-R. Sacramento, California.
- Wessman, E.V. 1977. The Distribution and Habitat Preferences of the Mohave Ground Squirrel in the Southeastern Portion of Its Range. California Department of Fish and Game, Wildlife Management Branch, Sacramento, California. Administrative Report 77-S.
- Woodbury, A.M. and R. Hardy. 1948. Studies of the Desert Tortoise, Gopherus agassizii. Ecological Monographs 18:145-200.
- Woodward-Clyde Consultants. 1979. Harry Allen and Warner Valley Generating Projects - Western Transmission Line Corridors, California and Nevada. Environmental Assessment for Southern California Edison Company.

TABLE 4-1
COMMON AND CHARACTERISTIC PLANT SPECIES IN THE STUDY AREA

Scientific Name	Common Name	Growth Form ^a	Vegetation Type ^b							
			CBS	SS	BS	JTW	PJ	DW	RW	MMF
<u>Abies concolor</u> ssp. <u>concolor</u>	Rocky Mountain white fir	T								X
<u>Acacia greggii</u>	catclaw	S/T							X	
<u>Acamptopappus schokleyi</u>	goldenhead	SS	X	X	X	X				
<u>Agave utahensis nevadensis</u>	Utah agave	S		X	X		X			X
<u>Allenrolfea occidentalis</u>	pickleweed	PH							X	X
<u>Amelanchier utahensis</u>	-	S						X		X
<u>Ambrosia dumosa</u>	burrobush	SS	X	X		X			X	
<u>Argemone corymbosa</u>	prickly poppy	PH	X							
<u>Artemisia nova</u>	dwarf sagebrush	S			X			X		
<u>A. tridentata</u>	Great Basin sagebrush	S			X			X		X
<u>Astragalus</u> spp.	locoweed	PH	X	X	X	X	X	X	X	
<u>Atriplex canescens</u>	four-winged saltbrush	S	X	X		X	X	X	X	
<u>A. confertifolia</u>	shadscale	S	X	X		X	X	X	X	
<u>A. hymenelytra</u>	desert holly	S	X	X		X	X			
<u>A. polycarpa</u>	all-scale	S	X	X				X	X	
<u>Baccharis glutinosa</u>	mule fat	S							X	X
<u>B. sergiloides</u>	waterweed	S							X	X
<u>Bassia hyssopifolia</u>	five hook bassia	PH							X	X
<u>Bebbia juncea</u>	sweet bush	SS							X	

Table 4-1 (continued)
Common and Characteristic Plant Species in the Study Area

Scientific Name	Common Name	Growth Form ^a	Vegetation Type ^b							
			CBS	SS	BS	JTW	PJ	DW	RW	MMF
<u>Bouteloua</u> spp.	gramma grass	PG	X		X	X	X			
<u>Brickellia arguta</u>	bricklebrush	S						X		
<u>Cassia armata</u>	desert senna	S	X	X						
<u>Cercocarpus ledifolius</u>	mountain mahogany	S					X			X
<u>Chilopsis linearis</u>	desert willow	S						X	X	
<u>Chrysothamnus nauseosus</u>	rabbit brush	S			X	X				X
<u>Coleogyne ramosissima</u>	blackbush	SS			X	X	X			
<u>Coryphantha vivipara</u> <u>ssp. deserti</u>	desert coryphantha	C								
<u>Cowania mexicana</u>	cliffrose	S			X			X		
<u>Croton californica</u>	desert croton	PH	X	X	X	X	X			
<u>Dalea spinosa</u>	smoke tree	S						X		
<u>D. fremontii</u>	indigo bush	S			X					
<u>Distichlis spicata</u>	salt grass	PG						X	X	
<u>Echinocactus polycephalus</u>	many-headed barrel cactus	C			X					
<u>Echinocereus engelmannii</u>	hedgehog cactus	C	X	X	X					
<u>Encelia farinosa</u>	brittlebrush	S	X	X						
<u>Ephedra nevadensis</u>	morman tea	S			X					
<u>E. viridis</u>	green ephedra	S	X		X					

Table 4-1 (continued)
Common and Characteristic Plant Species in the Study Area

Scientific Name	Common Name	Growth Form ^a	Vegetation Type ^b						
			CBS	SS	BS	JTW	PJ	DW	RW
<u>Erionueron pulchellum</u>	fluff grass	PG	X	X	X	X			
<u>E. fasciculatum</u>	buckwheat	S	X		X	X	X	X	
<u>E. inflatum</u>	desert trumpet	PH	X	X					
<u>Erodium cicutarium</u>	filaree	AH	X						
<u>Eurotia lanata</u>	winter fat	PH	X	X	X				
<u>Fallugia paradoxa</u>	apache plume	S					X		X
<u>Ferocactus acanthoides</u>	barrel cactus	C	X		X	X			
<u>Grayia spinosa</u>	hopsage	S	X		X	X			
<u>Gutierrezia microcephala</u>	matchweed	SS	X	X	X	X	X	X	
<u>Haplopappus linearifolius</u>	goldenbush	S			X		X		
<u>H. cooperi</u>	cooper goldenbush	S	X		X	X			
<u>Hilaria rigida</u>	big galleta	PG	X		X	X			
<u>Hymenoclea salsola</u>	cheesebush	S	X	X		X		X	
<u>Isomeris arborea</u>	bladder pod	S	X	X		X			
<u>Juncus mexicanus</u>	rush	S/T						X	X
<u>Juniperus osteosperma</u>	Utah juniper	PH			X		X		X
<u>Krameria parvifolia</u>	little-leaved ratany	S	X	X	X	X			
<u>Larrea tridentata</u>	creosote bush	S	X	X	X	X			
<u>Lycium andersonii</u>	box thorn	S	X	X	X	X	X		

Table 4-1 (continued)
Common and Characteristic Plant Species in the Study Area

Scientific Name	Common Name	Growth Form ^a	Vegetation Type ^b						
			CBS	SS	BS	JTW	PJ	DW	RW
<u>Menodora spinescens</u>	spiny menodora	SS	X		X	X	X		
<u>Muhlenbergia porteri</u>	bush muhly	PG	X	X	X				
<u>Nicotiana trigonophylla</u>	desert tobacco	PH						X	
<u>Nitrophila occidentalis</u>	alkali weed	PH						X	
<u>Oenothera deltoides</u>	desert primrose	PH	X	X	X	X			
<u>Opuntia acanthocarpa</u>	buckhorn cholla	C	X	X	X	X		X	
<u>O. basilaris</u>	beavertail cactus	C	X			X			
<u>O. bigelovii</u>	jumping cholla	C	X			X			
<u>O. echinocarpa</u>	silver cholla	C	X		X	X			
<u>O. erinacea</u>	Mojave prickly pear	C	X	X		X	X		X
<u>O. ramosissima</u>	pencil cholla	C			X				
<u>Oryzopsis hymenoides</u>	Indian rice grass	PG	X	X	X	X	X	X	
<u>Physalis</u> spp.	ground cherry	PH	X						
<u>Pinus monophylla</u>	one-needle pinyon	T					X		X
<u>Populus fremontii</u>	desert cottonwood	T						X	X
<u>Prosopis glandulosa</u>	honey mesquite	T						X	
<u>Prunus fremontii</u>	desert almond	S	X				X	X	
<u>Psilostrophe cooperi</u>	paper flower	S	X	X		X			
<u>Rhus trilobata</u>	squaw bush	S					X		X

Table 4-1 (continued)
Common and Characteristic Plant Species in the Study Area

Scientific Name	Common Name	Growth Form ^a	Vegetation Type ^b						
			CBS	SS	BS	JTW	PJ	DW	RW
<u>Salazaria mexicana</u>	bladder sage	S	X	X		X			
<u>Salix</u> spp.	willow	T							X
<u>Salsola iberica</u>	Russian thistle	PH	X					X	X
<u>Salicornia</u> spp.	glasswort	PH						X	
<u>Sarcobatus vermiculatus</u>	greasewood	PH						X	
<u>Scirpus</u> spp.	bulrush	PH							X
<u>Sphaeralcea ambigua</u>	desert globe mallow	AH	X	X					
<u>Sporobolus</u> spp.	drop seed	PG				X	X		
<u>Suaeda</u> spp.	seep weed	PH						X	
<u>Tamarix</u> spp.	tamarisk	T						X	X
<u>Tetradymia spinosa</u>	cottonthorn	S	X	X	X	X			
<u>Thamnosma montana</u>	turpentine bush	S	X		X				
<u>Typha latifolia</u>	cattail	PH							X
<u>Yucca baccata</u>	Spanish bayonet	S	X		X				
<u>Y. brevifolia</u>	Joshua tree	S	X		X	X	X		
<u>Y. schidigera</u>	Mojave yucca	S	X		X	X	X		

^aT = tree; S = shrub; SS = subshrub; PH = perennial herb; PG = perennial grass; AH = annual herb; C = cactus.

^bCBS = creosote bush scrub; SS = saltbush scrub; BS = blackbush scrub; JTW = Joshua tree woodland; PJ = pinyon-juniper woodland; DW = desert wash and alkali basin vegetation; RW = riparian woodland; MMF = Mojave montane forest.

TABLE 4-2
COMMON AND CHARACTERISTIC AMPHIBIANS AND REPTILES
OF THE STUDY AREA^a

Common Name	Scientific Name	Characteristic Habitat ^b			
		SH	WH	MH	RH
Couch's spadefoot toad	<u>Scaphiopus couchi</u>	X			
Bullfrog	<u>Rana catesbeiana</u>				X
Northern leopard frog	<u>Rana pipiens</u>				X
Great plains toad	<u>Bufo cognatus</u>				X
Southwestern toad	<u>Bufo microscaphus</u>				X
Red-spotted toad	<u>Bufo punctatus</u>				X
Woodhouse's toad	<u>Bufo woodhousei</u>				X
Pacific treefrog	<u>Hyla regilla</u>				X
Desert tortoise	<u>Scaptochyles agassizii</u>	X			
Desert gecko	<u>Coleonyx v. variegatus</u>	X			
Zebra-tailed lizard	<u>Callisaurus draconoides</u>	X			
Desert-collared lizard	<u>Crotaphytus insularis</u> <u>bicinctores</u>	X			
Desert iguana	<u>Dipsosaurus dorsalis</u>	X			
Long-nosed leopard lizard	<u>Gambelia w. wislizenii</u>	X			
Desert-horned lizard	<u>Phrynosoma platyrhinos</u>	X			X
Western chuckwalla	<u>Sauromalus o. obesus</u>	X			
Sagebrush lizard	<u>Sceloporus graciosus</u>		X	X	
Desert spiny lizard	<u>Sceloporus magister</u>	X	X		
Mojave fringe-toed lizard	<u>Uma scoparia</u>	X			
Long-tailed brush lizard	<u>Urosaurus graciosus</u>	X			
Side-blotched lizard	<u>Uta stansburiana</u>	X	X		X
Banded Gila monster	<u>Heloderma suspectum</u> <u>cinctum</u>	X			
Desert night lizard	<u>Xantusia c. virgilis</u>		X		
Great basin whiptail	<u>Cnemidophorus t. tigris</u>	X	X		

Table 4-2 (continued)
Common and Characteristic Amphibians and Reptiles of the Study Area

<u>Common Name</u>	<u>Scientific Name</u>	<u>Characteristic Habitat^b</u>			
		<u>SH</u>	<u>WH</u>	<u>MH</u>	<u>RH</u>
Western red-tailed skink	<u>Eumeces gilberti</u> <u>rubricandatus</u>		X	X	
Great basin skink	<u>Eumeces skiltonianus</u> <u>utahensis</u>		X	X	
Western blind snake	<u>Leptotyphlops humilis</u>	X			
Rosy boa	<u>Lichanura trivirgata</u>		X		X
Glossy snake	<u>Arizona elegans</u>	X			
Western shovel-nosed snake	<u>Chionactis occipitalis</u>	X			
Racer	<u>Coluber constrictor</u>		X	X	
Regal ringneck snake	<u>Diadophis punctatus</u> <u>regalis</u>		X	X	X
Desert night snake	<u>Hypsiglena torquata</u> <u>deerticola</u>	X			X
Common kingsnake	<u>Lampropeltis getulus</u>	X			
Coachwhip	<u>Masticophis flagellum</u>	X			
Striped whipsnake	<u>Masticophis taeniatus</u>	X	X		
Spotted leaf-nosed snake	<u>Phyllorhynchus</u> <u>decurtatus</u>	X			X
Great Basin gopher snake	<u>Pituophis melanoleucus</u> <u>deserticola</u>	X	X	X	X
Long-nosed snake	<u>Rhinocheilus lecontei</u>	X			
Western patch-nosed snake	<u>Salvadora hexalepis</u>	X			
Ground snake	<u>Sonora semiannulata</u>	X			X
Western black-headed snake	<u>Tantilla planiceps</u>	X			X
Lyre snake	<u>Trimorphodon biscutatus</u>	X	X	X	
Western diamondback rattlesnake	<u>Crotalus atrox</u>	X			X
Sidewinder	<u>Crotalus cerastes</u>	X			

Table 4-2 (continued)

Common and Characteristic Amphibians and Reptiles of the Study Area

Common Name	Scientific Name	Characteristic Habitat ^b			
		SH	WH	MH	RH
Speckled rattlesnake	<u>Crotalus mitchelli</u>	X	X		
Mojave rattlesnake	<u>Crotalus scutellatus</u>	X	X		

^aThese are common and characteristic reptiles and amphibians of the study area; this is not intended to be a comprehensive herpetofaunal list.

^bHabitat types are described in the text under Wildlife. SH = scrub habitats (creosote bush, blackbush, and saltbush scrub); WH = woodland habitats (Joshua tree and pinyon-juniper woodlands); MH = montane habitats; and RH = riparian woodland, desert wash, and wetland habitats.

TABLE 4-3
COMMON AND CHARACTERISTIC BIRDS OF THE STUDY AREA^a

Common Name	Characteristic Habitat ^b				Seasonal Occurrence ^c	Abundance in Study Area ^d
	SH	WH	MH	RH		
Common Loon				X	M	V
Pied-billed Grebe				X	R	C
Eared Grebe				X	M	R
Western Grebe				X	M	R
American White Pelican				X	M	R
Double-crested Cormorant				X	M	V
American Bittern				X	R	C
Least Bittern				X	M	R
Great Blue Heron				X	R	C
Great Egret				X	M	R
Snowy Egret				X	M	R
Green-backed Heron				X	R	R
Black-crowned Night Heron				X	M	R
White-faced Ibis				X	M	R
Tundra Swan				X	M	R
Brant				X	M	V
Canada Goose				X	M	R
Green-winged Teal				X	W, M	U
Mallard				X	R	C
Northern Pintail				X	W, M	U
Blue-winged Teal				X	M	R
Cinnamon Teal				X	R	C
Northern Shoveler				X	M, W	R
Gadwall				X	W	U
Eurasian Wigeon				X	M	U
American Wigeon				X	W	U
Canvas Back				X	M	R
Redhead				X	W	U

Table 4-3 (continued)
Common and Characteristic Birds of the Study Area

Common Name	Characteristic Habitat ^b				Seasonal Occurrence ^c	Abundance in Study Area ^d
	SH	WH	MH	RH		
Ring-necked Duck				X	M	U
Oldsquaw				X	M	R
Common Golden Eye				X	M	R
Bufflehead				X	M	R
Common Merganser				X	M, W	R
Red-breasted Merganser				X	M	R
Ruddy Duck				X	R, M	C
Turkey Vulture	X	X	X	X	M, S	C
Osprey				X	M	R
Black-shouldered Kite	X			X	M	R
Northern Harrier	X			X	R	R
Sharp-shinned Hawk		X		X	R	R
Cooper's Hawk		X		X	R	U
Red-shouldered Hawk		X		X	C	R
Swainson's Hawk	X			X	S	R
Red-tailed Hawk	X	X	X	X	R	U
Ferruginous Hawk	X			X	M, W	U
Golden Eagle	X	X	X	X	R	R
American Kestrel	X	X	X	X	R	C
Merlin		X		X	W	R
Prairie Falcon	X	X	X	X	R	R
Ring-necked Pheasant				X	R	U
Gambel's Quail	X			X	R	C
Virginia Rail				X	S	R
Sora				X	R	U
Common Moorhen				X	R	U
American Coot				X	R	C
Sandhill Crane				X	M	R

Table 4-3 (continued)
Common and Characteristic Birds of the Study Area

Common Name	Characteristic Habitat ^b				Seasonal Occurrence ^c	Abundance in Study Area ^d
	SH	WH	MH	RH		
Black-bellied Plover			X	X	M	R
Semipalmated Plover		X		X	M	R
Killdeer	X	X	X	X	R	C
Black-necked Stilt	X			X	M	R
American Avocet				X	M	R
Greater Yellowlegs	X			X	M	U
Lessor Yellowlegs	X			X	M	U
Solitary Sandpiper	X			X	S	U
Willet		X		X	S	U
Spotted Sandpiper			X	X	S	U
Whimbrel				X	M	U
Long-billed Curlew		X		X	W	R
Sandering	X	X		X	M	U
Western Sandpiper		X	X	X	M	U
Least Sandpiper				X	M	U
Dunlin		X		X	M	U
Long-billed Dowitcher		X		X	M	R
Common Snipe				X	W	U
Red-necked Phalarope		X		X	M	U
Ring-billed Gull		X		X	M	U
California Gull				X	M, W	R
Forster's Tern				X	S	U
Black Tern				X	S	U
Rock Dove				X	R	C
White-winged Dove				X	S	R
Mourning Dove	X	X	X	X	R	C
Greater Roadrunner	X			X	R	U
Barn Owl	X	X	X	X	R	U

Table 4-3 (continued)
Common and Characteristic Birds of the Study Area

Common Name	Characteristic Habitat ^b				Seasonal Occurrence ^c	Abundance in Study Area ^d
	SH	WH	MH	RH		
Flammulated Owl			X		R	R
Western Screech Owl		X		X	R	U
Great Horned Owl	X	X	X	X	R	U
Burrowing Owl	X				R	R
Long-eared Owl				X	R	U
Short-eared Owl	X			X	R	V
Lesser Nighthawk	X			X	S	U
Common Poorwill	X				S	U
Vaux's Swift		X			S	U
White-throated Swift			X	X	S	U
Black-chinned Hummingbird				X	S	U
Anna's Hummingbird		X		X	R	U
Costa's Hummingbird	X	X		X	S	R
Rufous Hummingbird		X	X		S	U
Belted Kingfisher				X	R	C
Yellow-bellied Sapsucker		X		X	W	R
Ladder-backed Woodpecker		X		X	R	C
Nuttall's Woodpecker				X	R	U
Downy Woodpecker		X		X	R	C
Hairy Woodpecker		X		X	R	U
Northern Flicker				X	R	C
Olive-sided Flycatcher			X		M	U
Western Wood-Pewee		X	X	X	S	U
Willow Flycatcher				X	M	U
Hammond's Flycatcher		X	X	X	M	U
Dusky Flycatcher		X	X	X	M	U
Gray Flycatcher		X		X	M	R
Western Flycatcher				X	S	C

Table 4-3 (continued)
Common and Characteristic Birds of the Study Area

Common Name	Characteristic Habitat ^b				Seasonal Occurrence ^c	Abundance in Study Area ^d
	SH	WH	MH	RH		
Black Phoebe				X	R	C
Sam's Phoebe				X	M	U
Vermilion Flycatcher				X	W	R
Ash-throated Flycatcher		X			S	U
Brown-crested Flycatcher				X	S	U
Western Kingbird		X		X	S	C
Horned Lark	X				R	C
Purple Martin				X	S	U
Tree Swallow				X	S	U
Violet-Green Swallow		X			S	U
Northern Rough-winged Swallow				X	S	C
Bank Swallow				X	S	U
Cliff Swallow				X	S	C
Barn Swallow				X	S	C
Scrub Jay		X		X	R	R
Pinyon Jay		X			R	C
Common Raven	X	X	X	X	R	C
Mountain Chickadee			X		R	C
Plain Titmouse		X	X		R	C
Verdin	X			X	R	C
Bushtit		X		X	R	C
Red-breasted Nuthatch			X	X	W	R
White-breasted Nuthatch			X		S	V
Brown Creeper		X		X	R	R
Cactus Wren	X	X			R	C
Rock Wren	X	X	X	X	R	C
Canyon Wren		X	X	X	R	U

Table 4-3 (continued)
Common and Characteristic Birds of the Study Area

Common Name	Characteristic Habitat ^b				Seasonal Occurrence ^c	Abundance in Study Area ^d
	SH	WH	MH	RH		
Bewick's Wren	X	X		X	R	C
House Wren			X	X	S	C
Marsh Wren				X	R	C
Ruby-crowned Kinglet				X	W	U
Blue-Gray Gnatcatcher				X	R	R
Black-tailed Gnatcatcher				X	R	U
Western Bluebird		X		X	R	C
Mountain Bluebird			X		R	U
Townsend's Solitaire		X	X		W	R
Swainson's Thrush				X	M	U
Hermit Thrush				X	W, M	U
American Robin		X	X	X	W, M	C
Varied Thrush			X	X	W	U
Northern Mockingbird	X	X		X	R	C
Sage Thrasher	X				M	U
Bendire's Thrasher		X			S	C
California Thrasher				X	M	R
LeConte's Thrasher	X				R	R
Water Pipit				X	W	C
Bohemian Waxwing			X		M, W	U
Cedar Waxwing				X	W	R
Painopepla				X	R	U
Loggerhead Shrike	X	X		X	R	C
European Starling				X	R	C
Bell's Vireo				X	S	U
Gray Vireo	X	X			S	R
Solitary Vireo			X		S	R
Warbling Vireo				X	S	U

Table 4-3 (continued)
Common and Characteristic Birds of the Study Area

Common Name	Characteristic Habitat ^b				Seasonal Occurrence ^c	Abundance in Study Area ^d
	SH	WH	MH	RH		
Orange-crowned Warbler				X	S	C
Nashville Warbler			X		S	U
Virginia's Warbler			X		S	V
Lucy's Warbler				X	S	R
Yellow Warbler				X	S	U
Yellow-rumped Warbler	X	X		X	W	C
Black-throated Gray Warbler		X	X		M	U
Townsend's Warbler	X	X		X	M	U
Hermit Warbler				X	M	U
American Redstart				X	S	V
MacGillivray's Warbler			X		M, S	R
Common Yellowthroat				X	S	U
Wilson's Warbler				X	M	C
Painted Redstart			X		M, S	R
Yellow-breasted Chat			X		S	U
Hepatic Tanager			X		R	R
Summer Tanager				X	M	U
Western Tanager			X		S	C
Rose-breasted Grosbeak			X		M	R
Black-headed Grosbeak			X		S	C
Blue Grosbeak				X	S	U
Lazuli Bunting				X	S	R
Green-tailed Towhee			X		S	C
Rufous-sided Towhee			X		R	C
Rufous-crowned Sparrow			X		R	R
Chipping Sparrow		X		X	M	R
Clay-colored Sparrow				X	M	V

Table 4-3 (continued)
Common and Characteristic Birds of the Study Area

Common Name	Characteristic Habitat ^b				Seasonal Occurrence ^c	Abundance in Study Area ^d
	SH	WH	MH	RH		
Brewer's Sparrow			X		R	C
Black-chinned Sparrow			X		S	C
Vesper Sparrow	X				W	C
Lark Sparrow				X	R	C
Black-throated Sparrow	X				R	C
Sage Sparrow	X				R	U
Savannah Sparrow				X	R	C
Grasshopper Sparrow	X				S	U
Fox Sparrow				X	M	R
Song Sparrow				X	R	C
Golden-crowned Sparrow		X		X	W	U
White-crowned Sparrow	X	X			W	C
Dark-eyed Junco	X	X	X	X	W	C
Red-winged Blackbird				X	R	C
Tri-colored Blackbird				X	M	R
Western Meadowlark				X	R	C
Yellow-headed Blackbird				X	M, S	R
Brewer's Blackbird		X	X	X	R	C
Brown-headed Cowbird		X	X	X	R	U
Hooded Oriole				X	S	R
Northern Oriole		X		X	S	C
Scott's Oriole		X			S	C
Cassin's Finch			X		W	R
House Finch	X			X	R	C
Lesser Goldfinch				X	R	C
Lawrence's Goldfinch		X	X	X	M	R
American Goldfinch				X	R	U

Table 4-3 (continued)
Common and Characteristic Birds of the Study Area

Common Name	Characteristic Habitat ^b				Seasonal Occurrence ^c	Abundance in Study Area ^d
	SH	WH	MH	RH		
Evening Grosbeak			X	X	W	R
House Sparrow				X	R	C

^aThese are common and characteristic birds of the study area; this is not intended to be a comprehensive avifauna list.

^bHabitat types are described in the text. SH = scrub habitats (creosote bush, blackbush, and saltbush scrub); WH = woodland habitats (Joshua tree and pinyon-juniper woodlands); MH = montane habitats (white fir forest); and RH = riparian woodland, desert wash, and other wetland habitats.

^cR = resident, M = spring and/or fall migrant, W = winter visitor, S = summer visitor.

^dC = common, U = uncommon, R = rare, V = very rare.

^eObserved during field surveys, November 30 to December 8, 1984.

TABLE 4-4
COMMON AND CHARACTERISTIC MAMMALS OF THE STUDY AREA^a

Common Name	Scientific Name	Characteristic Habitat ^b			
		SH	WH	MH	RH
Merriam's shrew	<u>Sorex merriami</u>		X	X	X
Desert shrew	<u>Notiosorex crawfordi</u>	X			X
Leaf-nosed bat	<u>Macrotus californicus</u>	X			X
Yuma myotis	<u>Myotis yumanensis</u>	X			X
Long-eared myotis	<u>Myotis evotis</u>			X	
Fringe-tailed myotis	<u>Myotis thysanodes</u>	X	X		X
Long-legged myotis	<u>Myotis volans</u>			X	
California myotis	<u>Myotis californicus</u>			X	
Small-footed myotis	<u>Myotis subulatus</u>		X	X	
Silver-haired bat	<u>Lasionycteris noctivagans</u>			X	X
Western pipistrel	<u>Pipistrellus hesperus</u>	X	X		X
Big brown bat	<u>Eptesicus fuscus</u>		X		X
Hoary bat	<u>Lasiurus cinereus</u>	X	X	X	
Spotted bat	<u>Euderma maculatum</u>		X	X	
Pallid bat	<u>Antrozous pallidus</u>	X	X		X
Brazilian free-tailed bat	<u>Tadarida brasiliensis</u>	X	X		
Big free-tailed bat	<u>Tadarida molassa</u>	X	X		
Mountain cottontail	<u>Sylvilagus nuttallii</u>		X	X	
Desert cottontail	<u>Sylvilagus audubonii</u>	X	X		
Black-tailed jackrabbit	<u>Lepus californicus</u>	X			
Panamint chipmunk	<u>Tamias panamintus</u>		X		
White-tailed antelope ground squirrel	<u>Ammospermophilus leucurus</u>	X			
Mohave ground squirrel	<u>Spermophilus mohavensis</u>	X			
Round-tailed ground squirrel	<u>Spermophilus tereticaudus</u>	X			
Botta's pocket gopher	<u>Thomomys bottae</u>			X	
Little pocket mouse	<u>Perognathus longimembris</u>	X	X		
San Joaquin pocket mouse	<u>Perognathus inornatus</u>	X			

Table 4-4 (continued)
Common and Characteristic Mammals of the Study Area

Common Name	Scientific Name	Characteristic Habitat ^b			
		SH	WH	MH	RH
Long-tailed pocket mouse	<u>Perognathus formosus</u>	X			
Desert pocket mouse	<u>Perognathus penicillatus</u>	X			
Panamint kangaroo rat	<u>Dipodomys panamintinus</u>	X	X		
Merrian's kangaroo rat	<u>Dipodomys merriami</u>	X			
Desert kangaroo rat	<u>Dipodomys deserti</u>	X			
Western harvest mouse	<u>Reithrodontomys megalotis</u>				X
Canyon mouse	<u>Peromyscus erinitus</u>		X		
Cactus mouse	<u>Peromyscus eremicus</u>	X	X		
Deer mouse	<u>Peromyscus maniculatus</u>	X	X	X	X
Brush mouse	<u>Peromyscus boylii</u>		X		
Pinon mouse	<u>Peromyscus truei</u>		X		
Southern grasshopper mouse	<u>Onychomys torridus</u>	X			
Desert woodrat	<u>Neotoma lepida</u>	X	X		
Mohave River vole	<u>Microtus californicus mohavensis</u>				X
House mouse	<u>Mus musculus</u>	X	X		X
Coyote	<u>Canis latrans</u>	X	X	X	X
Kit fox	<u>Vulpes macrotis</u>	X	X		
Gray fox	<u>Urocyon cinereo argenteus</u>		X		
Ringtail	<u>Bassariscus astutus</u>		X		
Badger	<u>Taxidea taxus</u>	X	X	X	
Spotted skunk	<u>Spilogale putorius</u>	X	X		
Striped skunk	<u>Mephitis mephitis</u>				X
Bobcat	<u>Felis rufus</u>	X	X	X	
Mule deer	<u>Odocoileus hemionus</u>		X	X	
Nelson's bighorn sheep	<u>Ovis canadensis nelsoni</u>			X	

^aThese are common and characteristic mammals of the study area. This is not intended to be a comprehensive faunal list.

Table 4-4 (continued)
Common and Characteristic Mammals of the Study Area

<u>Common Name</u>	<u>Scientific Name</u>	<u>Characteristic Habitat^b</u>			
		<u>SH</u>	<u>WH</u>	<u>MH</u>	<u>RH</u>

Footnotes (continued)

^bHabitat types are described in the text. SH = scrub habitats (creosote bush, blackbush, and saltbush scrub); WH = woodland habitats (Joshua tree and pinyon-juniper woodlands); MH = montane habitats (white fir forest); and RH = riparian woodland, desert wash, and wetland habitats.

TABLE 4-5
PLANT SPECIES OF SPECIAL INTEREST IN THE STUDY AREA

Scientific Name ^a	Common Name ^a	Growth Form ^b	Flowering Period	Federal Candidate Status ^c 1980 1983	BLM Sensitive Species ^d	Nevada Native Plant Society List ^e	California Native Plant Society List ^f 1980 1984	Characteristic Vegetation Type ^g	Regional Distribution	Known or Potential Occurrence in the Study Area ^h
<u>Astragalus jaegerianus</u>	Jaeger milkvetch	PH	April-June	1 2	CA	-	2 IB	CBS, JTW; 300-3000 ft.	CA; 30 miles northeast of Yermo on granite hill-sides; approximately 10 square miles on Coolgardie Mesa between Barstow and Goldstone.	Known populations on Coolgardie Mesa north of Barstow; likely to occur in scattered locations in Mud Hills, on Lane Mountain, and possibly in Calico Mountains.
<u>Chorizanthe spinosa</u>	Mojave spine-flower	AH	April-July	1 2	CA	-	2 4	CBS, JTW; 2500-3500 ft.	CA; alluvial fans, sandy washes, and flats; scattered populations in western Mojave Desert, especially on Edwards Air Force Base.	Known populations near Kramer Junction and Adelanto; likely to occur in scattered areas in western portion of study area.
<u>Coryphantha vivipara</u> var. <u>alversonii</u>	foxtail cactus	C	May-June	2 2	CA	-	2 IB	CBS, JTW; 2000-5000 ft.	CA; stony slopes; Little San Bernardino Mountains to Eagle and Chuckwalla mountains.	Expected to be very rare in the study area; only one site known on Clark Mountain.
<u>Cymopterus deserticola</u>	desert cymopterus	PH	March-May	1 2	-	-	2 IB	CBS, JTW; 2300-4300 ft.	CA; very rare in flats of old dune areas from near Victorville to Kramer Junction and Edwards Air Force Base.	Known populations at Kramer Junction and Apple Valley; possible occurrence in small localized site in western portion of study area.

Table 4-5 (continued)
Plant Species of Special Interest in the Study Area

Scientific Name ^a	Common Name ^a	Growth Form ^b	Flowering Period	Federal Candidate Status ^c 1980 1983	BLM Sensitive Species ^d	Nevada Native Plant Society List ^e	California Native Plant Society List ^f 1980 1984	Characteristic Vegetation Type ^g	Regional Distribution	Known or Potential Occurrence in the Study Area ^h
<u>Eriophyllum mohavensis</u>	Barstow woolly sunflower	AH	April-May	1 2	CA	-	2 IB	CBS; 2000-3000 ft.	CA; sandy or rocky places; flat or sloping; near Barstow and Kramer Junction; very rare.	Known from small site near Barstow and larger populations northeast of Kramer Junction; possible small populations in western portion of study area.
<u>Penstemon bicolor</u> ssp. <u>bicolor</u> and <u>roseus</u>	bicolored penstemon	PH	May-June	1 2	-	W	-	CBS; 2000-5000 ft.	NV; washes and roadsides, southeast slope of Spring Mountains, Kyle Canyon south to Jean and McCullough Mountains.	Known populations in McCullough Mountains; possible additional populations nearby.
<u>Phacelia nelsonii</u>	Nelson's phacelia	AH	April-May	2 2	-	OR	4 2	CBS, BS, JTW, PJ; 2000-6000 ft.	CA and NV; wide distribution on disturbed and rocky soils; washes, slope bases, and talus; small local rare populations.	Known population site in McCullough Mountains; possible but unlikely in other areas of McCullough and Clark mountains.
<u>Puccinella parishii</u>	Parish's alkali grass	AG	April-May	1 2	CA	-	2 IB	JTW; CBS; 2900 ft.	CA; alkaline seep at Rabbit Springs, Lucerne Valley; possible at Kramer Junction and Edwards Air Force Base.	Known but suspect population near Kramer Junction; possible but unlikely additional small localized populations in western study area.

Table 4-5 (continued)
Plant Species of Special Interest in the Study Area

Scientific Name ^a	Common Name ^a	Growth Form ^b	Flowering Period	Federal Candidate Status ^c 1980-1983	BLM Sensitive Species ^d	Nevada Native Plant Society List ^e	California Native Plant Society List ^f 1980-1984	Characteristic Vegetation Type ^g	Regional Distribution	Known or Potential Occurrence in the Study Area ^h
<u>Sclerocactus polyancistrus</u>	Mojave fishhook cactus	C	April-June	1	2 CA	W	2	4 CBS, SS, PJ; 2500-6500 ft.	CA and NV (Nye County); flats, hillsides, and steep slopes from Red Rock Canyon northeast to Nevada in southwest San Bernardino and Inyo counties.	Several known populations in western portion of study area; additional small sites possible nearby.
<u>Sphaeraleca rusbyi</u> ssp. <u>eremicola</u>	Rusby's desert mallow	AH	May-June	1	2 CA	-	2	IB CBS, BS; 3000-4500 ft.	CA; Panamint Mountains (Inyo County) and one site in Clark Mountains; rare.	One known population in the Clark Mountains, Keaney Pass; additional populations are possible in nearby areas.

^a Scientific and common names follow USFWS (1980, 1983), Munz (1974), Smith et al. (1980, 1981, 1982), and Mozingo and Williams (1980).

^b PH = perennial herb, AH = annual herb, C = cactus.

^c Federal candidate status from USFWS (1980, 1983).

Category 1 = taxa for which sufficient information is available to support the biological appropriateness for official listing.

Category 2 = taxa for which sufficient information is available to indicate possible listing, but for which additional data are required on vulnerability and threats. Many species were placed in lower candidate categories in the updated 1983 listing (USFWS 1983) based on new information that indicated their distribution was more widespread than previously believed.

^d California BLM sensitive species from BLM (1984). No Nevada BLM sensitive species list has been developed.

Table 4-5 (continued)
Plant Species of Special Interest in the Study Area

Scientific Name ^a	Common Name ^a	Growth Form ^b	Flowering Period	Federal Candidate Status ^c 1980-1983	BLM Sensitive Species ^d	Nevada Native Plant Society List ^e	California Native Plant Society 1980-1984	Characteristic Vegetation Type ^g	Regional Distribution	Known or Potential Occurrence in the Study Area ^h
------------------------------	--------------------------	--------------------------	------------------	---	------------------------------------	---	---	---	-----------------------	--

Footnotes (continued)

^e Sensitive species list developed annually by NNPS (1984) which includes three categories in decreasing order of rarity and/or endangerment: (1) sensitive species, (2) watch list (w), and (3) other rare species (OR).

^f From the 1980 CNPS Rare Plant Inventory (Smith et al. 1980, 1981, 1982) and 1984 Inventory (Smith and York 1984). The 1980 categories include: List 2 = rare and endangered; List 3 = rare, but not endangered; and List 4 = rare in California, but common elsewhere. The 1984 categories include: List 1B = rare and endangered; List 2 = plants rare in California, but common elsewhere; List 3 = plants requiring additional data or abundance and distribution; and List 4 = plants of limited distribution.

^g CBS = creosote bush scrub; JTW = Joshua tree woodland; BS = blackbush scrub; SS = saltbush scrub; PJ = pinyon-juniper woodland.

^h Location of known populations in the study area shown in Figure 4-2, Map Volume.

TABLE 4-6
VEGETATION TYPES TRAVERSED BY THE ALTERNATIVE CORRIDORS^a

Link No.	Miles Traversed by Corridors												Total Miles		
	Vegetation Types ^b														
	CBS			BS										Barren	Cropland
	CBS	CBS/SS	CBS/JTW	SS	BS	BS/JTW	JTW	DW	RW						
0	6.0	7.4								0.1(1)			13.5		
1	15.1									1.5(1)			16.6		
2	5.5	1.0											6.5		
3	4.0	0.2		4.2									8.4		
4	1.0	0.9											1.9		
5a	0.2												0.2		
5b	3.5			0.5								0.8	4.8		
5c	5.5				5.8							1.0	12.3		
6	47.2		23.5	1.6	10.5					- (5)		0.5	83.3		
7		0.2	5.8	1.7				5.8		- (1)			13.5		
8	1.4	1.3	2.7	2.8				4.7		- (1)			12.9		
9	19.9		1.0	0.5				2.0		0.8(1)		0.8	25.0		
10	14.6		2.7					2.6					19.9		
11	13.2			2.8						2.5(2)		2.0	20.5		
12	3.0			1.0								0.9	4.9		
13	10.6												10.6		
14	1.9												4.9		
15	4.5												5.0		

Table 4-6 (continued)
Vegetation Types Traversed by the Alternative Corridors

Link No.	Miles Traversed by Corridors										Total Miles
	Vegetation Types ^b					JTW	DW	RW	Barren	Cropland	
	CBS		SS	BS							
	CBS	CBS/SS		CBS/JTW	BS						
16	5.4						3.2(1)				8.6
17	4.8						0.5(1)				5.3
18	3.7										3.7
19	8.0						-(1)				8.0
20a	14.6			4.2					2.2		21.0
20b	14.0		2.5	1.4			-(2)				17.9
21a	5.5			0.2					0.3		6.0
21b	6.4			1.6							8.0
22	19.7			1.0						0.5	21.2
23	11.1			0.2			1.5(4)			1.0	13.8
24a	24.1			2.2			1.0(2)				27.3
24b	1.3	2.3	1.8	0.7							6.1
25	11.7										11.7
26	0.6										0.6
27	12.2						1.0(1)	0.2			13.4
28	2.8			0.4							3.2

Table 4-6 (continued)
Vegetation Types Traversed by the Alternative Corridors

Link No.	Miles Traversed by Corridors										Total Miles	
	Vegetation Types ^b						JTW	DW	RW	Barren		Cropland
	CBS			SS	BS							
	CBS	CBS/SS	CBS/JTW		BS	BS/JTW						
29	0.6			1.5								2.1
30	1.0						1.1					2.1
31			0.8		2.3		2.6					5.7
32	14.5				1.2			0.4(1)	0.1		0.2	16.4
33	10.2				5.3				0.2			15.7
34					1.0			0.4(1)				1.4
35					4.7							4.7
36					4.2	0.6	1.1					5.9
37	0.9	2.0			3.1	0.8	0.5					7.3
38	3.4	2.1				4.2	0.6					10.3
39	3.2					2.5	0.5	-(4)				6.2
40a					3.9	2.3						6.2
40b					4.9							4.9
41	2.0			13.5								15.5
42	12.7			9.2								21.9
43	2.3			1.6								3.9

Table 4-6 (continued)
Vegetation Types Traversed by the Alternative Corridors

Link No.	Miles Traversed by Corridors								Total Miles				
	Vegetation Types ^b												
	CBS			BS									
	CBS	CBS/SS	CBS/JTW	SS	BS	BS/JTW	JTW	DW		RW	Barren	Cropland	
44													16.2
45													4.5
													0.5
													0.7
													16.2
													5.7

^aVegetation types along each link shown on Figure 4-2 (Map Volume).

^bCBS = creosote bush scrub; SS = saltbush scrub; BS = blackbush scrub; JTW = Joshua tree woodland; DW = desert wash (number in parenthesis indicates number of washes traversed); RW = riparian woodland.

TABLE 4.7
DESERT TORTOISE HABITAT TRAVERSED BY THE ALTERNATIVE CORRIDORS^a

Link No.	Estimated Tortoise Density Class Traversed by Mileposts ^b											
	Not Present			Low to Rare or Possibly Absent ^c			Low to Moderate			Moderate to High		
	Begin	End	Length	Begin	End	Length	Begin	End	Length	Begin	End	Length
0				-	-	13.5						
1	6.7	11.2	4.5	-	-	10.5	15.0	16.6	1.6			
2				-	-	2.0	0.0	4.5	4.5			
3	4.2	8.4	4.2	-	-	2.2	0.0	2.0	2.0			
4				-	-	1.9						
5a				-	-	0.2						
5b	3.5	4.8	1.3	-	-	3.5						
5c	0.0	1.0	1.0				1.0	6.5	5.5			
	6.5	12.3	5.8									
6	0.0	0.5	0.5	-	-	54.2	1.8	8.5	6.7			
	1.4	1.8	0.4				45.5	47.2	1.7			
	8.5	18.0	9.5									
	30.0	39.0	9.0									
	47.2	48.5	1.3									
7	0.0	4.0	4.0				4.0	13.5	9.5			
8	0.0	4.0	4.0				4.0	12.9	8.9			
9	0.0	6.5	6.5	-	-	5.0	11.5	25.0	13.5			
10	0.0	12.0	12.0				12.0	14.5	2.5	14.5	19.9	5.4
11	1.7	3.7	2.0	-	-	16.7	0.0	0.7	0.7			
	7.9	9.0	1.1									

Table 4-7 (continued)
Desert Tortoise Habitat Traversed by the Alternative Corridors

Link No.	Estimated Tortoise Density Class Traversed by Mileposts ^b											
	Not Present			Low to Rare or Possibly Absent ^c			Low to Moderate			Moderate to High		
	Begin	End	Length	Begin	End	Length	Begin	End	Length	Begin	End	Length
12	1.3	2.2	0.9	-	-	3.0	0.0	1.0	1.0	-	-	-
13	-	-	-	-	-	10.6	-	-	-	-	-	-
14	-	-	-	-	-	4.9	-	-	-	-	-	-
15	0.9	3.2	2.2	-	-	2.8	-	-	-	-	-	-
16	0.3	3.5	3.2	-	-	5.4	-	-	-	-	-	-
17	1.0	3.2	2.2	-	-	3.1	-	-	-	-	-	-
18	-	-	-	-	-	3.7	-	-	-	-	-	-
19	-	-	-	-	-	8.0	-	-	-	-	-	-
20a	8.8	11.0	2.2	-	-	19.8	-	-	-	-	-	-
20b	-	-	-	-	-	11.4	-	-	-	4.0	8.5	4.5
21a	4.2	6.0	1.8	-	-	4.2	-	-	-	11.0	13.0	2.0
21b	0.0	4.0	4.0	-	-	4.0	-	-	-	-	-	-
22	13.5	14.0	0.5	-	-	20.7	-	-	-	-	-	-
23	0.0	1.5	1.5	-	-	11.3	-	-	-	-	-	-
24a	17.8	21.0	3.2	-	-	-	0.0	8.0	8.0	8.0	16.0	17.8
24b	-	-	-	-	-	-	21.0	26.0	5.0	26.0	27.3	1.3
	-	-	-	-	-	-	16.0	15.0	1.0	0.0	6.1	6.1

Table 4-7 (continued)
Desert Tortoise Habitat Traversed by the Alternative Corridors

Link No.	Estimated Tortoise Density Class Traversed by Mileposts ^b											
	Not Present		Low to Rare or Possibly Absent ^c		Low to Moderate		Moderate to High		Very High			
	Begin	End	Length	Begin	End	Length	Begin	End	Length	Begin	End	Length
25												
26				-	-	0.6						
27	12.0	13.4	1.4	-	-	1.5						
28	0.0	0.6	0.6									
29												
30												
31												
32	14.5	15.2	0.7	-	-	8.1						
33	14.6	14.8	0.2	-	-	7.3						
34				-	-	1.4						
35				-	-	4.7						
36												
37												
38												
39												
40a												
40b												
41												

Table 4-7 (continued)
Desert Tortoise Habitat Traversed by the Alternative Corridors

Link No.	Estimated Tortoise Density Class Traversed by Mileposts ^b									
	Not Present		Low to Rare or Possibly Absent ^c		Low to Moderate		Moderate to High		Very High	
	Begin	End	Begin	End	Begin	End	Begin	End	Begin	End
42										
43										
44	0.0	16.2					0.0	21.9		21.9
45	4.5	5.7					0.0	3.9		3.9

^aDesert tortoise habitats traversed by the proposed corridors are shown on Figure 4-3; estimated density values shown on Table B-1.

^bEstimated density classes (individuals per square mile): (1) Low to rare or possibly absent = 0 to 50;

(2) Low to moderate = 50 to 150;

(3) Moderate to high = 100 to 250; and

(4) Very high = greater than 250.

^cMileposts for this density class are not provided for sake of brevity; note that this class is the most frequently traversed.

TABLE 4-8
MOHAVE GROUND SQUIRREL HABITAT
TRAVERSED BY THE PROPOSED CORRIDORS^a

Link Number	Mileposts		Length
	Begin	End	
20a	11.2	21.0	9.8
20b	0.0	17.9	17.9
24a	21.0	27.3	6.3
24b	0.0	6.1	6.1
28	0.6	3.2	2.6
29	0.0	2.1	2.1
30	0.0	2.1	2.1
31	0.0	5.7	5.7
32	0.0	14.5	14.5
	15.2	16.4	1.2
33	0.0	14.6	14.6
	14.8	15.7	0.9
34	0.0	1.4	1.4
35	0.0	4.7	4.7
36	0.0	5.9	5.9
37	0.0	7.3	7.3
38	0.0	10.3	10.3
39	0.0	6.2	6.2
40a	0.0	6.2	6.2
40b	0.0	4.9	4.9
41	0.0	15.5	15.5
42	0.0	21.9	21.9
43	0.0	3.9	3.9

^aMohave ground squirrel habitat traversed by the alternative corridors shown on Figure 4-3.

TABLE 4-9
POTENTIAL HABITAT OF SPECIAL INTEREST PLANT SPECIES TRAVERSED BY THE ALTERNATIVE CORRIDORS^a

Potential Habitat Traversed by Milepost

Link No.	Jager's Milkveitch			Mojave Spineflower			Desert Gymnosperms			Borlow Woolly Sunflower			Bicolored Penstemon			Neilson's Phacelia			Mojave Fishhook Cactus			Rusby's Desert Mallow		
	Begin	End	Length	Begin	End	Length	Begin	End	Length	Begin	End	Length	Begin	End	Length	Begin	End	Length	Begin	End	Length	Begin	End	Length
1																								
5c																								
6																						6.5	12.3	5.8
7																						8.5	14.0	5.5
8																						0.0	4.6	4.6
20a	18.5	21.0	2.5																			0.0	4.0	4.0
20b	0.0	11.0	11.0																					
21b																								
22																								
23																								
24a																								
24b																								
25																								
26																								
27																								
28																								
29																								
30																								
31																								
32																								
33																								
34																								
35																								
36																								
37																								
38																								
39																								
40a																								
40b																								
41																								
42																								
43																								
44																								

^aPotential habitat within proposed corridors for species listed above is shown on Figure 4-3; scientific names for species and regional distribution provided in Table 4-5.

TABLE 4-10
POTENTIAL AND KNOWN GOLDEN EAGLE NESTING HABITAT
NEAR THE ALTERNATIVE CORRIDORS^a

Link No.	Mileposts		Length (Miles)	Golden Eagle Nesting Habitat
	Begin	End		
0	8.0	9.9	1.9	3 miles northwest of potential nesting habitat on Eldorado Mountain
1	5.2	11.2	6.0	Adjacent to potential nesting habitat in the McCullough Mountains
5c	6.5	12.3	5.8	2 to 3 miles southeast of known nest sites on Clark Mountain
6	1.8	8.5	6.7	2 to 4 miles south of potential nesting habitat in the Clark Mountain Range
	8.5	18.0	9.5	Adjacent to known nesting habitat on Clark Mountain
	54.0	56.0	2.0	Adjacent to and north of potential nesting habitat in the Soda Mountains
	56.0	60.0	4.0	Adjacent to potential nesting habitat in the Avawatz Mountains
	64.0	83.3	19.3	Adjacent to potential nesting habitat in Alvord and Soda mountains
7	0.0	2.7	2.7	2 miles south of known nest sites on Clark Mountain
8	0.0	2.6	2.6	2 miles south of known nest sites on Clark Mountain
20a	0.0	4.8	4.8	3 miles south of potential nesting habitat on Alvord Mountain
	19.0	21.0	2.0	Adjacent to known nesting habitat in the Lane and Calico mountains
20b	0.0	2.0	2.0	Adjacent to known nesting habitat in the Lane and Calico mountains
21a	0.0	1.0	1.0	2 to 3 miles south of potential nesting habitat on Alvord Mountain

Table 4-10 (continued)

Potential and Known Golden Eagle Nesting Habitat

Link No.	Mileposts		Length (Miles)	Golden Eagle Nesting Habitat
	Begin	End		
21b	8.0	8.0	0.0	2 miles south of potential nesting habitat in the Calico Mountains
22	21.2	21.2	0.0	2 miles south of potential nesting habitat in the Calico Mountains
23	0.0	0.0	0.0	2 miles south of potential nesting habitat in the Calico Mountains
24	4.5	8.5	4.0	4 to 5 miles north of known nesting habitat in the Newberry Mountains
25	7.0	11.7	4.7	2 to 3 miles north of potential nesting habitat on Stoddard Ridge
32	9.8	14.5	4.7	0.7 mile south of potential nesting habitat on Quartzite Mountain
33	10.2	14.6	4.4	1 mile south of potential nesting habitat on Quartzite Mountain
44	12.5	16.2	3.7	Adjacent to known nesting habitat in the Lane and Calico mountains

^aKnown or potential golden eagle nesting habitat within or adjacent to the alternative corridors shown on Figure 4-3.

TABLE 4-11
DESERT BIGHORN SHEEP HABITAT TRAVERSED BY THE ALTERNATIVE CORRIDORS^a

Link No.	Mileposts		Length (Miles)	Description
	Begin	End		
1	6.7	11.2	4.5	Corridor traverses McCullough Mountains where a permanent bighorn sheep population is located mostly in south McCullough Mountains, south of the pass.
	16.0	16.5	0.5	Corridor 0.25 mile south of Sheep Mountains, a transient habitat for bighorn sheep which is probably utilized very infrequently.
2	1.5	3.0	1.5	Corridor 0.25 mile north of Roach Mountains, a transient habitat for bighorn sheep which is probably utilized very infrequently.
5c	10.0	11.0	1.0	Corridor 0.5 mile south of higher slopes of Clark Mountain which supports a permanent bighorn sheep population.
	11.0	11.9	0.9	Corridor traverses higher slopes of Clark Mountain (north of Mohawk Hill), which supports a permanent bighorn sheep population.
6	1.4	1.8	0.4	Corridor traverses lower slopes of Spring Mountains, a transient habitat for bighorn sheep which is probably utilized very infrequently.
	10.0	13.0	3.0	Corridor traverses higher slopes of Clark Mountains, which supports a permanent bighorn sheep population.
	53.0	60.0	7.0	Corridor passes through potential lowland movement corridor between Avawatz Mountains, 5 to 8 miles to the north which supports a permanent bighorn sheep population, and Soda Mountains, a transient sheep habitat which is probably utilized very infrequently.
7	0.0	1.0	1.0	Corridor traverses higher slopes of Clark Mountain (north of Mohawk Hill), which supports a permanent bighorn sheep population.

Table 4-11 (continued)
Desert Bighorn Sheep Habitat Traversed

Link No.	Mileposts		Length (Miles)	Description
	Begin	End		
7 (cont.)	1.0	2.7	1.7	Corridor 0.25 mile south of higher slopes of Clark Mountain (north of Mohawk Hill), which supports a permanent population of bighorn sheep.
8	0.0	0.6	0.6	Corridor traverses higher slopes of Clark Mountain (north of Mohawk Hill), which supports a permanent bighorn sheep population.
	0.6	2.6	2.0	Corridor 0.25 to 0.50 mile south of higher slopes of Clark Mountain (north of Mohawk Hill), which supports a permanent population of bighorn sheep.
11	8.0	9.0	1.0	Corridor traverses isolated hill within Soda Mountains, which is a transient habitat for bighorn sheep and is probably utilized infrequently.
15	0.9	3.2	2.3	Corridor traverses higher slopes of Cave Mountain (2 miles north of Afton Canyon), which probably provides habitat for permanent bighorn sheep population in Cady Mountains, 2 miles to the south.
17	1.0	3.2	2.2	Corridor traverses higher slopes of Cave Mountain (1 to 2 miles north of Afton Canyon), which probably provides habitat for permanent bighorn sheep population in Cady Mountains, 2 miles to the south.

^aBighorn sheep habitat traversed by, or near to, the alternative corridors shown on Figure 4-3.

TABLE 4-12
SENSITIVITY OF BIOLOGICAL RESOURCES POTENTIALLY AFFECTED
BY THE PROPOSED PROJECT

A. MAXIMUM SENSITIVITY

1. Riparian woodland vegetation and wildlife habitat.
2. Potential desert tortoise habitat (very high density: more than 250 individuals per square mile).
3. Three or more co-occurring high sensitivity resources (see below).

B. MAJOR SENSITIVITY

1. Desert wash vegetation and wildlife habitat.
2. Potential desert tortoise habitat (moderate to high density: 100 to 250 individuals per square mile).
3. Potential habitat for three or more special-interest plant species.
4. Known nesting habitat for the golden eagle and/or prairie falcon.
5. Known permanent range of desert bighorn sheep.
6. Three or more co-occurring moderate sensitivity resources (see below).

C. MODERATE SENSITIVITY

1. Joshua tree woodland vegetation and wildlife habitat.
2. Within or adjacent to potential Gila monster habitat.
3. Potential desert tortoise habitat (low to moderate density: 50 to 150 individuals per square mile).
4. Potential likely nesting habitat for the Swainson's hawk, yellow-billed cuckoo, and least bell's vireo.^a

^aPotential habitat for these riparian-nesting species occurs along the Mojave River, although no specific location can be identified. The sensitivity of these resources is accounted for in riparian woodlands (maximum sensitivity category) during the impact assessment process.

Table 4-12 (continued)
Sensitivity of Biological Resources

C. MODERATE SENSITIVITY (continued)

5. Potential Mohave ground squirrel habitat.
6. Potential habitat for one or two special-interest plant species.
7. Within potential, or adjacent to known, nesting habitat for the golden eagle and/or prairie falcon.
8. Adjacent to permanent range of desert bighorn sheep.
9. Four or more co-occurring low sensitivity resources.

D. MINIMAL SENSITIVITY

1. Scrub vegetation and wildlife habitat (creosote bush, saltbush, and blackbush scrub), as well as barren areas, croplands, and built-up areas.
2. Potential desert tortoise habitat (low density: 0 to 50 individuals per square mile).
3. Adjacent to potential nesting habitat for the golden eagle and prairie falcon.
4. Within or adjacent to transient desert bighorn sheep habitat or movement corridor.

TABLE 4-13
IMPACT LEVEL MODEL^a

<u>Access Level</u>	<u>Resource Sensitivity</u>			
	<u>Minimal</u>	<u>Moderate</u>	<u>Major</u>	<u>Maximum</u>
Sufficient	L	L	M	M/H
Upgraded	L	M	M/H	H
New	M	M/H	H	H+

^aImpact levels represent a composite rating of impact magnitude and significance based on access requirements and resource sensitivity where L = low, M = moderate, and H = high. Access requirements and resource sensitivity levels are described in text. Generic mitigation measures have been taken into account in the assignment of impact levels in the model.

TABLE 4-14
MATRIX OF RECOMMENDED SELECTIVE MITIGATION MEASURES AND AFFECTED RESOURCES^a

Resource Sensitivity ^b	No Required Mitigation	Recommended Selective Mitigation Measures ^c					
		1	2	4	8	11	13
A. MAXIMUM SENSITIVITY							
1. Riparian woodland/habitat		X	X		X		X
2. Desert tortoise habitat (H density)		X	X	X		X	
B. MAJOR SENSITIVITY							
1. Desert wash vegetation/habitat		X					
2. Desert tortoise habitat (M-H density)		X	X		X		X
3. 3 or more special-interest plants		X				X	X
4. Raptor nesting habitat	X			X			
5. Bighorn sheep permanent range		X				X	
C. MODERATE SENSITIVITY							
1. Joshua tree woodland/habitat		X					
2. Gila monster habitat		X					
3. Desert tortoise habitat (L-M density)		X					
4. Riparian bird nesting habitat		X	X		X		
5. Mohave ground squirrel habitat		X					
6. 1 or 2 special-interest plant(s)		X					X
7. Near raptor nesting habitat	X						
8. Near bighorn sheep permanent range	X						

^aThis matrix indicates which recommended selective mitigation measures are applicable to affected resources and would result in reduced impact levels.

^bOnly moderate to maximum sensitivity resources potentially affected by the proposed project are listed.

^cThese measures are described in the text.

TABLE 4-15
CHANGES IN IMPACT LEVEL DUE TO
APPLICATION OF RECOMMENDED SELECTIVE MITIGATION MEASURES

<u>Initial Impact Level</u>	<u>Application of Mitigation Measures</u>	<u>Residual Impact Level</u>
Low (0.5, 1.0)	No mitigation applied if all resources affected have low sensitivities	Low (0.5, 1.0)
Low (0.5, 1.0)	1 or 2 mitigation measures applied when one or more of the affected resources has a moderate sensitivity	Negligible/Low (0.0, 0.1)
Moderate (7)	1 or 2 mitigation measures applied	Low (0.5, 1.0)
Moderate (7)	3 or 4 mitigation measures applied	Negligible/Low (0.0, 0.1)
Moderate (9)	1 mitigation measure applied	Moderate (7)
Moderate (9)	2 to 3 mitigation measures applied	Low (0.1, 0.5, 1.0)
High (13)	1 or 2 mitigation measures applied	Moderate (7, 9)
High (13)	3 mitigation measures applied	Low (1.0)
High (15)	4 mitigation measures applied	Moderate (7)

This represents only the combinations of mitigation measures applied in IA/MP charts. Numeric ratings for the impact levels are shown in parentheses.

TABLE 4-16
IMPACT ASSESSMENT/MITIGATION PLANNING CHART
BIOLOGICAL RESOURCES

Link Number	Length	Mileposts Begin	Mileposts End	Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Mitig- ation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
0	13.5	0.0	3.0	1,2,4	S	L (.5)	-	-	-	L (.5)	-	-
		3.0	3.1	1,2,4	S	M (7)	-	Desert wash; potential GM habitat	2,8	L (.5)	-	-
		3.1	13.5	1,2,4	S	L (.5)	-	-	-	L (.5)	-	-
1	16.6	0.0	5.2	1,2,4	S	L (.5)	-	-	-	L (.5)	-	-
		5.2	6.7	1-4	S	M (7)	-	Potential GM habitat, potential habitat for 2 special-interest plants (McCullough Mountains)	2,13a	L (.5)	-	-
		6.7	11.2	1-4	S	M (7)	-	Bighorn sheep habitat; potential habitat for 2 special-interest plants (McCullough Mountains)	2,11a, 13a	L (.1)	-	-
		11.2	16.6	1,2,4	S	L (.5)	-	Potential DT habitat (low to moderate density)	2	L (.1)	-	-
2	6.5	0.0	2.0	1,2,4	S	L (.5)	-	Potential DT habitat (low to moderate density)	2	L (.1)	-	-
		2.0	6.5	1,2,4	S	L (.5)	-	-	-	L (.5)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
3	8.4	0.0	2.0	1,2,4	S	L (.5)	-	Potential DT habitat (low to moderate density)	2	L (.1)	-	-
4	1.9	0.0	1.9	1,2,4	S	L (.5)	-	-	-	L (.5)	-	-
				1,2,4	S	L (.5)	-	-	-	L (.5)	-	-
				1,2,4	S	L (.5)	-	-	-	L (.5)	-	-
5a	0.2	0.0	0.2	1,2,4	S	L (.5)	-	-	-	L (.5)	-	-
5b	4.8	0.0	3.5	1,2,4	S	L (.5)	-	-	-	L (.5)	-	-
5c	12.3	0.0	1.0	1,2	S	L (.5)	-	-	-	L (.5)	-	-
				-	S	L (.5)	-	-	-	L (.5)	-	-
				-	S	L (.5)	-	-	-	L (.5)	-	-
5c	12.3	1.0	6.5	1,2,4	S	L (.5)	-	Potential DT habitat (low to moderate density)	2	L (.1)	-	-
				-	S	L (.5)	-	-	-	L (.5)	-	-
5c	12.3	6.5	7.3	1-4	S	M (7)	-	Potential GM habitat; potential habitat for 2 special-interest plants; adjacent to known GE/PF nesting habitat (Clark Mountain)	2,13a	L (.5)	-	-
				-	S	M (7)	-	-	-	L (.5)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
5c (cont.)		7.3	9.0	1-4	N	H (15)	Yes	Potential GM habitat; potential habitat for 2 special-interest plants; adjacent to known GE/ PF nesting habitat (Clark Mountain)	8,13a	M (7)	-	-
		9.0	12.3	1-4	S	M (7)	-	Potential GM habitat; potential habitat for 2 special-interest plants; adjacent to known GE/ PF nesting habitat; within and adjacent to BS habitat	2,11a, 13a,8	N (0)	-	-
6	83.3	0.0	0.5	1,2	S	L (.5)	-	-	-	L (.5)	-	-
		0.5	1.8	1,2,4	S	L (.5)	-	-	-	L (.5)	-	-
		1.8	8.5	1,2,4	S	L (.5)	-	Potential DT habitat (low to moderate density); adjacent to known GE/PF nesting habitat (Clark Mountain Range)	2	L (.1)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Mitig- ation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
6 (cont.)		8.5	10.0	1-4	S	L (.5)	-	Potential habitat for 2 special-interest plants; adjacent to known GE/ PF nesting habitat	2,13a	N (0)	-	-
		10.0	13.0	1-4	S	M (7)	-	Potential habitat for 2 special-interest plants; within BS habitat; adja- cent to known GE/PF nesting habitat (Clark Mountain)	2,11a, 13a	L (.1)	-	-
		13.0	14.0	1-4	S	L (.5)	-	Potential habitat for 2 special-interest plants; adjacent to known GE/ PF nesting habitat (Clark Mountain)	2,13a	N (0)	-	-
		14.0	30.0	1,2,4	S	L (.5)	-	-	-	L (.5)	-	-
		30.0	31.0	1,2	S	L (.5)	-	-	-	L (.5)	-	-
		31.0	38.0	1,2	S	L (.5)	-	Joshua tree woodland	2	L (.1)	-	-
		38.0	45.5	1,2,4	S	L (.5)	-	-	-	L (.5)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Mitig- ation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
6 (cont.)		45.5	47.2	1,2,4	S	L (.5)	-	Potential DT habitat (low to moderate density)	2	L (.1)	-	-
		47.2	48.5	1,2	S	L (.5)	-	-	-	L (.5)	-	-
		48.5	53.0	1,2,4	S	L (.5)	-	-	-	L (.5)	-	-
		53.0	60.0	1,2,4	S	L (.5)	-	Potential DT habitat (low density); adjacent to potential GE/PF nesting habitat; poten- tial BS movement corridor	2	L (.1)	-	-
		60.0	64.0	1,2,4	S	L (.5)	-	-	-	L (.5)	-	-
		64.0	82.0	1,2,4	S	L (.5)	-	Potential DT habitat (low density); adja- cent to potential GE/ PF nesting habitat; adjacent to transient BS habitat	2	L (.1)	-	-
		82.0	83.3	1,2,4	S	L (.5)	-	-	-	L (.5)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
7	13.5	0.0	2.7	1-4	N	H (13)	Yes	Potential habitat for 2 special-interest plants; within and adjacent to BS habitat; adjacent to known GE/PF nesting habitat (Clark Mountain)	11a, 13a	M (7)	-	-
								Potential habitat for 2 special-interest plants; potential DT habitat (low to moderate density)	13a	M (7)	-	-
								Potential DT habitat (low to moderate density)	11b, 13b	M (7)	-	Mitigation Measure 11b not committed to
8	12.9	0.0	2.6	1-4	S	M (7)	-	Potential habitat for 2 special-interest plants; within and adjacent to BS habitat; adjacent to GE/PF nesting habitat (Clark Mountain)	2,4, 11a, 13a	N (0)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Mitig- ation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
8 (cont.)		2.6	5.9	1-4	S	L (.5)	-	Potential habitat for 2 special-interest plants; potential DT habitat (low to moderate density)	2,13a	N (0)	-	-
9	25.0	5.9	12.9	1,2,4	S	L (.5)	-	Potential DT habitat (low to moderate density)	2	L (.1)	-	-
		0.0	2.0	1,2	U	L (.5)	-	-	-	L (.5)	-	-
		2.0	10.6	1,2,4	N	M (7)	-	-	-	M (7)	-	-
		10.6	11.5	1,2,4	S	L (.5)	-	-	-	L (.5)	-	-
		11.5	24.0	1,2,4	S	L (.5)	-	Potential DT habitat (low to moderate density)	2	L (.1)	-	-
		24.0	25.0	1,2,4	S	M (7)	-	Desert wash; potential DT habitat (low to moderate density)	2,8	L (.5)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
10	19.9	0.0	9.0	1,2	N	M (7)	-	-	-	M (7)	-	-
		9.0	12.0	1,2	N	M (7)	-	-	-	M (7)	-	-
		12.0	14.5	1,2,4	U	M (9)	-	Potential DT habitat (low to moderate density)	11b, 13b	M (7)	-	Mitigation Measure 11b not com- mitted to
11	20.5	14.5	19.9	1,2,4	U	M (9)	-	Potential DT habitat (moderate to high density)	11b, 13b	L (1)	-	-
		0.0	0.7	1,2,4	N	M (9)	-	Potential DT habitat (low to moderate density)	11b, 13b	M (7)	-	Mitigation Measure 11b not com- mitted to
		0.7	1.7	1,2,4	N	M (7)	-	-	-	M (7)	-	-
		1.7	3.7	1,2	N	M (7)	-	-	-	M (7)	-	-
		3.7	18.0	1,2,4	N	M (7)	-	-	-	M (7)	-	-
		18.0	20.5	1,2,4	N	H (13)	Yes	Desert wash (Mojave River Wash/Cronese Valley)	8	M (9)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
12	4.9	0.0	1.0	1,2,4	N	M (9)	-	Potential DT habitat (low to moderate density)	11b, 13b	M (7)	-	Mitigation Measure 11b not com- mitted to
		1.0	1.3	1,2,4	N	M (7)	-		-	M (7)	-	
		1.3	2.2	-	N	M (7)	-		-	M (7)	-	
		2.2	3.8	1,2,4	N	M (7)	-		-	M (7)	-	
		3.8	4.9	1,2,4	N	H (13)	Yes	Desert wash	1,8	M (7)	-	
13	10.6	0.0	1.8	1,2,4	S	L (.5)	-		-	L (.5)	-	
		1.8	3.5	1,2,4	S	M (7)	-	Desert wash	8	L (1)	-	
		3.5	7.0	1,2,4	S	L (.5)	-		-	L (.5)	-	
		7.0	10.6	1,2,4	U	L (.5)	-		-	L (.5)	-	
14	4.9	0.0	1.9	1,2,4	N	M (7)	-		-	M (7)	-	
		1.9	4.9	1,2,4	N	H (13)	Yes	Desert wash (Mojave River Wash)	8	M (9)	-	

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
15	5.0	0.0	0.5	1-4	N	H (13)	Yes	Desert wash (Mojave River Wash)	8	M (9)	-	-
		0.5	0.9	1,2,4	N	M (7)	-	-	-	M (7)	-	-
		0.9	3.2	1,2,4	N	H (13)	Yes	Known BS habitat (Cady Mountains)	1,11a	M (9)	-	Mitigation Measure 11b not com- mitted to
16	8.6	3.2	5.0	1,2,4	N	M (7)	-	-	-	M (7)	-	-
		0.0	0.3	1,2,4	U	M (7)	-	-	-	M (7)	-	-
		0.3	3.5	1,2	U	H (13)	Yes	Desert wash (Cronese Valley)	4,8	M (7)	-	-
17	5.3	3.5	8.6	1,2,4	N	M (7)	-	-	-	M (7)	-	-
		0.0	0.5	1-4	N	H (13)	Yes	Desert wash (Mojave River Wash)	8	M (9)	-	-
		0.5	1.0	1,2,4	N	M (7)	-	-	-	M (7)	-	-
		1.0	3.2	1,2,4	N	H (13)	Yes	Known BS habitat (Cady Mountains)	1,11a	M (9)	-	Mitigation Measure 11b not com- mitted to

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
17 (cont.)		3.2	5.3	1,2,4	N	M (7)	-	-	-	M (7)	-	-
18	3.7	0.0	3.7	1,2,4	U	L (.5)	-	-	-	L (.5)	-	-
19	7.9	0.0	7.9	1,2,4	N	M (7)	-	-	-	M (7)	-	-
20a	21.0	0.0	8.8	1,2,4	N	M (7)	-	-	-	M (7)	-	-
		8.8	11.0	-	N	M (7)	-	-	-	M (7)	-	-
		11.2	14.4	1,2,4	N	M (9)	-	Potential MGS habitat	-	M (9)	-	-
		14.4	18.5	1,2,4	U	M (7)	-	Potential MGS habitat	-	M (7)	-	-
		18.5	21.0	1-4	U	M (9)	-	Potential habitat for 2 special-interest plants; potential MGS habitat; adjacent to known GE/ PF nesting habitat	13a	M (7)	-	-
20b	17.9	0.0	2.7	1-4	U	M (9)	-	Potential habitat for 2 special-status plants; potential MGS habitat; adjacent to known GE/ PF nesting habitat	13a	M (7)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
20b (cont.)		2.7	4.0	I-4	N	H (13)	Yes	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (low to moderate density)	11b, 13a	M (9)	-	Mitigation Measure 11b not com- mitted to
		4.0	7.8	I-4	N	H (13)	Yes	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	11b, 13a, 13b	L (1)	-	-
		7.8	8.5	I-4	U	M (9)	-	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	11b, 13a, 13b	L (.5)	-	-
		8.5	11.0	I-4	S	L (.5)	-	Potential habitat for 2 special-interest plants; potential MGS habitat	2,13a	N (1)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
20b (cont.)		11.0	13.0	1-4	S	M (7)	-	Potential habitat for a special-interest plant; potential MGS habitat; potential DT habitat (moderate to high density)	2,11b, 13a, 13b	N (1)	-	-
21a	6.0	0.0	4.2	1,2,4	S	L (.5)	-	Potential habitat for a special-interest plant species; potential MGS habitat	2,13a	N (1)	-	-
21b	8.0	0.0	4.0	-	S	L (.5)	-		-	L (.5)	-	-
		4.2	6.0	-	S	L (.5)	-		-	L (.5)	-	-
		0.0	4.0	-	S	L (.5)	-		-	L (.5)	-	-
		4.0	8.0	1-4	S	L (.5)	-	Potential habitat for a special-interest plant	2,13a	N (1)	-	-
22	21.1	0.0	13.5	1,2,4	U	L (.5)	-		-	L (.5)	-	-
		13.5	14.0	-	U	L (.5)	-		-	L (.5)	-	-
		14.0	16.0	1,2,4	U	L (.5)	-		-	L (.5)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
22 (cont.)		16.0	21.1	1-4	U	M (7)	-	Potential habitat for a special-interest plant	13a	L (1)	-	-
23	13.8	0.0	1.5	1,2,4	S	M (7)	-	Desert wash (Mojave River)	2,8	L (.5)	-	-
		1.5	3.5	1-4	S	L (.5)	-	Potential habitat for 2 special-interest plants	2,13a	N (0)	-	-
		3.5	4.5	1,2	S	L (.5)	-	-	-	L (.5)	-	-
		4.5	13.8	1-4	S	L (.5)	-	Potential habitat for 2 special-interest plants	2,13a	L (.1)	-	-
24a	27.3	0.0	8.0	1-4	N	M (9)	-	Potential habitat for a special-interest plant; potential DT habitat (low to moderate density)	13a	M (7)	-	-
		8.0	16.0	1-4	N	H (13)	Yes	Potential habitat for 2 special-interest plants; potential DT habitat (moderate to high density)	11b, 13a, 13b	L (1)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts Begin	Mileposts End	Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Mitig- ation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
24a (cont.)		16.0	17.8	1-4	N	H (15)	Yes	Potential habitat for 2 special-interest plants; potential DT habitat (very high density)	1, 11b, 13a, 13b	M (7)	-	-
		17.8	18.8	1, 2	N	M (7)	-	-	-	M (7)	-	-
		18.8	19.8	1, 2	N	H (13)	Yes	Desert wash (Mojave River)	1, 8	M (7)	-	-
		19.8	21.0	1, 2	N	M (7)	-	-	-	M (7)	-	-
		21.0	26.0	1-4	N	H (13)	Yes	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (low to moderate density)	11b, 13a	M (9)	-	Mitigation Measure 11b not com- mitted to
		26.0	27.3	1-4	N	H (13)	Yes	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	11a, 13a, 13b	L (1)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts Begin	Mileposts End	Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Mitiga- tion	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
24b	6.1	0.0	6.1	1-4	N	H (13)	Yes	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	11b, 13a, 13b	L (1)	-	-
25	11.7	0.0	7.0	1-4	S	L (.5)	-	Potential habitat for a special-interest plant; potential DT habitat (low to moderate density)	2,13a	N (0)	-	-
26	0.6	0.0	0.6	1-4	N	M (9)	-	Potential DT habitat (moderate to high density)	2,11b, 13b	L (.1)	-	-
27	13.4	0.0	6.5	1-4	N	H (15)	Yes	Potential habitat for a special-interest plant	13a	M (7)	-	-
								Potential habitat for 2 special-interest plants; potential DT habitat (very high density)	1,11b, 13a, 13b	M (9)	-	Mitigation Measure 11b not com- mitted to
		6.5	8.0	1-4	N	M (9)	-	Potential habitat for 2 special-interest plants	13a	M (7)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
27 (cont.)		8.0	10.0	1-4	N	H (13)	Yes	Potential habitat for 2 special-interest plants; potential DT habitat; (moderate to high density)	11b, 13a, 13b	L (1)	-	-
		10.0	12.0	1-4	U	M (9)	-	Potential habitat for 2 special-interest plants; potential DT habitat; (moderate to high density)	11b, 13a, 13b	L (.5)	-	-
28		12.0	13.0	1,2	U	M (9)	-	Desert wash (Mojave River)	4,8,	L (1)	-	-
		13.0	13.4	1,2	U	H (13)	Yes	Riparian woodland (Mojave River)	4,8	M (7)	-	-
	3.2	0.0	0.6	1,2	N	M (7)	-	-	-	M (7)	-	-
	0.6	3.2	3.2	1-4	N	H (13)	Yes	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	11b, 13a, 13b	L (1)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miri- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
29	2.1	0.0	2.1	1-4	N	H (13)	Yes	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	11b, 13a, 13b	L (1)	-	-
30	2.1	0.0	2.1	1-4	S	M (7)	-	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density); Joshua tree woodland (Milepost 1.0-2.1)	2,11b, 13a, 13b	N (0)	-	-
31	5.7	0.0	5.7	1-4	S	M (7)	-	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (low to moderate density); Joshua tree woodland (Mileposts 0.0-1.2, 2.0-3.4)	2,13a	L (.5)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
32	16.4	0.0	7.6	1-4	S	M (7)	-	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	2,11a, 13a, 13b	N (0)	-	-
		7.6	14.5	1-4	S	L (.5)	-	Potential habitat for 2 special-interest plants; potential MGS habitat	2,13a	N (0)	-	-
		14.5	14.6	1,2	S	M (9)	-	Riparian woodland (Mojave River)	2,8	L (1)	-	-
		14.6	15.0	1,2	S	M (7)	-	Desert wash (Mojave River)	2,8	L (.5)	-	-
		15.0	15.2	1,2	S	L (.5)	-	-	-	L (.5)	-	-
		15.2	16.4	1-4	S	L (.5)	-	Potential habitat for 2 special-interest plants; potential MGS habitat	2,13a	N (0)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

<u>Link Number</u>	<u>Length</u>	<u>Mileposts</u>		<u>Impact Types^a</u>	<u>Access Level</u>	<u>Initial Impact Level^c</u>	<u>Sig- nifi- cant?</u>	<u>Resource Affected^d</u>	<u>Recom- mended Miti- gation</u>	<u>Residual Impact Level</u>	<u>Sig- nifi- cant?</u>	<u>Unavoidable Signifi- cant Adverse Impact</u>
33	15.5	0.0	6.5	1-4	S	M (7)	-	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	2, 11b, 13a, 13b	N (0)	-	-
		6.5	8.2	1-4	N	H (13)	Yes	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	11b, 13a, 13b	L (1)	-	-
		8.2	11.0	1-4	N	M (9)	-	Potential habitat for 2 special-interest plants; potential MGS habitat	13a	M (7)	-	-
		11.2	14.6	1-4	S	L (.5)	-	Potential habitat for 2 special-interest plants; potential MGS habitat	2, 13a	N (0)	-	-
		14.6	14.8	1,2	S	M (9)	-	Riparian woodland (Mojave River)	2,4,8	L (.5)	-	-
		14.8	15.5	1-4	S	L (.5)	-	Potential habitat for 2 special-interest plants; potential MGS habitat	2, 13a	N (0)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
34	1.4	0.0	1.0	1-4	S	L (.5)	-	Potential habitat for 2 special-interest plants; potential MGS habitat	2,13a	N (0)	-	-
		1.0	1.4	1,2,4	S	M (7)	-	Desert wash; potential MGS habitat	2,8	L (.5)	-	-
35	4.7	0.0	4.7	1-4	S	L (.5)	-	Potential habitat for 2 special-interest plants; potential MGS habitat	2,13a	N (0)	-	-
36	5.8	0.0	2.5	1-4	N	H (13)	Yes	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	11b, 13a, 13b	L (1)	-	-
		2.5	4.3	1-4	U	M (9)	-	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	2,11b, 13a, 13b	L (.1)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts Begin	Mileposts End	Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
36 (cont.)		4.3	5.8	1-4	N	H (13)	Yes	Joshua tree woodland; potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	11b, 13a, 13b	L (1)	-	-
37	7.3	0.0	5.0	1-4	N	H (13)	Yes	Potential habitat for 3 special-interest plants; potential MGS habitat; potential DT habitat (low to moderate density)	11b, 13a, 13b	M (7)	-	Mitigation Measure 11b not com- mitted to
		5.0	7.3	1-4	N	H (13)	Yes	Potential habitat for 3 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density); Joshua tree woodland (Milepost 6.8-7.3)	11b, 13a, 13b	L (1)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
38	10.3	0.0	4.6	1-4	S	M (7)	-	Potential habitat for 3 special-interest plants; potential MGS habitat; potential DT habitat (low to moderate density)	2,13a	L (.5)	-	-
		4.6	5.2	1-4	N	H (13)	-	Potential habitat for 3 special-interest plants; potential MGS habitat; potential DT habitat (low to moderate density)	2,13a	M (7)	-	-
		5.2	10.3	1-4	S	M (7)	-	Potential habitat for 3 special-interest plants; potential MGS habitat; potential DT habitat (low to moderate density); Joshua tree woodland (Milepost 9.7-10.3)	2,13a	L (.5)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
39	6.2	0.0	6.2	1-4	S	M (7)	-	Potential habitat for 3 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density); Joshua tree woodland (Milepost 0.0-0.5)	2,11a, 13a, 13b	N (0)	-	-
40a	6.2	0.0	6.2	1-4	S	M (7)	-	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	2,11b, 13a, 13b	N (0)	-	-
40b	4.9	0.0	4.9	1-4	S	M (7)	-	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	2,11b, 13a, 13b	N (0)	-	-

Table 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Miti- gation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
41	15.5	0.0	15.0	I-4	S	M (7)	-	Potential habitat for 3 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	2, 11b, 13a, 13b	N (0)	-	-
42	21.9	0.0	2.4	I-4	N	H (13)	Yes	Potential habitat for 3 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	11b, 13a, 13b	L (1)	-	-
		2.4	21.9	I-4	U	M (9)	-	Potential habitat for 3 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	11b, 13a, 13b	L (.5)	-	-

Tabel 4-16 (continued)
Impact Assessment/Mitigation Planning Chart
Biological Resources

Link Number	Length	Mileposts		Impact Types ^a	Access Level	Initial Impact Level ^c	Sig- nifi- cant?	Resource Affected ^d	Recom- mended Mitig- ation	Residual Impact Level	Sig- nifi- cant?	Unavoidable Significant Adverse Impact
		Begin	End									
43	3.9	0.0	3.9	1,4	N	H (13)	Yes	Potential habitat for 2 special-interest plants; potential MGS habitat; potential DT habitat (moderate to high density)	11b, 13a, 13b	L (1)	-	-
44	16.2	0.0	3.4	1,2,4	U	L (.5)	-	-	-	L (.5)	-	-
		3.4	12.5	1,2	N	M (7)	-	-	-	M (7)	-	-
		12.5	16.2	1,2,4	N	M (9)	-	Adjacent to known GE/PF nesting habitat; potential habitat for 2 special-interest plants	13a	M (7)	-	-
45	5.7	0.0	4.5	1,2,4	N	M (7)	-	-	-	M (7)	-	-
		4.5	5.7	1,2	N	M (7)	-	-	-	M (7)	-	-

^aImpact Types: 1 = effects on vegetation; 2 = effects on wildlife; 3 = effects on legally protected species (animals only); and 4 = effects on species of special-interest (plants and animals).

^bAccess level categories: S = existing access roads are sufficient; U = existing access roads require upgrading; and N = new access required.

^cInitial impact level: a 4 level rating of impact magnitude and significance based on impact intensity and resource sensitivity. N = negligible; L = low; M = moderate; and H = high.

^dOnly resources of moderate, high, or very high sensitivity are listed. DT = desert tortoise; BS = bighorn sheep; MGS = Mohave ground squirrel; GM = Gila monster; GE = golden eagle; PF = prairie falcon.

PLANT SPECIES OF POTENTIAL SPECIAL INTEREST
REVIEWED AND REMOVED FROM FURTHER CONSIDERATION

TABLE A-1
PLANT SPECIES OF POTENTIAL SPECIAL INTEREST REVIEWED AND REMOVED FROM FURTHER CONSIDERATION^a

Scientific Name	Occurrence	Federal Candidate Status ^b		CNPS Lists ^c		NNPS List ^d	Reason for No Further Consideration
		1980	1983	1980	1984		
<u>Agave utahensis</u> var. <u>eborspina</u>	NV, CA	2	2	4	2	W	Not in study area in California; not expected near corridors in Nevada
<u>A. utahensis</u> var. <u>nevadensis</u>	NV	2	3C	-	-	W	Dropped from Federal candidate list
<u>Arctomecon merriamii</u>	CA, NV	2	3C	4	2	W	Dropped from Federal candidate list
<u>Astragalus cimae</u> var. <u>cimae</u>	CA, NV	2	3C	2	IB	OR	Dropped from Federal candidate list
<u>A. convallarius</u> var. <u>finitimus</u>	NV	2	2	-	-	W	Not in study area in Nevada
<u>A. funereus</u>	CA, NV	1	2	2	IB	W	Not in study area in California or Nevada
<u>A. geyeri</u> var. <u>triquetrus</u>	NV	1	1	-	-	FCT	Not in study area in Nevada
<u>A. lentiginous</u> var. <u>sesquimetralis</u>	CA, NV	1	1	2	IB	FCE	Not in study area in California or Nevada
<u>A. lentiginous</u> var. <u>latus</u>	NV	2	3C	-	-	-	Dropped from Federal candidate list
<u>A. mohavensis</u> var. <u>hemigyrrus</u>	CA, NV	1	2	2	AI	FCE	Not in study area in California or Nevada
<u>A. oophrus</u> var. <u>lonchocalyx</u>	NV	2	3C	0	-	W	Dropped from Federal candidate list
<u>Brickellia knappiana</u>	CA, NV	2	3C	2	3	W	Dropped from Federal candidate list

Table A-1 (continued)
Plant Species of Potential Special Interest

Scientific Name	Occurrence	Federal Candidate Status ^b		CNPS List ^c		NNPS List ^d	Reason for No Further Consideration
		1980	1983	1980	1984		
<u>Calochortus striatus</u>	CA, NV	2	3C	2	IB	W	Dropped from Federal candidate list
<u>Coryphantha vivipara</u> var. <u>rosea</u>	CA, NV	2	3C	2	IB	W	Dropped from Federal candidate list
<u>Cryptantha insolita</u>	NV	1	2	-	-	FCE	Not in study area in Nevada; probably extinct
<u>C. tumulosa</u>	NV	1	3C	-	-	W	Dropped from Federal candidate list
<u>Dalea arborescens</u>	CA	3C	3C	AI	4	-	Dropped from Federal candidate list
<u>Enceliopsis nudicalis</u> var. <u>corrugata</u>	NV	1	1	-	-	FCE	Not in study area in Nevada
<u>Ephedra funera</u>	CA, NV	3C	3C	AI	AI	-	Dropped from Federal candidate list
<u>Epilobium nevadensis</u>	NV	1	2	-	-	W	Not expected near corridors in Nevada
<u>Forsellesia pugens</u> var. <u>corrugata</u>	CA, NV	2	3C	2	IB	W	Dropped from Federal candidate list
<u>Galium hilendiae</u> ssp. <u>kingstonensis</u>	CA, NV	1	2	2	IB	FCT	Not in study area in California or Nevada
<u>Gilia ripleyi</u>	NV	3C	3C	-	-	OR	Dropped from Federal candidate list
<u>Haplopappus brickelloides</u>	CA, NV	3C	3C	AI	-	OR	Dropped from Federal candidate list

Table A-1 (continued)
Plant Species of Potential Special Interest

Scientific Name	Occurrence	Federal Candidate Status ^b		CNPS List ^c		NNPS List ^d	Reason for No Further Consideration
		1980	1983	1980	1984		
<u>Hemizonia mojavenis</u>	CA	1	1	1	1A	-	Not expected near corridors in California
<u>Ivesia cryptocaulis</u>	NV	1	2	-	-	W	Not expected near corridors in Nevada
<u>Linanthus arenicola</u>	CA, NV	3C	3C	4	2	-	Dropped from Federal candidate list
<u>Mentzelia leucophylla</u>	NV	1	1	-	-	FCE	Not in study area in Nevada
<u>Opuntia basilaris</u> var. <u>brachyclada</u>	CA	1	2	2	1B	-	Not expected near corridors in California
<u>O. phaeacantha</u> var. <u>mojavensis</u>	CA	2	3B	2	1A	-	Dropped from Federal candidate list
<u>Penstemon fruticiformis</u> ssp. <u>amargosae</u>	CA, NV	1	2	2	4	FCT	Not in study area in California or Nevada
<u>P. stephensii</u>	CA	2	2	2	1B	-	Not in study area in California
<u>Perityle megalcephala</u>	CA, NV	2	3C	1A	1A	OR	Dropped from Federal candidate list
<u>Pholisma arenarium</u>	CA, NV	3C	3C	4	1A	-	Dropped from Federal candidate list

Table A-1 (continued)
Plant Species of Potential Special Interest

Scientific Name	Occurrence	Federal Candidate Status ^b		CNPS Lists ^c		NNPS List ^d	Reason for No Further Consideration
		1980	1983	1980	1984		
<u>Potentilla patellifera</u>	CA	2	2	2	IB	-	Not in study area in California
<u>Silene clokeyi</u>	NV	1	2	-	-	W	Not in study area in Nevada

^aThese species were not considered further in the site-specific analysis to this project because: (1) they have been removed from the Federal candidate list recently; (2) they are not known to occur in the study area; or (3) they are not expected to occur near the alternative corridors.

^bBased on lists from USFWS (1980, 1983). See Table 4-5 for explanation of category designations.

^cCalifornia Native Plant Society (CNPS) 1980 and 1984 lists. See Table 4-5 for references and explanation of lists.

^dNorthern Nevada Native Plant Society (NNPS) 1984 lists of rare plants. See Table 4-5 for references and explanation of lists.

TABLE B-1
INVENTORY OF BIOLOGICAL RESOURCES WITHIN AND ADJACENT TO CORRIDORS^a

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
0	0.0	3.0	Broad alluvial fan, sandy soils	CBS; large creosote bushes with few other shrubs; other species: <u>Oenothera</u> spp., <u>Baileya</u> spp. and <u>Brickellia</u> spp.	Scrub habitat species expected; 3 ravens observed	None	Potential desert tortoise habitat (0 to 50 density class)
	3.0	3.1	Broad sandy wash	DW; dominated by creosote bush and <u>Acacia greggii</u>	Scrub habitat species expected	None	Potential Gila monster habitat at Milepost 2.7, and eastward; potential desert tortoise habitat (0 to 50 density class)
	3.1	5.3	Broad alluvial fan	CBS; see species list, Table C-1	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
	5.3	8.0	Edge of dry lake	CBS and SS transition; by <u>Atriplex polycarpa</u>	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
	8.0	9.9	Edge of dry lake	CBS and SS transition; dominated by creosote bush and alkali grasses: <u>Bouteloua</u> spp. and <u>Erioneuron pulchellum</u>	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class); 3 miles northwest of potential golden eagle and prairie falcon nesting habitat on Eldorado Mountain
	9.9	12.7	Lower alluvial fan	CBS and SS transition; dominated by <u>Atriplex polycarpa</u>	Scrub habitat species expected; horn larks observed	None	Potential desert tortoise habitat (0 to 50 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
0 (cont.)	12.7	13.5	Lower alluvial fan	CBS; no associated subshrubs	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
1	0.0	2.2	Lower alluvial fan	CBS; few associated subshrubs; <u>Bouteloua</u> spp. common	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
	2.2	3.2	Lower alluvial fan	CBS	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
	3.2	5.2	Upper alluvial fan	CBS; other species include <u>Psilostrophe</u> <u>cooperi</u> , <u>Chrysothamnus</u> <u>spp.</u> , <u>Hymenoclea salsola</u> , and <u>Lycium cooperi</u>	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
	5.2	6.7	Narrow canyon floor	DW; dominated by <u>Acacia greggii</u> and <u>Opuntia acanthocarpa</u>	Scrub habitat species expected; ravens, cactus wren nest, and white-tailed antelope squirrel observed	Potential habitat for bicolored pen- stemons and Nelson's phacelia	Potential desert tortoise habitat (0 to 50 density class); potential Gila monster habitat; adjacent to potential golden eagle and prairie falcon nesting habitat
	6.7	11.2	Rugged moun- tain slopes (McCullough Mountain)	CBS; species-rich phase (see species list, Table C-2)	Scrub habitat species expected	Potential habitat for bicolored pen- stemons and Nelson's phacelia	Known bighorn sheep habitat on McCullough Mountain; adjacent to golden eagle and prairie falcon nesting habitat

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
1 (cont.)	11.2	15.0	Upper alluvial fan; sandy soils	CBS; associated species include <u>Opuntia ramossissima</u> , <u>O. echinocarpa</u> , <u>Hilaria rigida</u> , <u>Sporobolus contractus</u> , and <u>Lycium cooperi</u>	Scrub habitat species expected; house finches observed	None	Potential desert tortoise habitat (0 to 50 density class)
	15.0	16.6	Bajadas and slope bases	CBS (as above) with stand of <u>Yucca schidigera</u> between Milepost 16.0 to 16.6; surrounding hills sparsely vegetated but abundant with cacti	Scrub habitat species expected	None	Potential desert tortoise habitat (40 to 90 density class); adjacent to transient bighorn sheep habitat, Milepost 16.0 to 16.5 (Sheep Mountain)
2	0.0	4.5	Alluvial fan and slope base	CBS with stand of <u>Yucca schidigera</u> between Mileposts 0.0 and 2.0	Scrub habitat species expected; American kestrel sighted at Milepost 4.0	None	Potential desert tortoise habitat (40 to 90 density class to Milepost 2.0; 0 to 50 to Milepost 4.5); traverses edge of transient bighorn sheep habitat along Milepost 1.5 to 3.0 (Roach Mountains)
	4.5	5.5	Edge of playa	CBS and SS transition; SS contains <u>A. canescens</u> and <u>A. confertifolia</u>	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
	5.5	6.5	Edge of playa	CBS	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
3	0.0	4.0	Edge of playa	CBS; Yucca schidigera between Mileposts 0.0 and 2.0	Scrub habitat species expected; horned larks and raven observed	None	Potential desert tortoise habitat (40 to 90 density class to Milepost 2.0; 0 to 50 density class to Milepost 4.0); prairie falcon observed
	4.0	4.2	Edge of playa	CBS and SS transition	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class); prairie falcon observed nearby
4	4.2	8.4	Edge and center of playa (Roach Dry Lake)	SS and barren lands	Scrub habitat species expected in low numbers; raven observed	None	None
	0.0	1.0	Edge of playa	CBS	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
	1.0	1.9	Edge of playa	CBS and SS	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
5a	0.0	0.2	Broad sandy plain	CBS	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
5b	0.0	3.5	Broad sandy plain	CBS	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
	3.5	4.0	Edge of playa	SS	Scrub habitat species expected in low numbers	None	None

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
5b (cont.)	4.0	4.8	Dry lake (Ivanpah Dry Lake)	Barren	None	None	None
5c	0.0	1.0	Dry lake (Ivanpah Dry Lake)	Barren	None	None	None
	1.0	6.5	Edge of playa, lower bajada	CBS at lower slopes; CBS with enriched flora at high slopes (see species list, Table C-3)	Scrub habitat species expected; red-tailed hawk and cactus nests observed	None	Potential desert tortoise habitat (40 to 155 density class)
	6.5	12.3	Lower mountain slopes (Clark Mountain)	BS (see species list, Table C-4)	Scrub habitat species expected; white-tailed antelope squirrels observed	Potential habitat for Nelson's phacelia and Rusby's desert mallow	Potential Gila monster habitat up to 4,000 feet elevation; known golden eagle nesting sites 2 to 3 miles northwest on Clark Mountain; bighorn sheep habitat adjacent from Milepost 10.0 to 11.0 and traversed from Milepost 11.0 to 11.9
6	0.0	0.5	Dry lake (Roach Dry lake)	Barren	None	None	None
	0.5	0.8	Dry lake edge	SS	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (0 to 50 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
6 (cont.)	0.8	1.4	Bajada	CBS	Scrub habitat species expected; 2 American kestrels observed	None	Potential desert tortoise habitat (0 to 50 density class)
	1.4	1.8	Small ridge	CBS, sparsely vegetated with cacti (see species list, Table C-4)	Scrub habitat species expected	None	Transient bighorn sheep habitat traversed (Spring Mountains)
	1.8	8.5	Bajada	CBS, enriched with many species (see species list, Table C-3); barren hills above to north	Scrub habitat species expected	None	Potential desert tortoise habitat (40 to 155 density class); known golden eagle and prairie falcon nesting habitat 0.5 to 4 miles north in Clark Mountain Range
	8.5	18.0	Mountain slopes and canyon bottoms (Kearny Pass on Clark Mountain)	BS with JTW on higher slopes above corridor and scattered Joshua trees below; see species list, Tables C-4 and C-5; scattered <u>Pinus monophylla</u> and <u>Juniperus osteosperma</u> on higher ridges and in canyons	Scrub and woodland habitat species expected, with elements of montane forest; mule deer, white-tailed antelope squirrel, ravens, rock wren, red-shafted flicker, and black-throated sparrows observed	Potential habitat for Nelson's phacelia and Rusby's desert mallow, Milepost 8.5 to 14.0	Known bighorn sheep habitat Milepost 10.0 to 13.0 and in surrounding slopes; known golden eagle nesting sites south on Clark Mountain

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
6 (cont.)	18.0	20.0	Upper alluvial slope	CBS; few associated scrub shrub; <u>Bouteloua</u> spp. understory with scattered <u>Opuntia</u> spp; scattered Joshua trees	Scrub habitat species expected; loggerhead shrike observed	None	Potential desert tortoise habitat (0 to 50 density class)
	20.0	30.0	Broad valley and surrounding slopes	CBS with scattered Joshua trees	Scrub habitat species expected; white- crowned sparrow, cactus wren, Brewer's sparrow, and logger- head shrike observed	None	Potential desert tortoise habitat (0 to 50 density class)
	30.0	31.0	Gentle slopes	BS	Scrub habitat species expected	None	None
	31.0	44.5	Mountain slopes, canyons	CBS with scattered Joshua; see species list, Table C-6; DW at Milepost 37.6 and adjacent to corridor to Milepost 39.0; barren hills above	Scrub habitat species expected with elements of woodland habitat assemblage; cactus wren and red-tailed hawk observed	None	Potential desert tortoise habitat Milepost 39.0 to 44.5 (0 to 50 density class)
	44.5	45.5	Narrow alluvial fan	CBS with high density of <u>Hymenoclea salsola</u> and <u>Atriplex hymenelytra</u>	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost			Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End	Length				Plants	Wildlife
6 (cont.)	45.5	47.2	1.7	Narrow alluvial fan and playa edge	CBS with tamarisk trees at substation Milepost 46.3	Scrub habitat species expected; sage sparrow observed	None	Potential desert tortoise habitat (40 to 155 density class); prairie falcon observed
	47.2	48.5	1.3	Playa (Silver Dry Lake)	SS and barren areas	Scrub habitat species expected in low numbers	None	None
	48.5	63.2	14.7	Low mountain slopes and broad alluvial fans	CBS with DW at Mileposts 50.5 and 50.8 where greater diversity of shrubs present; barren hills in surrounding areas	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class); adjacent to potential golden eagle and prairie falcon nesting habitat north in Avawatz Mountains, near Milepost 56.0 to 60.0; adjacent to and north of potential golden eagle and prairie falcon nesting habitat in Soda Mountains between Milepost 54.0 and 56.0. Potential bighorn sheep movement from Avawatz Mountains to transient range in Soda Mountains between Mileposts 53.0 and 60.0
	63.2	64.0	0.8	Edge of dry playa (Red Pass Lake)	CBS with understory of <u>Hilaria rigida</u>	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class); adjacent to and north of potential golden eagle and prairie falcon nesting habitat in Soda Mountains between Mileposts 64.0 and 65.0

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
6 (cont.)	64.0	82.0	Bajadas, canyons, rugged slopes	CBS (see species list, Table C-7); most of this length in sparsely vegetated, especially on rugged slopes; sandy DW at Milepost 68.8; northern most stand of crucifixion thorn (<u>Castela emoryi</u>) 2 miles southeast of Milepost 75.0; several minor washes traversed between Mileposts 79.0 and 82.0	Scrub habitat species expected	None	Scattered potential desert tortoise habitat (0 to 50 density class); potential golden eagle and prairie falcon nesting habitat in adjacent Soda and Alvord mountains; nearby Soda Mountains are transient range for bighorn sheep
	82.0	83.3	Gentle rocky slope	CBS with numerous associated species including <u>Hilaria rigida</u> , <u>Ephedra californica</u> , <u>Psilostrophe cooperi</u> , <u>Eriogonum fasciculatum</u> and <u>E. inflatum</u>	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class); potential golden eagle and prairie falcon habitat 2 miles north on Alvord Mountain
7	0.0	1.0	Steep rugged mountain slopes	BS with scattered Joshua trees	Scrub habitat species expected	Potential habitat for Nelson's phacelia and Rusby's desert mallow	Bighorn sheep habitat near Mohawk Hill; known golden eagle nesting habitat 2 miles north on Clark Mountain

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
7 (cont.)	1.0	2.7	1.7	BS with scattered Joshua trees	Scrub habitat species expected	Potential habitat for Nelson's phacelia and Rusby's desert mallow	Adjacent to bighorn sheep habitat and near known golden eagle nest area on Clark Mountain
	2.7	5.6	2.9	CBS with scattered Joshua trees; area of <u>Yucca schidigera</u> between Mileposts 2.7 and 4.0	Scrub and woodland habitat species expected	Potential habitat for Nelson's phacelia and Rusby's desert mallow to Milepost 4.0	Potential desert tortoise habitat between Mileposts 4.0 and 5.6 (40 to 155 density class)
	5.6	5.8	0.2	CBS and SS transition with <u>Ambrosia dumosa</u> and <u>Hymenoclea salsola</u>	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (40 to 155 density class)
	5.8	7.5	1.7	SS with DW at Milepost 7.0 dominated by <u>Chilopsis linearis</u> , <u>Tamarix</u> spp., <u>Atriplex</u> spp. and <u>Gutierrezia</u> spp.	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (40 to 155 density class)
	7.5	10.4	2.9	CBS and JTW	Scrub and woodland habitat species expected	None	Potential desert tortoise habitat (40 to 155 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
7 (cont.)	10.4	13.5	Upper bajada	BS with scattered Joshua trees	Scrub habitat species expected	None	Potential desert tortoise habitat (40 to 155 density class)
8	0.0	0.6	Mountain slopes (Clark Mountain)	BS with scattered Joshua trees	Scrub and woodland habitat species expected	Potential habitat for Nelson's phacelia and Rusby's desert mallow	Bighorn sheep habitat near Mohawk Hill; known golden eagle nesting habitat 2 miles north on Clark Mountain
	0.6	2.6	Narrow canyon bottom	BS with scattered Joshua trees	Scrub and woodland habitat species expected	Potential habitat for Nelson's phacelia and Rusby's desert mallow	Adjacent to bighorn sheep habitat and near golden eagle nesting habitat on Clark Mountain
	2.6	5.9	Alluvial fan	CBS with scattered Joshua trees to Milepost 4.5 and groups of <i>Yucca schidigera</i> between Mileposts 2.6 and 3.8	Scrub and woodland habitat species expected; raven observed	Potential habitat for Nelson's phacelia and Rusby's desert mallow to Milepost 4.0	Potential desert tortoise habitat between Mileposts 4.0 to 5.9 (40 to 155 density class)
	5.9	6.3	Edge of basin	CBS and SS transition with abundant <i>Ambrosia dumosa</i> and <i>Hymenoclea salsola</i>	Scrub habitat species expected	None	Potential desert tortoise habitat (40 to 155 density class)
	6.3	9.1	Low-lying basin	SS with DW at Milepost 7.3 dominated by <i>Chilopsis linearis</i> , <i>Tamarix</i> spp., <i>Atriplex</i> spp. and <i>Gutierrezia</i> spp.	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (40 to 155 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
8 (cont.)	9.1	10.0	Lower alluvial plain	CBS and SS transition with scattered Joshua trees beginning at Milepost 9.5	Scrub habitat species expected	None	Potential deer tortoise habitat (40 to 155 density class)
	10.0	10.8	Alluvial plain	CBS with <u>Bouteloua</u> spp. understory and scattered Joshua trees	Scrub and woodland habitat species expected	None	Potential desert tortoise habitat (40 to 155 density class)
	10.8	12.9	Upper bajada	BS with scattered Joshua trees	Scrub and woodland habitat species expected	None	Potential desert tortoise habitat (40 to 155 density class)
9	0.0	2.0	Broad canyon bottom	BS with scattered Joshua trees	Scrub and woodland habitat species expected	None	None
	2.0	3.0	Steep mountain slopes	CBS; sparse vegetation with scattered Joshua trees	Scrub and woodland habitat species expected	None	None
	3.0	6.5	Steep mountain slopes	CBS, sparsely vegetated	Scrub habitat species expected in low numbers; horned lark observed	None	None

Date	Time			Location	Description	Remarks
	Start	End	Duration			
17/11	17:00	18:00	1:00	100m SW of 100m	100m SW of 100m	100m SW of 100m
18/11	18:00	19:00	1:00	100m SW of 100m	100m SW of 100m	100m SW of 100m
19/11	19:00	20:00	1:00	100m SW of 100m	100m SW of 100m	100m SW of 100m
20/11	20:00	21:00	1:00	100m SW of 100m	100m SW of 100m	100m SW of 100m
21/11	21:00	22:00	1:00	100m SW of 100m	100m SW of 100m	100m SW of 100m
22/11	22:00	23:00	1:00	100m SW of 100m	100m SW of 100m	100m SW of 100m
23/11	23:00	24:00	1:00	100m SW of 100m	100m SW of 100m	100m SW of 100m
24/11	24:00	25:00	1:00	100m SW of 100m	100m SW of 100m	100m SW of 100m
25/11	25:00	26:00	1:00	100m SW of 100m	100m SW of 100m	100m SW of 100m
26/11	26:00	27:00	1:00	100m SW of 100m	100m SW of 100m	100m SW of 100m
27/11	27:00	28:00	1:00	100m SW of 100m	100m SW of 100m	100m SW of 100m
28/11	28:00	29:00	1:00	100m SW of 100m	100m SW of 100m	100m SW of 100m
29/11	29:00	30:00	1:00	100m SW of 100m	100m SW of 100m	100m SW of 100m
30/11	30:00	31:00	1:00	100m SW of 100m	100m SW of 100m	100m SW of 100m

Continued on page 2

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
9 (cont.)	6.5	20.2	Alluvial fans, gentle slopes	CBS; sparsely vegetated and homogeneous; tamarisk trees along nearby highway	Scrub habitat species expected; ravens and Brewer's blackbirds observed	None	Potential desert tortoise habitat, 0 to 50 density class between Mileposts 6.5 to 11.5, then 40 to 155 density class to Milepost 20.2
	20.2	20.7	Low-lying basin (between Silver and Soda dry lakes)	SS	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (40 to 155 density class)
	20.7	25.0	Gentle to steep sloping hillside	CBS; barren from Mileposts 20.7 to 21.5; DW at Milepost 24.0 to 24.8 with <i>Dalea spinosa</i> , <i>Hymenoclea salsola</i> , <i>Atriplex hymenelytra</i> , and <i>Encelia</i> spp.	Scrub habitat species expected	None	Potential desert tortoise habitat (40 to 155 density class)
10	0.0	2.6	Low hills	BS with scattered Joshua trees	Scrub and woodland habitat species expected	None	None
	2.6	5.3	Upper alluvial fan	CBS with scattered Joshua trees; sparse vegetation	Scrub and woodland habitat species expected	None	None

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
10 (cont.)	5.3	12.0	Broad alluvial fan	CBS, sparsely vegetated	Scrub habitat species expected	None	None
	12.0	14.5	Broad alluvial fan	CBS	Scrub habitat species expected	None	Potential desert tortoise habitat (40 to 155 density class)
	14.5	19.9	Broad alluvial fan	CBS; tamarisk along highway to the north	Scrub habitat species expected; Brewer's blackbird; starlings, and house sparrows observed near Baker	None	Potential desert tortoise habitat (greater than 155 density class)
11	0.0	0.7	Broad plain	CBS with <u>Atriplex</u> <u>hymenelytra</u> , <u>Hymenoclea</u> <u>salsola</u> , and <u>Encelia</u> spp.	Scrub habitat species expected	None	Potential desert tortoise habitat (40 to 155 density class)
	0.7	1.7	Edge of playa	SS with <u>Atriplex</u> spp. and <u>Grayia spinosa</u>	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (0 to 50 density class)
	1.7	3.7	Dry lake bed (Soda Lake)	Barren	None	None	None
	3.7	5.5	Edge of playa	SS	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (0 to 50 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
11 (cont.)	5.5	7.9	Narrow bajada	CBS with several minor DW dominated by <u>Dalea spinosa</u>	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
	7.9	9.0	Mountain slopes	CBS, sparse vegetation on mountain slopes	Scrub habitat species expected	None	None
	9.0	18.0	Bajadas and gentle slopes; Mojave River Wash	CBS, sparse and homogeneous vegetation	Scrub habitat species expected; raven observed	None	Potential desert tortoise habitat (0 to 50 density class); traverses edge of Soda Mountains between Milepost 8.0-9.0 which is transient bighorn sheep habitat
	18.0	20.5	Mojave River Wash and Cronese Valley (south)	DW; sparse vegetation dominated by <u>Atriplex</u> spp., <u>Ambrosia dumosa</u> , <u>Croton californicus</u> ; centered to the north near Cronese Valley; <u>Prosopis glandulosa</u> and <u>Chilopsis linearis</u> stands also present	Scrub habitat species expected with DW species	<u>Chilopsis linearis</u> and <u>Prosopis glandulosa</u> sparsely distributed	Potential desert tortoise habitat (0 to 50 density class)
12	0.0	1.0	Alluvial plain	CBS with <u>Atriplex</u> <u>hymenelytra</u> , <u>Hymenoclea salsola</u> , and <u>Encelia</u> spp.	Scrub habitat species expected	None	Potential desert tortoise habitat (40 to 155 density class)

(continued) 1 of 2
 continued on page 2

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 14

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d		
	Begin	End				Plants	Wildlife	
12 (cont.)	1.0	1.3	0.3	Edge of dry lake	SS with <u>Atriplex</u> spp. and <u>Grayia spinosa</u>	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (0 to 50 density class)
	1.3	2.2	0.9	Dry lake bed (Soda Lake)	Barren	None	None	None
	2.2	2.9	0.7	Edge of dry lake	SS	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (0 to 50 density class)
	2.9	4.9	2.0	Gentle to sloping hillside	CBS with several minor DW at Milepost 3.8 to 4.8 dominated by <u>Dalea spinosa</u>	Scrub habitat species expected; raven observed	None	Potential desert tortoise habitat (0 to 50 density class)
13	0.0	10.6	10.6	Upper bajadas below barren hills	CBS; open and homogeneous with <u>Opuntia basilaris</u> and <u>Opuntia echinocarpa</u> ; several minor DW at Milepost 1.8 to 3.5 dominated by <u>Dalea spinosa</u> ; CBS very sparse at Milepost 7.3 to 9.0; sandy CBS at Milepost 9.0 to 10.5	Scrub habitat species expected; raven observed	None	Potential desert tortoise habitat (0 to 50 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
14	0.0	1.9	Edge of dry lake (East Cronese Lake)	CBS	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
	1.9	4.9	Dry lake bed and wash (Mojave River Wash and lower East Cronese Lake)	DW dominated by <u>Chilopsis linearis</u> and <u>Prosopis glandulosa</u> hummocks; other species: <u>Atriplex</u> spp., <u>Croton californica</u> , and <u>Argemone corymbosa</u>	Scrub and wash habitat species expected	<u>Prosopis glandulosa</u> and <u>Chilopsis linearis</u>	Potential desert tortoise habitat (0 to 50 density class)
15	0.0	0.5	Gentle slope	DW as above	Scrub and wash habitat species expected	<u>Prosopis glandulosa</u> and <u>Chilopsis linearis</u>	Potential desert tortoise habitat (0 to 50 density class)
	0.5	0.9	Slope base	CBS	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
	0.9	3.2	Rugged mountains (Cave Mountains)	CBS, sparse vegetation on slopes	Scrub habitat species expected in low numbers	None	Potential bighorn sheep habitat; north extension of Cady Mountain population, 2 miles north of Afton Canyon
	3.2	5.0	Lower mountain slopes	CBS on sandy soil with <u>Hilaria rigida</u> and <u>Bouteloua</u> spp.	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (0 to 50 density class)

11. 10. 11

No.	Dimensions			Material	Remarks	Remarks	Remarks	Remarks
	Length	Breadth	Height					
1	3.5	2.0	1.8	Concrete	Foundation for the main building	Foundation for the main building	Foundation for the main building	Foundation for the main building
2	3.0	1.5	1.5	Concrete	Foundation for the main building	Foundation for the main building	Foundation for the main building	Foundation for the main building
3	2.5	1.5	1.5	Concrete	Foundation for the main building	Foundation for the main building	Foundation for the main building	Foundation for the main building
4	2.0	1.5	1.5	Concrete	Foundation for the main building	Foundation for the main building	Foundation for the main building	Foundation for the main building
5	1.5	1.5	1.5	Concrete	Foundation for the main building	Foundation for the main building	Foundation for the main building	Foundation for the main building
6	1.5	1.5	1.5	Concrete	Foundation for the main building	Foundation for the main building	Foundation for the main building	Foundation for the main building
7	1.5	1.5	1.5	Concrete	Foundation for the main building	Foundation for the main building	Foundation for the main building	Foundation for the main building
8	1.5	1.5	1.5	Concrete	Foundation for the main building	Foundation for the main building	Foundation for the main building	Foundation for the main building
9	1.5	1.5	1.5	Concrete	Foundation for the main building	Foundation for the main building	Foundation for the main building	Foundation for the main building
10	1.5	1.5	1.5	Concrete	Foundation for the main building	Foundation for the main building	Foundation for the main building	Foundation for the main building

Foundation for the main building

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
16	0.0	0.3	Edge of dry lake.	CBS	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (0 to 50 density class)
	0.3	3.5	Dry lake bottom (East Cronese Lake)	DW dominated by <u>Chilopsis linearis</u> and <u>Prosopis glandulosa</u> hummocks; other species: <u>Atriplex</u> spp., <u>Croton californica</u> , and <u>Argemone corymbosa</u>	Scrub wash habitat species expected	<u>Chilopsis linearis</u> and <u>Prosopis glandulosa</u>	None
	3.5	8.6	Large canyon bottom (Cronese Valley)	CBS; sandy soils; other species: <u>Hilaria rigida</u> and <u>Bouteloua</u> spp.; sparsely vegetated hills above; sandy washout at Milepost 6.2 to 6.8	Scrub habitat species expected; white-tailed antelope squirrels observed	None	Potential desert tortoise habitat (0 to 50 density class)
17	0.0	0.5	Dry lake bottom and wash area (Mojave River Wash)	DW dominated by <u>Chilopsis linearis</u> and <u>Prosopis glandulosa</u>	Scrub and wash habitat species expected	<u>Chilopsis linearis</u> and <u>Prosopis glandulosa</u>	Potential desert tortoise habitat (0 to 50 density class)
	0.5	1.0	Lower mountain slopes	CBS	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
17 (cont.)	1.0	3.2	Rugged mountains (Cave Mountains)	CBS, sparse vegetation	Scrub habitat species expected in low numbers	None	Potential bighorn sheep habitat; north extension of Cady Mountain population; 1 to 2 miles north of Afton Canyon
	3.2	5.3	Lower mountain slopes	CBS with sandy soils; other species: <u>Hilaria rigida</u> and <u>Bouteloua</u> spp.	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class); 1 to 2 miles north of Afton Canyon
18	0.0	3.7	Broad hillsides and shallow canyons	CBS; sandy and rocky soils with variable plant cover; scattered washes with greater cover	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (0 to 50 density class); 0.5 to 1.0 mile north of Mojave River
19	0.0	8.0	Upper bajadas	CBS; sandy soil and <u>Hilaria rigida</u> with <u>Bouteloua</u> spp. between Mileposts 0.0 and 3.1; species-rich CBS between Mileposts 3.1 and 8.0 with <u>Hilaria rigida</u> , <u>Ephedra californica</u> , <u>Psilostrophe cooperi</u> , and <u>Eriogonum fasciculatum</u> ; sandy DW at Milepost 6.5	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost			Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End	Length				Plants	Wildlife
20a	0.0	4.8	4.8	Alluvial fan	CBS	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class); potential golden eagle and prairie falcon nesting habitat 3 miles north on Alvord Mountain
	4.8	8.8	4.0	Alluvial fan	SS, sparse vegetation	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (0 to 50 density class)
	8.8	11.0	2.2	Dry lake bed (Coyote Lake)	Barren	None	None	None
	11.0	11.2	0.2	Edge of dry lake	SS	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (0 to 50 density class)
	11.2	21.0	9.8	Alluvial fans, small bajadas and canyons	CBS; relatively homogeneous with <u>Bouteloua</u> spp. understorey and scattered <u>Opuntia</u> <u>ramosissima</u> ; occasional Joshua trees; surrounding hills are mostly barren	Scrub habitat species expected	Potential habitat for Jaeger's milk-vetch and Mojave fishhook cactus from Milepost 18.5 to 21.0	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (0 to 50 density class); known golden eagle and prairie falcon nesting habitat in surrounding hills (Lane and Calico mountains) from Milepost 19.0; prairie falcon observed

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
20b	0.0	8.5	Alluvial fans, small bajadas and canyons	CBS; relatively homogeneous with <u>Bouteloua</u> spp. understorey and scattered <u>Opuntia ramosissima</u> ; occasional <u>Joshua</u> trees; surrounding hills are mostly barren	Scrub habitat species expected	Potential habitat for Jaeger's milk-vetch and Mojave fishhook cactus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (0 to 50 density class from Milepost 0.0 to 4.0; 100 to 250 density class between Mileposts 4.0 and 8.5); known golden eagle and prairie falcon habitat in surrounding hills (Lane and Calico mountains) between Mileposts 0.0 and 2.0
	8.5	11.0	Mountain pass (Waterman hills)	CBS and JTW; see species list, Table C-8	Scrub and woodland habitat species expected; red-tailed hawk observed	Potential habitat for Jaeger's milk-vetch and Mojave fishhook cactus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (0 to 50 density class)
	11.0	16.0	Broad alluvial fan	CBS and DW at Mileposts 11.5 and 12.0; scattered Joshua trees	Scrub habitat species expected	Potential habitat for Mojave spine-flower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat, 100 to 250 density class between Mileposts 11.0 and 13.0; 0 to 50 density class between Mileposts 13.0 and 16.0
	16.0	16.5	Valley floor	SS; dominated by <u>Atriplex polycarpa</u> , <u>A. canescens</u> , <u>A. confertifolia</u> , and <u>Grayia sp.</u>	Scrub habitat species expected in low numbers	Potential habitat for Mojave spine-flower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (0 to 50 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
20b (cont.)	16.5	17.0	Small knoll	CBS; other species include: <u>Ambrosia dumosa</u> , <u>Hilaria rigida</u> , <u>Croton californica</u> , <u>Salsola iberica</u> , <u>Oryzopsis</u> spp., and <u>Sporobolus</u> spp.	Scrub habitat species expected	Potential habitat for Mojave spine-flower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (0 to 50 density class)
21a	17.0	17.9	Small depression	SS	Scrub habitat species expected in low numbers	Potential habitat for Mojave spine-flower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (0 to 50 density class)
	0.0	1.0	Hillside	CBS; sandy soils and abundant bare ground	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (0 to 50 density class); 2 to 3 miles south of potential golden eagle and prairie falcon nesting habitat on Alvord Mountain
	1.0	4.0	Broad alluvial fan	CBS	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
	4.0	4.2	Alluvial plain	SS; dominated by <u>Atriplex polycarpa</u> , <u>A. canescens</u> , <u>A. confertifolia</u> , <u>A. hymenelytra</u> , and <u>Grayia spinosa</u>	Scrub habitat species expected in low numbers	None	Potential desert tortoise habitat (0 to 50 density class)
	4.2	4.5	Small basin	Barren	None	None	None

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost			Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End	Length				Plants	Wildlife
21a (cont.)	4.5	6.0	1.5	Broad, flat sandy wash area	Sparsely vegetated CBS with agricultural fields and residences throughout	Scrub habitat species expected in low numbers	None	None
21b	0.0	4.0	4.0	Broad, flat sandy wash area	Sparsely vegetated CBS with agricultural fields and residences throughout	Scrub habitat species expected	None	None
	4.0	6.4	2.4	Broad, flat sandy wash area	CBS with rich understory of <u>Ambrosia dumosa</u> , <u>Grayia spinosa</u> , <u>Haplopappus</u> spp., <u>Hymenoclea salsola</u> , <u>Gutierrezia</u> spp., <u>Hilaria rigida</u> , <u>Bouteloua</u> spp., and <u>Atriplex canescens</u>	Scrub habitat species expected	Potential habitat for Barstow woolly sunflower	Potential desert tortoise habitat (0 to 50 density class)
	6.4	8.0	1.6	Edge of wash area, Mojave River	SS and barren areas near wash	Scrub habitat species expected in low numbers	Potential habitat for Barstow woolly sunflower	Potential desert tortoise habitat (0 to 50 density class); 2 miles south of potential golden eagle and prairie falcon nesting habitat in Calico Mountains at Milepost 14.0

Section 1: General Information		Section 2: Financial Data		Section 3: Operational Data	
Item	Value	Item	Value	Item	Value

1.01	1.01	2.01	2.01	3.01	3.01
1.02	1.02	2.02	2.02	3.02	3.02
1.03	1.03	2.03	2.03	3.03	3.03
1.04	1.04	2.04	2.04	3.04	3.04
1.05	1.05	2.05	2.05	3.05	3.05
1.06	1.06	2.06	2.06	3.06	3.06
1.07	1.07	2.07	2.07	3.07	3.07
1.08	1.08	2.08	2.08	3.08	3.08
1.09	1.09	2.09	2.09	3.09	3.09
1.10	1.10	2.10	2.10	3.10	3.10
1.11	1.11	2.11	2.11	3.11	3.11
1.12	1.12	2.12	2.12	3.12	3.12
1.13	1.13	2.13	2.13	3.13	3.13
1.14	1.14	2.14	2.14	3.14	3.14
1.15	1.15	2.15	2.15	3.15	3.15
1.16	1.16	2.16	2.16	3.16	3.16
1.17	1.17	2.17	2.17	3.17	3.17
1.18	1.18	2.18	2.18	3.18	3.18
1.19	1.19	2.19	2.19	3.19	3.19
1.20	1.20	2.20	2.20	3.20	3.20
1.21	1.21	2.21	2.21	3.21	3.21
1.22	1.22	2.22	2.22	3.22	3.22
1.23	1.23	2.23	2.23	3.23	3.23
1.24	1.24	2.24	2.24	3.24	3.24
1.25	1.25	2.25	2.25	3.25	3.25
1.26	1.26	2.26	2.26	3.26	3.26
1.27	1.27	2.27	2.27	3.27	3.27
1.28	1.28	2.28	2.28	3.28	3.28
1.29	1.29	2.29	2.29	3.29	3.29
1.30	1.30	2.30	2.30	3.30	3.30
1.31	1.31	2.31	2.31	3.31	3.31
1.32	1.32	2.32	2.32	3.32	3.32
1.33	1.33	2.33	2.33	3.33	3.33
1.34	1.34	2.34	2.34	3.34	3.34
1.35	1.35	2.35	2.35	3.35	3.35
1.36	1.36	2.36	2.36	3.36	3.36
1.37	1.37	2.37	2.37	3.37	3.37
1.38	1.38	2.38	2.38	3.38	3.38
1.39	1.39	2.39	2.39	3.39	3.39
1.40	1.40	2.40	2.40	3.40	3.40
1.41	1.41	2.41	2.41	3.41	3.41
1.42	1.42	2.42	2.42	3.42	3.42
1.43	1.43	2.43	2.43	3.43	3.43
1.44	1.44	2.44	2.44	3.44	3.44
1.45	1.45	2.45	2.45	3.45	3.45
1.46	1.46	2.46	2.46	3.46	3.46
1.47	1.47	2.47	2.47	3.47	3.47
1.48	1.48	2.48	2.48	3.48	3.48
1.49	1.49	2.49	2.49	3.49	3.49
1.50	1.50	2.50	2.50	3.50	3.50
1.51	1.51	2.51	2.51	3.51	3.51
1.52	1.52	2.52	2.52	3.52	3.52
1.53	1.53	2.53	2.53	3.53	3.53
1.54	1.54	2.54	2.54	3.54	3.54
1.55	1.55	2.55	2.55	3.55	3.55
1.56	1.56	2.56	2.56	3.56	3.56
1.57	1.57	2.57	2.57	3.57	3.57
1.58	1.58	2.58	2.58	3.58	3.58
1.59	1.59	2.59	2.59	3.59	3.59
1.60	1.60	2.60	2.60	3.60	3.60
1.61	1.61	2.61	2.61	3.61	3.61
1.62	1.62	2.62	2.62	3.62	3.62
1.63	1.63	2.63	2.63	3.63	3.63
1.64	1.64	2.64	2.64	3.64	3.64
1.65	1.65	2.65	2.65	3.65	3.65
1.66	1.66	2.66	2.66	3.66	3.66
1.67	1.67	2.67	2.67	3.67	3.67
1.68	1.68	2.68	2.68	3.68	3.68
1.69	1.69	2.69	2.69	3.69	3.69
1.70	1.70	2.70	2.70	3.70	3.70
1.71	1.71	2.71	2.71	3.71	3.71
1.72	1.72	2.72	2.72	3.72	3.72
1.73	1.73	2.73	2.73	3.73	3.73
1.74	1.74	2.74	2.74	3.74	3.74
1.75	1.75	2.75	2.75	3.75	3.75
1.76	1.76	2.76	2.76	3.76	3.76
1.77	1.77	2.77	2.77	3.77	3.77
1.78	1.78	2.78	2.78	3.78	3.78
1.79	1.79	2.79	2.79	3.79	3.79
1.80	1.80	2.80	2.80	3.80	3.80
1.81	1.81	2.81	2.81	3.81	3.81
1.82	1.82	2.82	2.82	3.82	3.82
1.83	1.83	2.83	2.83	3.83	3.83
1.84	1.84	2.84	2.84	3.84	3.84
1.85	1.85	2.85	2.85	3.85	3.85
1.86	1.86	2.86	2.86	3.86	3.86
1.87	1.87	2.87	2.87	3.87	3.87
1.88	1.88	2.88	2.88	3.88	3.88
1.89	1.89	2.89	2.89	3.89	3.89
1.90	1.90	2.90	2.90	3.90	3.90
1.91	1.91	2.91	2.91	3.91	3.91
1.92	1.92	2.92	2.92	3.92	3.92
1.93	1.93	2.93	2.93	3.93	3.93
1.94	1.94	2.94	2.94	3.94	3.94
1.95	1.95	2.95	2.95	3.95	3.95
1.96	1.96	2.96	2.96	3.96	3.96
1.97	1.97	2.97	2.97	3.97	3.97
1.98	1.98	2.98	2.98	3.98	3.98
1.99	1.99	2.99	2.99	3.99	3.99
2.00	2.00	3.00	3.00	4.00	4.00
2.01	2.01	3.01	3.01	4.01	4.01
2.02	2.02	3.02	3.02	4.02	4.02
2.03	2.03	3.03	3.03	4.03	4.03
2.04	2.04	3.04	3.04	4.04	4.04
2.05	2.05	3.05	3.05	4.05	4.05
2.06	2.06	3.06	3.06	4.06	4.06
2.07	2.07	3.07	3.07	4.07	4.07
2.08	2.08	3.08	3.08	4.08	4.08
2.09	2.09	3.09	3.09	4.09	4.09
2.10	2.10	3.10	3.10	4.10	4.10
2.11	2.11	3.11	3.11	4.11	4.11
2.12	2.12	3.12	3.12	4.12	4.12
2.13	2.13	3.13	3.13	4.13	4.13
2.14	2.14	3.14	3.14	4.14	4.14
2.15	2.15	3.15	3.15	4.15	4.15
2.16	2.16	3.16	3.16	4.16	4.16
2.17	2.17	3.17	3.17	4.17	4.17
2.18	2.18	3.18	3.18	4.18	4.18
2.19	2.19	3.19	3.19	4.19	4.19
2.20	2.20	3.20	3.20	4.20	4.20
2.21	2.21	3.21	3.21	4.21	4.21
2.22	2.22	3.22	3.22	4.22	4.22
2.23	2.23	3.23	3.23	4.23	4.23
2.24	2.24	3.24	3.24	4.24	4.24
2.25	2.25	3.25	3.25	4.25	4.25
2.26	2.26	3.26	3.26	4.26	4.26
2.27	2.27	3.27	3.27	4.27	4.27
2.28	2.28	3.28	3.28	4.28	4.28
2.29	2.29	3.29	3.29	4.29	4.29
2.30	2.30	3.30	3.30	4.30	4.30
2.31	2.31	3.31	3.31	4.31	4.31
2.32	2.32	3.32	3.32	4.32	4.32
2.33	2.33	3.33	3.33	4.33	4.33
2.34	2.34	3.34	3.34	4.34	4.34
2.35	2.35	3.35	3.35	4.35	4.35
2.36	2.36	3.36	3.36	4.36	4.36
2.37	2.37	3.37	3.37	4.37	4.37
2.38	2.38	3.38	3.38	4.38	4.38
2.39	2.39	3.39	3.39	4.39	4.39
2.40	2.40	3.40	3.40	4.40	4.40
2.41	2.41	3.41	3.41	4.41	4.41
2.42	2.42	3.42	3.42	4.42	4.42
2.43	2.43	3.43	3.43	4.43	4.43
2.44	2.44	3.44	3.44	4.44	4.44
2.45	2.45	3.45	3.45	4.45	4.45
2.46	2.46	3.46	3.46	4.46	4.46
2.47	2.47	3.47	3.47	4.47	4.47
2.48	2.48	3.48	3.48	4.48	4.48
2.49	2.49	3.49	3.49	4.49	4.49
2.50	2.50	3.50	3.50	4.50	4.50
2.51	2.51	3.51	3.51	4.51	4.51
2.52	2.52	3.52	3.52	4.52	4.52
2.53	2.53	3.53	3.53	4.53	4.53
2.54	2.54	3.54	3.54	4.54	4.54
2.55	2.55	3.55	3.55	4.55	4.55
2.56	2.56	3.56	3.56	4.56	4.56
2.57	2.57	3.57	3.57	4.57	4.57
2.58	2.58	3.58	3.58	4.58	4.58
2.59	2.59	3.59	3.59	4.59	4.59
2.60	2.60	3.60	3.60	4.60	4.60
2.61	2.61	3.61	3.61	4.61	4.61
2.62	2.62	3.62	3.62	4.62	4.62
2.63	2.63	3.63	3.63	4.63	4.63
2.64	2.64	3.64	3.64	4.64	4.64
2.65	2.65	3.65	3.65	4.65	4.65
2.66	2.66	3.66	3.66	4.66	4.66
2.67	2.67	3.67	3.67	4.67	4.67
2.68	2.68	3.68	3.68	4.68	4.68
2.69	2.69	3.69	3.69	4.69	4.69
2.70	2.70	3.70	3.70	4.70	4.70
2.71	2.71	3.71	3.71	4.71	4.71
2.72	2.72	3.72	3.72	4.72	4.72
2.73	2.73	3.73	3.73	4.73	4.73
2.74	2.74	3.74	3.74	4.74	4.74
2.75	2.75	3.75	3.75	4.75	4.75
2.76	2.76	3.76	3.76	4.76	4.76
2.77	2.77	3.77	3.77	4.77	4.77
2.78	2.78	3.78	3.78	4.78	4.78
2.79	2.79	3.79	3.79	4.79	4.79
2.80	2.80	3.80	3.80	4.80	4.80
2.81	2.81	3.81	3.81	4.81	4.81
2.82	2.82	3.82	3.82	4.82	4.82
2.83	2.83	3.83	3.83	4.83	4.83
2.84	2.84	3.84	3.84	4.84	4.84
2.85	2.85	3.85	3.85	4.85	4.85
2.86	2.86	3.86	3.86	4.86	4.86
2.87	2.87	3.87	3.87	4.87	4.87
2.88	2.88	3.88	3.88	4.88	4.88
2.89	2.89	3.89	3.89	4.89	4.89
2.90	2.90	3.90	3.90	4.90	4.90
2.91	2.91	3.91	3.91	4.91	4.91
2.92	2.92	3.92	3.92	4.92	4.92
2.93	2.93	3.93	3.93	4.93	4.93
2.94	2.94	3.94	3.94	4.94	4.94
2.95	2.95	3.95	3.95	4.95	4.95
2.96	2.96	3.96	3.96	4.96	4.96
2.97	2.97	3.97	3.97	4.97	4.97
2.98	2.98	3.98	3.98	4.98	4.98
2.99	2.99	3.99	3.99	4.99	4.99
3.00	3.00	4.00	4.00	5.00	5.00
3.01	3.01	4.01	4.01	5.01	5.01
3.02	3.02	4.02	4.02	5.02	5.02
3.03	3				

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
22	0.0	13.5	Sandy mesa and broad wash	CBS	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
	13.5	14.0	Flat sandy wash area	Croplands	None	None	None
	14.0	15.0	Flat sandy wash area	CBS	Scrub habitat species expected; raven observed	None	Potential desert tortoise habitat (0 to 50 density class); golden eagle observed
	15.0	16.0	Base of rocky hill (Harvard Hill)	CBS on flat area; slopes with CBS and <u>Opuntia echinocarpa</u> and <u>Opuntia engelmannii</u>	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class); golden eagle observed
16.0	20.2	20.2	Flat wash area (edge of Mohave River)	CBS with <u>Ambrosia dumosa</u> , <u>Grayia spinosa</u> , <u>Hymenoclea salsola</u> , <u>Gutierrezia</u> spp., <u>Bouteloua</u> spp., <u>Hilaria rigida</u> , and <u>Atriplex canescens</u>	Scrub habitat species expected	Potential habitat for Barstow woolly sunflower	Potential desert tortoise habitat (0 to 50 density class)
	20.2	21.2	Wash area, dunes (edge of Mohave River)	SS and DW with scattered <u>Acacia greggii</u>	Scrub habitat species expected in low numbers	Potential habitat for Barstow woolly sunflower	Potential desert tortoise habitat (0 to 50 density class); 2 miles south of potential golden eagle and prairie falcon nesting habitat in Calico Mountains at Milepost 21.2

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
23	0.0	1.5	Wash area with dunes (Mojave River crossing)	DW; scattered <u>Prosopis glandulosa</u> and <u>Acacia greggii</u>	Scrub habitat species expected in low numbers	None	2 miles south of potential golden eagle and prairie falcon nesting habitat in Calico Mountains at Milepost 0.0
	1.5	1.7	Edge of wash	SS	Scrub habitat species expected in low numbers	Potential habitat for Barstow woolly sunflower and Mojave fishhook cactus	Potential desert tortoise habitat (0 to 50 density class)
	1.7	3.5	Broad, flat valley bottom	CBS	Scrub habitat species expected	Potential habitat for Barstow woolly sunflower and Mojave fishhook cactus	Potential desert tortoise habitat (0 to 50 density class)
	3.5	4.5	Valley bottom	Croplands; tamarisks and cottonwoods planted at nearby ranch	None	None	None
	4.5	8.5	Hillside	CBS with <u>Ambrosia dumosa</u> , <u>Ephedra</u> spp., <u>Hymenoclea salsola</u> , <u>Gutierrezia</u> spp. and <u>Opuntia</u> spp.; tamarisks planted along nearby highway	Scrub habitat species expected	Potential habitat for Barstow woolly sunflower and Mojave fishhook cactus	Potential desert tortoise habitat (0 to 50 density class); 4 to 5 miles north of known golden eagle and prairie falcon nesting habitat in the Newberry Mountains

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost			Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End	Length				Plants	Wildlife
23 (cont.)	8.5	13.8	5.3	Upper bajadas and mountain slopes	CBS with numerous small DW at Mileposts 9.5, 10.0 and 11.6 dominated by <u>Acacia greggii</u> , <u>Lycium cooperi</u> , <u>Opuntia ramosissima</u> and <u>Opuntia basilaris</u> ; <u>Yucca schottii</u> - <u>gera</u> scattered in CBS between Mileposts 10.0 and 13.8	Scrub habitat species expected	Potential habitat for Mojave fishhook cactus	Potential desert tortoise habitat (0 to 50 density class); golden eagle nests previously recorded on transmission towers from Milepost 12.0
24a	0.0	5.5	5.5	Small hills, bajadas	CBS; sparse on hillsides; with <u>Bouteloua</u> spp. understory between Mileposts 2.7 and 5.5	Scrub habitat species expected	Potential habitat for Mojave fishhook cactus	Potential desert tortoise habitat (50 to 100 density class)
	5.5	8.0	2.5	Small bajadas	CBS	Scrub habitat species expected	Potential habitat for Mojave fishhook cactus	Potential desert tortoise habitat (50 to 100 density class)
	8.0	16.0	8.0	Various hills and narrow valleys	CBS; DW at Milepost 12.0 and <u>Atriplex</u> spp., <u>Gutierrezia</u> spp., <u>Hymenoclea salsola</u>	Scrub habitat species expected; loggerhead shrike, raven, and horned larks observed	Potential habitat for Mojave spineflower and Mojave fishhook cactus; nearest population of latter, 2 miles north of Milepost 14.0 at Hodge	Potential desert tortoise habitat (100 to 250 density class); prairie falcon observed at Mileposts 10.0 and 13.5

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
24a (cont.)	16.0	17.8	1.8	CBS; croplands and built-up areas within 0.5 mile	Scrub habitat species expected	Potential habitat for Mojave spineflower and Mojave fishhook cactus	Potential desert tortoise habitat (greater than 250 density class)
	17.8	18.8	1.0	SS; croplands and built-up areas within 0.5 mile	Scrub habitat species expected; mountain bluebird, white-crowned sparrow, house sparrow, western meadowlark, and greater roadrunner	None	None
	18.8	19.8	1.0	DW; sparsely vegetated with tamarisk trees and <u>Populus fremontii</u>	Scrub habitat expected; high density of birds expected; raven, northern harrier, ferruginous hawk, Brewer's blackbird, red-tailed hawk, redwing blackbird, starling, black-shouldered kite, and loggerhead shrike observed	None	Golden eagle observed
	19.8	21.0	1.2	SS and croplands; highly disturbed	Scrub habitat expected in low numbers	None	None

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
24a (cont.)	21.0	27.3	Gentle hills and small valleys; open and sandy; ends at Buckthorn Canyon	CBS with <u>Ambrosia dumosa</u> , <u>Lycium cooperi</u> , <u>Hymenoclea salsola</u> , <u>Gutierrezia</u> spp., and <u>Eriogonum inflatum</u> to Milepost 24.0; CBS with <u>Ambrosia dumosa</u> , <u>Croton californica</u> , <u>Hymenoclea salsola</u> , <u>Opuntia bigelovii</u> , and <u>Opuntia ramosissima</u>	Scrub habitat expected; ravens and kit fox scat observed	Potential habitat for desert cymopterus and Mojave spineflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (50 to 100 density class) between Mileposts 21.0 and 26.0; 100 to 250 density class between Mileposts 26.0 and 27.3; desert tortoise burrow observed at Milepost 23.0
24b	0.0	0.7	Hillside on edge of Buckthorn Canyon	SS disturbed by sheep grazing	Scrub habitat species expected	Potential habitat for Mojave spineflower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	0.7	2.0	Gentle slopes	CBS; with <u>Ambrosia dumosa</u> , <u>Croton californica</u> , <u>Hymenoclea salsola</u> , <u>Opuntia bigelovii</u> , <u>Opuntia ramosissima</u>	Scrub habitat species expected; white-tailed ground squirrel observed	Potential habitat for Mojave spineflower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	2.0	4.3	Gentle slopes	SS and CBS mixture	Scrub habitat species expected	Potential habitat for Mojave spineflower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
24b (cont.)	4.3	6.1	Hillside over wash (Fremont Wash)	CBS with infrequent and scattered Joshua trees	Scrub habitat species expected	Potential habitat for Mojave spine- flower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class); desert tortoise burrows at Mile- post 5.5
25	0.0	3.0	Broad alluvial fan	CBS with abundant <u>Hymenoclea salsola</u>	Scrub habitat species expected	Potential habitat for Mojave fish- hook cactus	Potential desert tortoise habitat (50 to 100 density class)
	3.0	7.0	Valley bottom (Stoddard Valley)	CBS; sparsely vegetated, homogeneous with <u>Bouteloua</u> spp. under- story	Scrub habitat species expected	Potential habitat for Mojave fish- hook cactus	Potential desert tortoise habitat (50 to 100 density class)
	7.0	11.7	Narrow valley	CBS with <u>Hymenoclea</u> <u>salsola</u> , <u>Ephedra cali-</u> <u>fornica</u> , and <u>Psilo-</u> <u>strophe cooperii</u> ; sur- rounding hills sparsely vegetated; <u>Yucca schidi-</u> <u>gera</u> and other species occur in upland areas from Milepost 11.0 (see species list, Table C-9)	Scrub habitat species expected	Potential habitat for Mojave fish- hook cactus	Potential desert tortoise habitat (100 to 250 density class); potential golden eagle and prairie falcon nesting habitat 2 to 3 miles south on Stoddard Ridge
26	0.0	0.6	Mountain pass	CBS; rich in species, see species list, Table C-9	Scrub habitat species expected	Potential habitat for Mojave fish- hook cactus	Potential desert tortoise habitat (0 to 50 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
27 (cont.)	0.0	5.5	Hillsides, narrow pass, and edge of valley	CBS with understory of <i>Hymenoclea salsola</i> , <i>Bouteloua</i> spp. and <i>Eriogonum inflatum</i> to Milepost 2.5; no under- story to Milepost 5.5	Scrub habitat species expected	Potential habitat for Mojave spine- flower	Potential desert tortoise habitat (greater than 250 density class)
	5.5	6.5	Upper alluvial fan	CBS with scattered <i>Yucca schidigera</i>	Scrub habitat species expected; white- tailed antelope squirrel observed	Potential habitat for Mojave fish- hook cactus and Mojave spineflower	Potential desert tortoise habitat (greater than 250 density class)
	6.5	8.0	Mountain pass	CBS with scattered <i>Yucca schidigera</i> and numerous other species (see species list, Table C-9); area of previous fire	Scrub habitat species expected; white- tailed antelope squirrel observed	Potential habitat for Mojave fish- hook cactus and Mojave spineflower	Potential desert tortoise habitat (0 to 50 density class)
	8.0	12.0	Broad alluvial	CBS	Scrub habitat species expected; American kestrel observed	Potential habitat for Mojave fish- hook cactus and Mojave spineflower; known population of the latter 1 mile north	Potential desert tortoise habitat (100 to 250 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost			Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End	Length				Plants	Wildlife
27 (cont.)	12.0	13.0	1.0	Open wash area; water present in narrow channel (Mojave River)	DW, sparsely vegetated	Scrub habitat species expected in low numbers	None	None
	13.2	13.4	0.2	Main channel Mojave River	RW on west side of river; see species list, Table C-10	Riparian woodland habitat species expected; cooper's hawk, killdeer, great blue heron, American kestrel, Say's phoebe, turkey vulture, house finch, Nuttall's woodpecker, black-billed gnatcatcher, raven, common loon, American root, common merganser, western grebe, red-billed grebe, gad-wall, American widgeon, and various ducks observed	None	None
	13.2	13.4	0.2	Bluffs overlooking river	Highly disturbed CBS; sparsely vegetated; see species list, Table C-11	Scrub habitat species expected in low numbers	None	None

Shooting 1st 5th
 secondly targeted to winners

Shooting 1st 5th		Shooting 2nd 4th		Shooting 3rd 6th		Shooting 4th 7th		Shooting 5th 8th		Shooting 6th 9th		Shooting 7th 10th		Shooting 8th 11th		Shooting 9th 12th		Shooting 10th 13th		Shooting 11th 14th		Shooting 12th 15th		Shooting 13th 16th		Shooting 14th 17th		Shooting 15th 18th		Shooting 16th 19th		Shooting 17th 20th		Shooting 18th 21st		Shooting 19th 22nd		Shooting 20th 23rd		Shooting 21st 24th		Shooting 22nd 25th		Shooting 23rd 26th		Shooting 24th 27th		Shooting 25th 28th		Shooting 26th 29th		Shooting 27th 30th		Shooting 28th 31st		Shooting 29th 32nd		Shooting 30th 33rd		Shooting 31st 34th		Shooting 32nd 35th		Shooting 33rd 36th		Shooting 34th 37th		Shooting 35th 38th		Shooting 36th 39th		Shooting 37th 40th		Shooting 38th 41st		Shooting 39th 42nd		Shooting 40th 43rd		Shooting 41st 44th		Shooting 42nd 45th		Shooting 43rd 46th		Shooting 44th 47th		Shooting 45th 48th		Shooting 46th 49th		Shooting 47th 50th		Shooting 48th 51st		Shooting 49th 52nd		Shooting 50th 53rd		Shooting 51st 54th		Shooting 52nd 55th		Shooting 53rd 56th		Shooting 54th 57th		Shooting 55th 58th		Shooting 56th 59th		Shooting 57th 60th		Shooting 58th 61st		Shooting 59th 62nd		Shooting 60th 63rd		Shooting 61st 64th		Shooting 62nd 65th		Shooting 63rd 66th		Shooting 64th 67th		Shooting 65th 68th		Shooting 66th 69th		Shooting 67th 70th		Shooting 68th 71st		Shooting 69th 72nd		Shooting 70th 73rd		Shooting 71st 74th		Shooting 72nd 75th		Shooting 73rd 76th		Shooting 74th 77th		Shooting 75th 78th		Shooting 76th 79th		Shooting 77th 80th		Shooting 78th 81st		Shooting 79th 82nd		Shooting 80th 83rd		Shooting 81st 84th		Shooting 82nd 85th		Shooting 83rd 86th		Shooting 84th 87th		Shooting 85th 88th		Shooting 86th 89th		Shooting 87th 90th		Shooting 88th 91st		Shooting 89th 92nd		Shooting 90th 93rd		Shooting 91st 94th		Shooting 92nd 95th		Shooting 93rd 96th		Shooting 94th 97th		Shooting 95th 98th		Shooting 96th 99th		Shooting 97th 100th		Shooting 98th 101st		Shooting 99th 102nd		Shooting 100th 103rd		Shooting 101st 104th		Shooting 102nd 105th		Shooting 103rd 106th		Shooting 104th 107th		Shooting 105th 108th		Shooting 106th 109th		Shooting 107th 110th		Shooting 108th 111st		Shooting 109th 112nd		Shooting 110th 113rd		Shooting 111th 114th		Shooting 112nd 115th		Shooting 113th 116th		Shooting 114th 117th		Shooting 115th 118th		Shooting 116th 119th		Shooting 117th 120th		Shooting 118th 121st		Shooting 119th 122nd		Shooting 120th 123rd		Shooting 121st 124th		Shooting 122nd 125th		Shooting 123rd 126th		Shooting 124th 127th		Shooting 125th 128th		Shooting 126th 129th		Shooting 127th 130th		Shooting 128th 131st		Shooting 129th 132nd		Shooting 130th 133rd		Shooting 131st 134th		Shooting 132nd 135th		Shooting 133rd 136th		Shooting 134th 137th		Shooting 135th 138th		Shooting 136th 139th		Shooting 137th 140th		Shooting 138th 141st		Shooting 139th 142nd		Shooting 140th 143rd		Shooting 141st 144th		Shooting 142nd 145th		Shooting 143rd 146th		Shooting 144th 147th		Shooting 145th 148th		Shooting 146th 149th		Shooting 147th 150th		Shooting 148th 151st		Shooting 149th 152nd		Shooting 150th 153rd		Shooting 151st 154th		Shooting 152nd 155th		Shooting 153rd 156th		Shooting 154th 157th		Shooting 155th 158th		Shooting 156th 159th		Shooting 157th 160th		Shooting 158th 161st		Shooting 159th 162nd		Shooting 160th 163rd		Shooting 161st 164th		Shooting 162nd 165th		Shooting 163rd 166th		Shooting 164th 167th		Shooting 165th 168th		Shooting 166th 169th		Shooting 167th 170th		Shooting 168th 171st		Shooting 169th 172nd		Shooting 170th 173rd		Shooting 171st 174th		Shooting 172nd 175th		Shooting 173rd 176th		Shooting 174th 177th		Shooting 175th 178th		Shooting 176th 179th		Shooting 177th 180th		Shooting 178th 181st		Shooting 179th 182nd		Shooting 180th 183rd		Shooting 181st 184th		Shooting 182nd 185th		Shooting 183rd 186th		Shooting 184th 187th		Shooting 185th 188th		Shooting 186th 189th		Shooting 187th 190th		Shooting 188th 191st		Shooting 189th 192nd		Shooting 190th 193rd		Shooting 191st 194th		Shooting 192nd 195th		Shooting 193rd 196th		Shooting 194th 197th		Shooting 195th 198th		Shooting 196th 199th		Shooting 197th 200th		Shooting 198th 201st		Shooting 199th 202nd		Shooting 200th 203rd		Shooting 201st 204th		Shooting 202nd 205th		Shooting 203rd 206th		Shooting 204th 207th		Shooting 205th 208th		Shooting 206th 209th		Shooting 207th 210th		Shooting 208th 211st		Shooting 209th 212nd		Shooting 210th 213rd		Shooting 211th 214th		Shooting 212nd 215th		Shooting 213th 216th		Shooting 214th 217th		Shooting 215th 218th		Shooting 216th 219th		Shooting 217th 220th		Shooting 218th 221st		Shooting 219th 222nd		Shooting 220th 223rd		Shooting 221st 224th		Shooting 222nd 225th		Shooting 223rd 226th		Shooting 224th 227th		Shooting 225th 228th		Shooting 226th 229th		Shooting 227th 230th		Shooting 228th 231st		Shooting 229th 232nd		Shooting 230th 233rd		Shooting 231st 234th		Shooting 232nd 235th		Shooting 233rd 236th		Shooting 234th 237th		Shooting 235th 238th		Shooting 236th 239th		Shooting 237th 240th		Shooting 238th 241st		Shooting 239th 242nd		Shooting 240th 243rd		Shooting 241st 244th		Shooting 242nd 245th		Shooting 243rd 246th		Shooting 244th 247th		Shooting 245th 248th		Shooting 246th 249th		Shooting 247th 250th		Shooting 248th 251st		Shooting 249th 252nd		Shooting 250th 253rd		Shooting 251st 254th		Shooting 252nd 255th		Shooting 253rd 256th		Shooting 254th 257th		Shooting 255th 258th		Shooting 256th 259th		Shooting 257th 260th		Shooting 258th 261st		Shooting 259th 262nd		Shooting 260th 263rd		Shooting 261st 264th		Shooting 262nd 265th		Shooting 263rd 266th		Shooting 264th 267th		Shooting 265th 268th		Shooting 266th 269th		Shooting 267th 270th		Shooting 268th 271st		Shooting 269th 272nd		Shooting 270th 273rd		Shooting 271st 274th		Shooting 272nd 275th		Shooting 273rd 276th		Shooting 274th 277th		Shooting 275th 278th		Shooting 276th 279th		Shooting 277th 280th		Shooting 278th 281st		Shooting 279th 282nd		Shooting 280th 283rd		Shooting 281st 284th		Shooting 282nd 285th		Shooting 283rd 286th		Shooting 284th 287th		Shooting 285th 288th		Shooting 286th 289th		Shooting 287th 290th		Shooting 288th 291st		Shooting 289th 292nd		Shooting 290th 293rd		Shooting 291st 294th		Shooting 292nd 295th		Shooting 293rd 296th		Shooting 294th 297th		Shooting 295th 298th		Shooting 296th 299th		Shooting 297th 300th		Shooting 298th 301st		Shooting 299th 302nd		Shooting 300th 303rd		Shooting 301st 304th		Shooting 302nd 305th		Shooting 303rd 306th		Shooting 304th 307th		Shooting 305th 308th		Shooting 306th 309th		Shooting 307th 310th		Shooting 308th 311st		Shooting 309th 312nd		Shooting 310th 313rd		Shooting 311th 314th		Shooting 312nd 315th		Shooting 313th 316th		Shooting 314th 317th		Shooting 315th 318th		Shooting 316th 319th		Shooting 317th 320th		Shooting 318th 321st		Shooting 319th 322nd		Shooting 320th 323rd		Shooting 321st 324th		Shooting 322nd 325th		Shooting 323rd 326th		Shooting 324th 327th		Shooting 325th 328th		Shooting 326th 329th		Shooting 327th 330th		Shooting 328th 331st		Shooting 329th 332nd		Shooting 330th 333rd		Shooting 331st 334th		Shooting 332nd 335th		Shooting 333rd 336th		Shooting 334th 337th		Shooting 335th 338th		Shooting 336th 339th		Shooting 337th 340th		Shooting 338th 341st		Shooting 339th 342nd		Shooting 340th 343rd		Shooting 341st 344th		Shooting 342nd 345th		Shooting 343rd 346th		Shooting 344th 347th		Shooting 345th 348th		Shooting 346th 349th		Shooting 347th 350th		Shooting 348th 351st		Shooting 349th 352nd		Shooting 350th 353rd		Shooting 351st 354th		Shooting 352nd 355th		Shooting 353rd 356th		Shooting 354th 357th		Shooting 355th 358th		Shooting 356th 359th		Shooting 357th 360th		Shooting 358th 361st		Shooting 359th 362nd		Shooting 360th 363rd		Shooting 361st 364th		Shooting 362nd 365th		Shooting 363rd 366th		Shooting 364th 367th		Shooting 365th 368th		Shooting 366th 369th		Shooting 367th 370th		Shooting 368th 371st		Shooting 369th 372nd		Shooting 370th 373rd		Shooting 371st 374th		Shooting 372nd 375th		Shooting 373rd 376th		Shooting 374th 377th		Shooting 375th 378th		Shooting 376th 379th		Shooting 377th 380th		Shooting 378th 381st		Shooting 379th 382nd		Shooting 380th 383rd		Shooting 381st 384th		Shooting 382nd 385th		Shooting 383rd 386th		Shooting 384th 387th		Shooting 385th 388th		Shooting 386th 389th		Shooting 387th 390th		Shooting 388th 391st		Shooting 389th 392nd		Shooting 390th 393rd		Shooting 391st 394th		Shooting 392nd 395th		Shooting 393rd 396th		Shooting 394th 397th		Shooting 395th 398th		Shooting 396th 399th		Shooting 397th 400th		Shooting 398th 401st		Shooting 399th 402nd		Shooting 400th 403rd		Shooting 401st 404th		Shooting 402nd 405th		Shooting 403rd 406th		Shooting 404th 407th		Shooting 405th 408th		Shooting 406th 409th		Shooting 407th 410th		Shooting 408th 411st		Shooting 409th 412nd		Shooting 410th 413rd		Shooting 411th 414th		Shooting 412nd 415th		Shooting 413th 416th		Shooting 414th 417th		Shooting 415th 418th		Shooting 416th 419th		Shooting 417th 420th		Shooting 418th 421st		Shooting 419th 422nd		Shooting 420th 423rd		Shooting 421st 424th		Shooting 422nd 425th		Shooting 423rd 426th		Shooting 424th 427th		Shooting 425th 428th		Shooting 426th 429th		Shooting 427th 430th		Shooting 428th 431st		Shooting 429th 432nd		Shooting 430th 433rd		Shooting 431st 434th		Shooting 432nd 435th		Shooting 433rd 436th		Shooting 434th 437th		Shooting 435th 438th		Shooting 436th 439th		Shooting 437th 440th		Shooting 438th 441st		Shooting 439th 442nd		Shooting 440th 443rd		Shooting 441st 444th		Shooting 442nd 445th		Shooting 443rd 446th		Shooting 444th 447th		Shooting 445th 448th		Shooting 446th 449th		Shooting 447th 450th		Shooting 448th 451st		Shooting 449th 452nd		Shooting 450th 453rd		Shooting 451st 454th		Shooting 452nd 455th		Shooting 453rd 456th		Shooting 454th 457th		Shooting 455th 458th		Shooting 456th 459th		Shooting 457th 460th		Shooting 458th 461st		Shooting 459th 462nd		Shooting 460th 463rd		Shooting 461st 464th		Shooting 462nd 465th		Shooting 463rd 466th		Shooting 464th 467th		Shooting 465th 468th		Shooting 466th 469th		Shooting 467th 470th		Shooting 468th 471st		Shooting 469th 472nd		Shooting 470th 473rd		Shooting 471st 474th		Shooting 472nd 475th		Shooting 473rd 476th		Shooting 474th 477th		Shooting 475th 478th		Shooting 476th 479th		Shooting 477th 480th		Shooting 478th 481st		Shooting 479th 482nd		Shooting 480th 483rd		Shooting 481st 484th		Shooting 482nd 485th		Shooting 483rd 486th		Shooting 484th 487th		Shooting 485th 488th		Shooting 486th 489th		Shooting 487th 490th		Shooting 488th 491st		Shooting 489th 492nd		Shooting 490th 493rd		Shooting 491st 494th		Shooting 492nd 495th		Shooting 493rd 496th		Shooting 494th 497th		Shooting 495th 498th		Shooting 496th 499th		Shooting 497th 500th		Shooting 498th 501st		Shooting 499th 502nd		Shooting 500th 503rd		Shooting 501st 504th		Shooting 502nd 505th		Shooting 503rd 506th		Shooting 504th 507th		Shooting 505th 508th		Shooting 506th 509th		Shooting 507th 510th		Shooting 508th 511st		Shooting 509th 512nd		Shooting 510th 513rd		Shooting 511th 514th		Shooting 512nd 515th		Shooting 513th 516th		Shooting 514th 517th		Shooting 515th 518th		Shooting 516th 519th		Shooting 517th 520th		Shooting 518th 521st		Shooting 519th 522nd		Shooting 520th 523rd		Shooting 521st 524th		Shooting 522nd 525th		Shooting 523rd 526th		Shooting 524th 527th		Shooting 525th 528th		Shooting 526th 529th		Shooting 527th 530th		Shooting 528th 531st		Shooting 529th 532nd		Shooting 530th 533rd		Shooting 531st 534th		Shooting 532nd 535th		Shooting 533rd 536th		Shooting 534th 537th		Shooting 535th 538th		Shooting 536th 539th		Shooting 537th 540th		Shooting 538th 541st		Shooting 539th 542nd		Shooting 540th 543rd		Shooting 541st 544th		Shooting 542nd 545th		Shooting 543rd 546th		Shooting 544th 547th		Shooting 545th 548th		Shooting 546th 549th		Shooting 547th 550th		Shooting 548th 551st		Shooting 549th 552nd		Shooting 550th 553rd		Shooting 551st 554th		Shooting 552nd 555th		Shooting 553rd 556th		Shooting 554th 557th		Shooting 555th 558th		Shooting 556th 559th		Shooting 557th 560th		Shooting 558th 561st		Shooting 559th 562nd		Shooting 560th 563rd		Shooting 561st 564th		Shooting 562nd 565th		Shooting 563rd 566th		Shooting 564th 567th		Shooting 565th 568th		Shooting 566th 569th		Shooting 567th 570th		Shooting 568th 571st		Shooting 569th 572nd		Shooting 570th 573rd		Shooting 571st 574th		Shooting 572nd 575th		Shooting 573rd 576th		Shooting 574th 577th		Shooting 575th 578th		Shooting 576th 579th		Shooting 577th 580th		Shooting 578th 581st		Shooting 579th 582nd		Shooting 580th 583rd		Shooting 581st 584th		Shooting 582nd 585th		Shooting 583rd 586th		Shooting 584th 587th		Shooting 585th 588th		Shooting 586th 589th		Shooting 587th 590th		Shooting 588th 591st		Shooting 589th 592nd		Shooting 590th 593rd		Shooting 591st 594th		Shooting 592nd 595th		Shooting 593rd 596th		Shooting 594th 597th		Shooting 595th 598th		Shooting 596th 599th		Shooting 597th 600th		Shooting 598th 601st		Shooting 599th 602nd		Shooting 600th 603rd		Shooting 601st 604th		Shooting 602nd 605th		Shooting 603rd 606th		Shooting 604th 607th		Shooting 605th 608th		Shooting 606th 609th		Shooting 607th 610th		Shooting 608th 611st		Shooting 609th 612nd		Shooting 610th 613rd		Shooting 611th 614th		Shooting 612nd 615th		Shooting 613th 616th		Shooting 614th 617th		Shooting 615th 618th		Shooting 616th 619th		Shooting 617th 620th		Shooting 618th 621st		Shooting 619th 622nd		Shooting 620th 623rd		Shooting 621st 624th		Shooting 622nd 625th		Shooting 623rd 626th		Shooting 624th 627th		Shooting 625th 628th		Shooting 626th 629th		Shooting 627th 630th		Shooting 628th 631st		Shooting 629th 632nd		Shooting 63	
------------------	--	------------------	--	------------------	--	------------------	--	------------------	--	------------------	--	-------------------	--	-------------------	--	-------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	--------------------	--	---------------------	--	---------------------	--	---------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	----------------------	--	-------------	--

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
28	0.0	0.6	Relatively flat terrain	Highly disturbed CBS; sparsely vegetated; see species list, Table C-11	Scrub habitat species expected in low numbers	None	None
	0.6	1.0	Relatively flat terrain	SS	Scrub habitat species expected	Potential habitat for Mojave spine-flower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	1.0	3.2	Gentle slopes and shallow depressions	CBS with small pockets of SS	Scrub habitat species expected	Potential habitat for Mojave spine-flower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
29	0.0	1.5	Sandy wash (Fremont Wash)	SS; highly disturbed by off-road vehicles	Scrub habitat species expected	Potential habitat for Mojave spine-flower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	1.5	2.1	Edge of wash	CBS	Scrub habitat species expected	Potential habitat for Mojave spine-flower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
30	0.0	1.0	Edge of wash	CBS	Scrub habitat species expected; red-tailed hawks observed	Potential habitat for Mojave spine-flower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)

5

No.	Location			Depth (m)	Date	Time	Duration (min)	Remarks
	Lat	Long	Alt					
1	0.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
2	1.2	5.1	0.2	0.2	0.2	0.2	0.2	0.2
3	0.0	1.2	1.2	1.2	1.2	1.2	1.2	1.2
4	1.0	2.5	3.3	3.3	3.3	3.3	3.3	3.3
5	0.0	1.1	0.0	0.0	0.0	0.0	0.0	0.0
6	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0

Downloaded 1-1-2019
10:00:00 AM
100.000.000.000

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
30 (cont.)	1.0	2.1	Fremont Wash area	JTW	Woodland habitat species expected	Potential habitat for Mojave spine-flower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
31	0.0	1.2	Broad, flat alluvial fan	JTW	Woodland habitat species expected	Potential habitat for Mojave spine-flower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (50 to 100 density class)
	1.2	2.0	Broad alluvial plain	CBS and SS mixture with scattered Joshua trees	Scrub habitat species expected	Potential habitat for Mojave spine-flower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (50 to 100 density class)
	2.0	3.4	Broad alluvial plain	JTW	Woodland habitat species expected	Potential habitat for Mojave spine-flower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (50 to 100 density class)
	3.4	5.7	Broad alluvial plain	CBS with scattered, infrequent Joshua trees	Scrub habitat species expected	Potential habitat for Mojave spine-flower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (50 to 100 density class)
32	0.0	1.7	Bajada	CBS with scattered <u>Yucca schidigera</u>	Scrub habitat species expected	Potential habitat for Mojave spine-flower and Mojave fishhook cactus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost			Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End	Length				Plants	Wildlife
32 (cont.)	1.7	2.6	0.9	Alluvial fan	CBS; species rich; see species list, Table C-9	Scrub habitat species expected	Potential habitat for Mojave spine- flower and Mojave fishhook cactus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	2.6	7.6	5.0	Gentle slopes and valleys	CBS	Scrub habitat species expected; raven and mountain bluebird observed	Potential habitat for Mojave spine- flower and Mojave fishhook cactus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class); prairie falcon observed
	7.6	9.8	2.2	Hills	CBS; species rich; see species list, Table C-9	Scrub habitat species expected	Potential habitat for Mojave spine- flower and Mojave fishhook cactus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (0 to 50 density class)
	9.8	14.5	4.7	Relatively steep hillsides	CBS with <u>Yucca schidi- gera</u> , <u>Hymenoclea salsola</u> , and scattered Joshua trees	Scrub habitat species expected	Potential habitat for Mojave spine- flower and Mojave fishhook cactus; nearest population of the latter at Milepost 11.0 and 0.7 mile north on Quartzite Mountain	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (0 to 50 density class); potential golden eagle and prairie falcon nesting habitat 0.7 mile north on Quartzite Mountain

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
32 (cont.)	14.5	14.6	Mojave River	RW; <u>Populus fremontii</u> and <u>tamarisk</u> with associated species (see species list, Table C-12); agriculture and built-up areas of Victorville in surrounding area	Riparian woodland habitat species expected; western bluebird, red-tailed hawk, lesser goldfinch, Say's phoebe, black phoebe, song sparrow, white-crowned sparrow, ruby-crowned knight, rock wren, red-shafted flicker, loggerhead shrike, and western meadowlark observed	None	None
	14.6	15.0	Mojave River	DW; no vegetation, little surface water	Scrub habitat species expected	None	None
	15.0	15.2	Edge of river	Cropland	None	None	None
	15.2	16.4	Hills	CBS with widely scattered Joshua trees	Scrub habitat species expected	Potential habitat for Mojave spineflower and Mojave fishhook cactus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (0 to 50 density class)
33	0.0	1.8	Bajada	CBS with scattered <u>Yucca schidigera</u>	Scrub habitat species expected	Potential habitat for Mojave spineflower and Mojave fishhook cactus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost			Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End	Length				Plants	Wildlife
33 (cont.)	1.8	2.6	0.8	Alluvial fan	CBS; species rich; see species list, Table C-9	Scrub habitat species expected	Potential habitat for Mojave spine- flower and Mojave fishhook cactus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	2.6	8.2	5.6	Gentle slopes and valleys	CBS	Scrub habitat species expected	Potential habitat for Mojave spine- flower and Mojave fishhook cactus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	8.2	10.2	2.0	Hills	CBS; species rich; see species list, Table C-9	Scrub habitat species expected	Potential habitat for Mojave spine- flower and Mojave fishhook cactus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (0 to 50 density class)
	10.2	14.6	4.4	Relatively steep hillsides	CBS with <u>Yucca schidi- gera</u> , <u>Hymenoclea salsola</u> , and scattered Joshua trees	Scrub habitat species expected	Potential habitat for Mojave spine- flower and Mojave fishhook cactus; nearest population of latter 1 mile on Quartzite Mountain	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (0 to 50 density class); potential golden eagle and prairie falcon nesting habitat 1 mile north on Quartzite Mountain

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost			Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End	Length				Plants	Wildlife
33 (cont.)	14.6	14.8	0.2	Mojave River	RW; <u>Populus fremontii</u> and <u>tamarisk</u> with associated species (see species list, Table C-12); agriculture and built-up areas of Victorville in surrounding area	Riparian woodland habitat species expected; western bluebird, red-tailed hawk, lesser goldfinch, Say's phoebe, black phoebe, song sparrow, white-crowned sparrow, ruby-crowned kinglet, rock wren, red-shafted flicker, loggerhead shrike, and western meadowlark observed	None	None
	14.8	15.7	0.9	Bluff above river	CBS with widely scattered Joshua trees; built-up areas of Victorville in surrounding area	Scrub habitat species expected in low numbers	Potential habitat for Mojave spineflower and Mojave fishhook cactus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (0 to 50 density class)
34	0.0	1.0	1.0	Low hill	CBS with widely scattered Joshua trees; adjacent to built-up areas; see species list, Table C-13	Scrub habitat species expected in low numbers	Potential habitat for Mojave spineflower and Mojave fishhook cactus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (0 to 50 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost			Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End	Length				Plants	Wildlife
34 (cont.)	1.0	1.4	0.4	Sandy wash	DW with infrequent <u>Populus fremontii</u> ; other species include <u>Distichlis spicata</u> , <u>Hymenoclea salsola</u> , <u>Chrysomaneus spp.</u> , <u>Gutierrezia spp.</u> , <u>Salsola iberica</u> , <u>Atriplex polycarpa</u> , and <u>Atriplex canescens</u>	Scrub habitat species expected; red-tailed hawk observed	None	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (0 to 50 density class)
35	0.0	4.7	4.7	Broad alluvial fan	CBS with widely scattered Joshua trees (see species list, Table C-12)	Scrub habitat species expected; raven and red-tailed hawk observed	Potential habitat for desert cymopterus and Mojave spineflower; nearest population of the latter 2.5 miles north near Adelanto	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (0 to 50 density class)
36	0.0	0.3	0.3	Bluffs above river	CBS with widely scattered Joshua trees	Scrub habitat species expected	Potential habitat for Mojave spineflower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	0.3	0.9	0.6	Relatively flat terrain	SS	Scrub habitat species expected	Potential habitat for Mojave spineflower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)

Table 1: Descriptive statistics of the data

Variable	Mean	Standard deviation	Minimum	Maximum	Skewness	Kurtosis
Age	35.2	12.5	18	65	0.1	3.2
Gender	0.5	0.5	0	1	0.0	3.0
Marital status	0.7	0.45	0	1	0.2	3.1
Education	12.5	2.5	8	16	-0.1	3.3
Income	25000	15000	5000	60000	0.5	3.5
Health	0.8	0.4	0	1	0.3	3.4
Smoking	0.3	0.45	0	1	0.1	3.2
Alcohol	0.2	0.4	0	1	0.2	3.3
Exercise	0.1	0.3	0	1	0.4	3.6
Stress	4.5	2.0	1	9	0.6	3.8
Depression	0.5	0.5	0	1	0.0	3.0
Loneliness	3.0	1.5	1	6	0.7	4.0
Social support	5.0	2.0	1	9	-0.2	3.4
Life satisfaction	6.0	1.0	4	8	-0.1	3.2

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost			Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End	Length				Plants	Wildlife
36 (cont.)	0.9	4.8	3.9	Relatively flat terrain	CBS with widely scat- tered Joshua trees	Scrub habitat species expected	Potential habitat for Mojave spine- flower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class); desert tortoise burrows observed
	4.8	5.9	1.1	Relatively flat terrain	JTW	Woodland habitat species expected	Potential habitat for Mojave spine- flower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
37	0.0	0.9	0.9	Broad alluvial plain	CBS	Scrub habitat species expected	Potential habitat for Mojave spine- flower, desert cymopterus, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (50 to 100 density class)
	0.9	1.7	0.8	Broad low-lying area	SS; see species list, Table C-14	Scrub habitat species expected	Potential habitat for Mojave spine- flower, desert cymopterus, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (50 to 100 density class);

Table 1. Summary of the data collected for the study.

Variable	Unit	Mean		SD		Range		Frequency	
		Mean	SD	Mean	SD	Min	Max	Min	Max
Age	Years	25.5	3.2	22.0	4.5	18	35	10	45
Gender	Male/Female	15/15	0	15/15	0	0	30	0	30
Education	Years	12.5	1.5	11.5	2.0	10	15	5	20
Occupation	Various	15	0	15	0	0	30	0	30
Income	\$/month	1500	200	1200	300	800	2000	500	2500
Marital Status	Married/Single	10/10	0	10/10	0	0	20	0	20
Religion	Various	15	0	15	0	0	30	0	30
Health Status	Good/Poor	10/10	0	10/10	0	0	20	0	20
Smoking Status	Smoker/Non-smoker	5/10	0	5/10	0	0	10	0	10
Alcohol Consumption	Times/week	2.5	1.5	1.5	2.0	0	5	0	10
Exercise Frequency	Times/week	1.5	1.0	1.0	1.5	0	3	0	5
Stress Level	Low/High	10/10	0	10/10	0	0	20	0	20
Depression Level	Low/High	10/10	0	10/10	0	0	20	0	20
Life Satisfaction	Low/High	10/10	0	10/10	0	0	20	0	20

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost			Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End	Length				Plants	Wildlife
37 (cont.)	1.7	3.7	2.0	Broad alluvial plain	CBS and SS mixture	Scrub habitat species expected	Potential habitat for Mojave spineflower, desert cymopterus, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (50 to 100 density class)
	3.7	6.8	3.1	Gentle alluvial fan	CBS with widely scattered Joshua trees	Scrub habitat species expected	Potential habitat for Mojave spineflower, desert cymopterus, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (50 to 100 density class) between Mileposts 3.7 and 5.0; 100 to 250 density class between Mileposts 5.0 and 6.8
	6.8	7.3	0.5	Low hills (Kramer Hills)	JTW	Woodland habitat species expected; red-tailed hawk observed	Potential habitat for Mojave spineflower, desert cymopterus, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
38	0.0	1.2	1.2	Broad alluvial fan	CBS	Scrub habitat species expected; red-tailed hawk observed	Potential habitat for Mojave spineflower, desert cymopterus, and Barstow woolly sunflower; nearest populations of latter 1.2 miles north in ACEC	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (50 to 100 density class)

Revised 1-11-2010
 Approved by the Board

Table 1: Summary of the data for the different groups

Group	N	Mean (SD)			t	p
		Age	Gender	Education		
Group 1	10	25.5 (2.1)	5 Male, 5 Female	High School	1.2	0.25
Group 2	10	26.5 (2.2)	6 Male, 4 Female	High School	1.5	0.15
Group 3	10	27.5 (2.3)	7 Male, 3 Female	High School	1.8	0.10
Group 4	10	28.5 (2.4)	8 Male, 2 Female	High School	2.1	0.05
Group 5	10	29.5 (2.5)	9 Male, 1 Female	High School	2.4	0.02
Group 6	10	30.5 (2.6)	10 Male, 0 Female	High School	2.7	0.01
Group 7	10	31.5 (2.7)	11 Male, 0 Female	High School	3.0	0.005
Group 8	10	32.5 (2.8)	12 Male, 0 Female	High School	3.3	0.001
Group 9	10	33.5 (2.9)	13 Male, 0 Female	High School	3.6	0.0005
Group 10	10	34.5 (3.0)	14 Male, 0 Female	High School	3.9	0.0001
Group 11	10	35.5 (3.1)	15 Male, 0 Female	High School	4.2	0.00005
Group 12	10	36.5 (3.2)	16 Male, 0 Female	High School	4.5	0.00001
Group 13	10	37.5 (3.3)	17 Male, 0 Female	High School	4.8	0.000005
Group 14	10	38.5 (3.4)	18 Male, 0 Female	High School	5.1	0.000001
Group 15	10	39.5 (3.5)	19 Male, 0 Female	High School	5.4	0.0000005
Group 16	10	40.5 (3.6)	20 Male, 0 Female	High School	5.7	0.0000001
Group 17	10	41.5 (3.7)	21 Male, 0 Female	High School	6.0	0.00000005
Group 18	10	42.5 (3.8)	22 Male, 0 Female	High School	6.3	0.00000001
Group 19	10	43.5 (3.9)	23 Male, 0 Female	High School	6.6	0.000000005
Group 20	10	44.5 (4.0)	24 Male, 0 Female	High School	6.9	0.000000001
Group 21	10	45.5 (4.1)	25 Male, 0 Female	High School	7.2	0.0000000005
Group 22	10	46.5 (4.2)	26 Male, 0 Female	High School	7.5	0.0000000001
Group 23	10	47.5 (4.3)	27 Male, 0 Female	High School	7.8	0.00000000005
Group 24	10	48.5 (4.4)	28 Male, 0 Female	High School	8.1	0.00000000001
Group 25	10	49.5 (4.5)	29 Male, 0 Female	High School	8.4	0.000000000005
Group 26	10	50.5 (4.6)	30 Male, 0 Female	High School	8.7	0.000000000001
Group 27	10	51.5 (4.7)	31 Male, 0 Female	High School	9.0	0.0000000000005
Group 28	10	52.5 (4.8)	32 Male, 0 Female	High School	9.3	0.0000000000001
Group 29	10	53.5 (4.9)	33 Male, 0 Female	High School	9.6	0.00000000000005
Group 30	10	54.5 (5.0)	34 Male, 0 Female	High School	9.9	0.00000000000001
Group 31	10	55.5 (5.1)	35 Male, 0 Female	High School	10.2	0.000000000000005
Group 32	10	56.5 (5.2)	36 Male, 0 Female	High School	10.5	0.000000000000001
Group 33	10	57.5 (5.3)	37 Male, 0 Female	High School	10.8	0.0000000000000005
Group 34	10	58.5 (5.4)	38 Male, 0 Female	High School	11.1	0.0000000000000001
Group 35	10	59.5 (5.5)	39 Male, 0 Female	High School	11.4	0.00000000000000005
Group 36	10	60.5 (5.6)	40 Male, 0 Female	High School	11.7	0.00000000000000001
Group 37	10	61.5 (5.7)	41 Male, 0 Female	High School	12.0	0.000000000000000005
Group 38	10	62.5 (5.8)	42 Male, 0 Female	High School	12.3	0.000000000000000001
Group 39	10	63.5 (5.9)	43 Male, 0 Female	High School	12.6	0.0000000000000000005
Group 40	10	64.5 (6.0)	44 Male, 0 Female	High School	12.9	0.0000000000000000001
Group 41	10	65.5 (6.1)	45 Male, 0 Female	High School	13.2	0.00000000000000000005
Group 42	10	66.5 (6.2)	46 Male, 0 Female	High School	13.5	0.00000000000000000001
Group 43	10	67.5 (6.3)	47 Male, 0 Female	High School	13.8	0.000000000000000000005
Group 44	10	68.5 (6.4)	48 Male, 0 Female	High School	14.1	0.000000000000000000001
Group 45	10	69.5 (6.5)	49 Male, 0 Female	High School	14.4	0.0000000000000000000005
Group 46	10	70.5 (6.6)	50 Male, 0 Female	High School	14.7	0.0000000000000000000001
Group 47	10	71.5 (6.7)	51 Male, 0 Female	High School	15.0	0.00000000000000000000005
Group 48	10	72.5 (6.8)	52 Male, 0 Female	High School	15.3	0.00000000000000000000001
Group 49	10	73.5 (6.9)	53 Male, 0 Female	High School	15.6	0.000000000000000000000005
Group 50	10	74.5 (7.0)	54 Male, 0 Female	High School	15.9	0.000000000000000000000001
Group 51	10	75.5 (7.1)	55 Male, 0 Female	High School	16.2	0.0000000000000000000000005
Group 52	10	76.5 (7.2)	56 Male, 0 Female	High School	16.5	0.0000000000000000000000001
Group 53	10	77.5 (7.3)	57 Male, 0 Female	High School	16.8	0.00000000000000000000000005
Group 54	10	78.5 (7.4)	58 Male, 0 Female	High School	17.1	0.00000000000000000000000001
Group 55	10	79.5 (7.5)	59 Male, 0 Female	High School	17.4	0.000000000000000000000000005
Group 56	10	80.5 (7.6)	60 Male, 0 Female	High School	17.7	0.0000000000000000000000000001
Group 57	10	81.5 (7.7)	61 Male, 0 Female	High School	18.0	0.00000000000000000000000000005
Group 58	10	82.5 (7.8)	62 Male, 0 Female	High School	18.3	0.00000000000000000000000000001
Group 59	10	83.5 (7.9)	63 Male, 0 Female	High School	18.6	0.000000000000000000000000000005
Group 60	10	84.5 (8.0)	64 Male, 0 Female	High School	18.9	0.000000000000000000000000000001
Group 61	10	85.5 (8.1)	65 Male, 0 Female	High School	19.2	0.0000000000000000000000000000005
Group 62	10	86.5 (8.2)	66 Male, 0 Female	High School	19.5	0.0000000000000000000000000000001
Group 63	10	87.5 (8.3)	67 Male, 0 Female	High School	19.8	0.00000000000000000000000000000005
Group 64	10	88.5 (8.4)	68 Male, 0 Female	High School	20.1	0.00000000000000000000000000000001
Group 65	10	89.5 (8.5)	69 Male, 0 Female	High School	20.4	0.000000000000000000000000000000005
Group 66	10	90.5 (8.6)	70 Male, 0 Female	High School	20.7	0.000000000000000000000000000000001
Group 67	10	91.5 (8.7)	71 Male, 0 Female	High School	21.0	0.0000000000000000000000000000000005
Group 68	10	92.5 (8.8)	72 Male, 0 Female	High School	21.3	0.0000000000000000000000000000000001
Group 69	10	93.5 (8.9)	73 Male, 0 Female	High School	21.6	0.00000000000000000000000000000000005
Group 70	10	94.5 (9.0)	74 Male, 0 Female	High School	21.9	0.00000000000000000000000000000000001
Group 71	10	95.5 (9.1)	75 Male, 0 Female	High School	22.2	0.000000000000000000000000000000000005
Group 72	10	96.5 (9.2)	76 Male, 0 Female	High School	22.5	0.000000000000000000000000000000000001
Group 73	10	97.5 (9.3)	77 Male, 0 Female	High School	22.8	0.0000000000000000000000000000000000005
Group 74	10	98.5 (9.4)	78 Male, 0 Female	High School	23.1	0.0000000000000000000000000000000000001
Group 75	10	99.5 (9.5)	79 Male, 0 Female	High School	23.4	0.00000000000000000000000000000000000005
Group 76	10	100.5 (9.6)	80 Male, 0 Female	High School	23.7	0.00000000000000000000000000000000000001
Group 77	10	101.5 (9.7)	81 Male, 0 Female	High School	24.0	0.000000000000000000000000000000000000005
Group 78	10	102.5 (9.8)	82 Male, 0 Female	High School	24.3	0.000000000000000000000000000000000000001
Group 79	10	103.5 (9.9)	83 Male, 0 Female	High School	24.6	0.0000000000000000000000000000000000000005
Group 80	10	104.5 (10.0)	84 Male, 0 Female	High School	24.9	0.0000000000000000000000000000000000000001
Group 81	10	105.5 (10.1)	85 Male, 0 Female	High School	25.2	0.00000000000000000000000000000000000000005
Group 82	10	106.5 (10.2)	86 Male, 0 Female	High School	25.5	0.00000000000000000000000000000000000000001
Group 83	10	107.5 (10.3)	87 Male, 0 Female	High School	25.8	0.000000000000000000000000000000000000000005
Group 84	10	108.5 (10.4)	88 Male, 0 Female	High School	26.1	0.000000000000000000000000000000000000000001
Group 85	10	109.5 (10.5)	89 Male, 0 Female	High School	26.4	0.0000000000000000000000000000000000000000005
Group 86	10	110.5 (10.6)	90 Male, 0 Female	High School	26.7	0.0000000000000000000000000000000000000000001
Group 87	10	111.5 (10.7)	91 Male, 0 Female	High School	27.0	0.00000000000000000000000000000000000000000005
Group 88	10	112.5 (10.8)	92 Male, 0 Female	High School	27.3	0.00000000000000000000000000000000000000000001
Group 89	10	113.5 (10.9)	93 Male, 0 Female	High School	27.6	0.000000000000000000000000000000000000000000005
Group 90	10	114.5 (11.0)	94 Male, 0 Female	High School	27.9	0.000000000000000000000000000000000000000000001
Group 91	10	115.5 (11.1)	95 Male, 0 Female	High School	28.2	0.0000000000000000000000000000000000000000000005
Group 92	10	116.5 (11.2)	96 Male, 0 Female	High School	28.5	0.0000000000000000000000000000000000000000000001
Group 93	10	117.5 (11.3)	97 Male, 0 Female	High School	28.8	0.00000000000000000000000000000000000000000000005
Group 94	10	118.5 (11.4)	98 Male, 0 Female	High School	29.1	0.00000000000000000000000000000000000000000000001
Group 95	10	119.5 (11.5)	99 Male, 0 Female	High School	29.4	0.000000000000000000000000000000000000000000000005
Group 96	10	120.5 (11.6)	100 Male, 0 Female	High School	29.7	0.000000000000000000000000000000000000000000000001
Group 97	10	121.5 (11.7)	101 Male, 0 Female	High School	30.0	0.0000000000000000000000000000000000000000000000005
Group 98	10	122.5 (11.8)	102 Male, 0 Female	High School	30.3	0.0000000000000000000000000000000000000000000000001
Group 99	10	123.5 (11.9)	103 Male, 0 Female	High School	30.6	0.00000000000000000000000000000000000000000000000005
Group 100	10	124.5 (12.0)	104 Male, 0 Female	High School	30.9	0.00000000000000000000000000000000000000000000000001

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
38 (cont.)	1.2	5.4	Broad, low- lying area	SS	Scrub habitat species expected; white-tailed antelope squirrel observed	Potential habitat for Mojave spine- flower, desert cymopterus, and Barstow woolly sunflower; nearest populations of the former two species 2 miles west of Kramer Junction; nearest population of last species 2 miles north in ACEC	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (50 to 100 density class)
	5.4	7.5	Broad, low- lying area	CBS and SS mixture	Scrub habitat species expected	Potential habitat for Mojave spine- flower, desert cymopterus, and Barstow woolly sunflower; nearest populations of the former two species 2 miles west of Kramer Junction	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (50 to 100 density class)

Table 1. Demographic characteristics of the study population

Characteristic	Number (n)	Percentage (%)	Age group (years)			
			18-24	25-34	35-44	45-54
Gender						
Male	120	60.0	30	40	30	20
Female	80	40.0	20	30	20	10
Ethnicity						
White	100	50.0	25	35	25	15
Black	60	30.0	15	20	15	10
Hispanic	40	20.0	10	15	10	5
Other	20	10.0	5	5	5	5
Marital status						
Married	100	50.0	25	35	25	15
Single	60	30.0	15	20	15	10
Divorced	40	20.0	10	15	10	5
Widowed	20	10.0	5	5	5	5
Education level						
High school or less	40	20.0	10	15	10	5
Some college	60	30.0	15	20	15	10
Bachelor's degree	80	40.0	20	30	20	10
Postgraduate	20	10.0	5	5	5	5
Annual income (\$)						
<10,000	40	20.0	10	15	10	5
10,000-20,000	60	30.0	15	20	15	10
20,000-30,000	80	40.0	20	30	20	10
>30,000	20	10.0	5	5	5	5

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
38 (cont.)	7.5	9.7	Broad alluvial fan	CBS	Scrub habitat species expected; red-tailed hawk observed	Potential habitat for Mojave spineflower, desert cymopterus, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (50 to 100 density class)
	9.7	10.3	Low hills (Kramer Hills)	JTW	Woodland habitat species expected	Potential habitat for Mojave spineflower, desert cymopterus, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (50 to 100 density class)
	0.0	0.5	Relatively flat area	JTW	Scrub habitat species expected	Potential habitat for Mojave spineflower, desert cymopterus, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
39	0.5	2.0	Relatively flat area	CBS; DW at Milepost 1.1	Scrub habitat species expected	Potential habitat for Mojave spineflower, desert cymopterus, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
39 (cont.)	2.0	3.0	Low-lying area	SS with scattered minor DWs	Scrub habitat species expected	Potential habitat for Mojave spineflower, desert cymopterus, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	3.0	4.7	Relatively flat area	CBS with DW at Milepost 3.3	Scrub habitat species expected; red-tailed hawk observed	Potential habitat for Mojave spineflower, desert cymopterus, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	4.7	6.2	Low-lying area	SS	Scrub habitat species expected; red-tailed hawk observed	Potential habitat for Mojave spineflower, desert cymopterus, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
40a	0.0	1.3	Low-lying area	SS	Scrub habitat species expected	Potential habitat for Mojave spineflower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
40a (cont.)	1.3	3.2	1.9	CBS with widely scattered Joshua trees	Scrub habitat species expected	Potential habitat for Mojave spineflower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	3.2	4.2	1.0	SS	Scrub habitat species expected	Potential habitat for Mojave spineflower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	4.2	6.2	2.0	CBS with widely scattered Joshua trees	Scrub habitat species expected	Potential habitat for Mojave spineflower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
40b	0.0	4.9	4.9	CBS with widely scattered Joshua trees	Scrub habitat species expected; raven, loggerhead shrike, red-tailed hawk, and white-tailed antelope squirrel observed	Potential habitat for Mojave spineflower and desert cymopterus	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class); desert tortoise shell observed
	0.0	7.0	7.0	SS; see species list, Table C-4; grazed area between Mileposts 3.0 and 4.5; cropland nearby between Mileposts 6.0 and 7.0; relatively	Scrub habitat species expected; red-tailed hawk and American kestrel observed	Potential habitat for Mojave spineflower, desert cymopterus, and Barstow woolly sunflower; popula-	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class); golden eagle and prairie falcon observed

Study 1: Overall Results		Study 2: Overall Results		Study 3: Overall Results		Study 4: Overall Results		Study 5: Overall Results	
Condition	Mean (SD)	Condition	Mean (SD)	Condition	Mean (SD)	Condition	Mean (SD)	Condition	Mean (SD)
Low	1.2 (0.4)	Low	1.1 (0.3)	Low	1.0 (0.2)	Low	0.9 (0.1)	Low	0.8 (0.1)
High	2.1 (0.5)	High	2.0 (0.4)	High	1.9 (0.3)	High	1.8 (0.2)	High	1.7 (0.1)
Analysis of Variance (ANOVA)									
Condition	F(1, 100) = 12.34, p < .001	Condition	F(1, 100) = 11.23, p < .001	Condition	F(1, 100) = 10.12, p < .001	Condition	F(1, 100) = 9.01, p < .001	Condition	F(1, 100) = 7.90, p < .001
Group	F(1, 100) = 3.45, p < .05	Group	F(1, 100) = 3.21, p < .05	Group	F(1, 100) = 2.98, p < .05	Group	F(1, 100) = 2.75, p < .05	Group	F(1, 100) = 2.52, p < .05
Interaction	F(1, 100) = 4.56, p < .01	Interaction	F(1, 100) = 4.32, p < .01	Interaction	F(1, 100) = 4.09, p < .01	Interaction	F(1, 100) = 3.86, p < .01	Interaction	F(1, 100) = 3.63, p < .01
Post-hoc Tests (Tukey's HSD)									
Low vs High	p < .001	Low vs High	p < .001	Low vs High	p < .001	Low vs High	p < .001	Low vs High	p < .001
Group 1 vs Group 2	p < .05	Group 1 vs Group 2	p < .05	Group 1 vs Group 2	p < .05	Group 1 vs Group 2	p < .05	Group 1 vs Group 2	p < .05
Interaction 1 vs 2	p < .01	Interaction 1 vs 2	p < .01	Interaction 1 vs 2	p < .01	Interaction 1 vs 2	p < .01	Interaction 1 vs 2	p < .01

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
41 (cont.)							
				sparse vegetation to Milepost 3.0; partial extension of Harper Dry Lake bed at Milepost 2.5; spring at Milepost 2.5 with tamarisk and <u>Populus fremontii</u>		tions of spineflower and sunflower 1.5 miles south of Milepost 7.0	
	7.0	8.5	1.5	Relatively flat area	Scrub habitat species expected	Potential habitat for desert cymopterus, Mojave spineflower, and Barstow woolly sunflower; populations of the latter species 1.5 miles south of Milepost 7.0	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class); desert tortoise burrows and golden eagle observed
	8.5	9.6	1.1	Relatively flat area	SS	Potential habitat for desert cymopterus, Mojave spineflower, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)

Table 1. Summary of the data collected from the field surveys.

Date	Time	Location	Number of individuals	Sex ratio	Age structure	Feeding behavior	Habitat preference	Notes
10/01/2010	08:00	Site A	15	1:1	10 adults, 5 juveniles	Feeding on algae	Shallow water, near rocks	First survey, good weather
10/02/2010	09:00	Site B	20	1:2	15 adults, 5 juveniles	Feeding on algae	Shallow water, near rocks	Second survey, cloudy
10/03/2010	08:30	Site A	18	1:1	12 adults, 6 juveniles	Feeding on algae	Shallow water, near rocks	Third survey, sunny
10/04/2010	09:15	Site B	22	1:1	16 adults, 6 juveniles	Feeding on algae	Shallow water, near rocks	Fourth survey, sunny
10/05/2010	08:45	Site A	16	1:1	11 adults, 5 juveniles	Feeding on algae	Shallow water, near rocks	Fifth survey, sunny
10/06/2010	09:00	Site B	21	1:2	15 adults, 6 juveniles	Feeding on algae	Shallow water, near rocks	Sixth survey, sunny
10/07/2010	08:30	Site A	19	1:1	13 adults, 6 juveniles	Feeding on algae	Shallow water, near rocks	Seventh survey, sunny
10/08/2010	09:15	Site B	23	1:1	17 adults, 6 juveniles	Feeding on algae	Shallow water, near rocks	Eighth survey, sunny
10/09/2010	08:45	Site A	17	1:1	12 adults, 5 juveniles	Feeding on algae	Shallow water, near rocks	Ninth survey, sunny
10/10/2010	09:00	Site B	24	1:2	18 adults, 6 juveniles	Feeding on algae	Shallow water, near rocks	Tenth survey, sunny

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
41 (cont.)	9.6	10.4	Sandy wash area	SS with high species diversity	Scrub habitat species expected	Potential habitat for desert cymopterus, Mojave spineflower, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	10.4	15.0	Broad, sandy alluvial fan	SS; sparsely vegetated	Scrub habitat species expected	Potential habitat for desert cymopterus, Mojave spineflower, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	15.0	15.5	Relatively flat area	CBS	Scrub habitat species expected	Potential habitat for desert cymopterus, Mojave spineflower, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (50 to 100 density class)
42	0.0	1.3	Low-lying area	SS dominated by <u>Atriplex canescens</u> , <u>Atriplex confertifolia</u> , and <u>Grayia spinosa</u>	Scrub habitat species expected	Potential habitat for desert cymopterus, Mojave spineflower, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost			Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End	Length				Plants	Wildlife
42 (cont.)	1.3	4.5	3.2	Base of slopes	CBS; sparse on hills; <i>Hilaria rigida</i> under- story; integrades with SS	Scrub habitat species expected	Potential habitat for desert cymop- terus, Mojave spineflower, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	4.5	5.0	0.5	Low-lying area	SS	Scrub habitat species expected	Potential habitat for desert cymop- terus, Mojave spineflower, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	5.0	6.2	1.2	Base of small hill	CBS	Scrub habitat species expected	Potential habitat for desert cymop- terus, Mojave spineflower, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	6.2	12.6	6.4	Broad low-lying areas	SS	Scrub habitat species expected	Potential habitat for desert cymop- terus, Mojave spineflower, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
42 (cont.)	12.6	20.9	Broad alluvial fan and flat areas	CBS with SS south of line between Mileposts 17.0 and 20.0	Scrub habitat species expected	Potential habitat for desert cymop- terus, Mojave spineflower, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	20.9	21.9	Relatively flat area	SS	Scrub habitat species expected; red-tailed hawk observed	Potential habitat for desert cymop- terus, Mojave spineflower, and Barstow woolly sunflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class); tortoise burrow observed
43	0.0	0.6	Edge of Buck- thorn Canyon	SS	Scrub habitat species expected	Potential habitat for desert cymop- terus and Mojave spineflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	0.6	1.5	Broad alluvial fan	CBS	Scrub habitat species expected	Potential habitat for desert cymop- terus and Mojave spineflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
	1.5	2.5	Broad alluvial fan	SS	Scrub habitat species expected	Potential habitat for desert cymop- terus and Mojave spineflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)

Document	Title	Author	Date	Location	Type	Status	Notes
1	Document 1-1	John Doe	1999	Room 101	Book	Available	Document 1-1
2	Document 1-2	John Doe	1999	Room 101	Book	Available	Document 1-2
3	Document 1-3	John Doe	1999	Room 101	Book	Available	Document 1-3
4	Document 1-4	John Doe	1999	Room 101	Book	Available	Document 1-4
5	Document 1-5	John Doe	1999	Room 101	Book	Available	Document 1-5
6	Document 1-6	John Doe	1999	Room 101	Book	Available	Document 1-6
7	Document 1-7	John Doe	1999	Room 101	Book	Available	Document 1-7
8	Document 1-8	John Doe	1999	Room 101	Book	Available	Document 1-8
9	Document 1-9	John Doe	1999	Room 101	Book	Available	Document 1-9
10	Document 1-10	John Doe	1999	Room 101	Book	Available	Document 1-10

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End				Plants	Wildlife
43 (cont.)	2.5	3.9	Broad alluvial fan	CBS	Scrub habitat species expected	Potential habitat for desert cymopterus and Mojave spineflower	Potential Mohave ground squirrel habitat; potential desert tortoise habitat (100 to 250 density class)
44	0.0	6.5	Broad, flat sandy area	Sparsely vegetated CBS with agricultural fields and occasional residence near Milepost 0.0 to 2.0	Scrub habitat species expected	None	None
	6.5	16.2	Alluvial fans, small bajadas and canyons	CBS with scattered <i>Opuntia ramosissima</i> ; much of surrounding hills are barren	Scrub habitat species expected	Potential habitat for Mojave fishhook cactus between Mileposts 12.5 and 16.2	Known golden eagle and prairie falcon nesting habitat in nearby Lane and Calico mountains from Milepost 12.5 to 16.2
45	0.0	4.5	Broad gentle slope	CBS	Scrub habitat species expected	None	Potential desert tortoise habitat (0 to 50 density class)
	4.5	5.0	Edge of playa	SS	Scrub habitat species expected in low numbers	None	None
	5.0	5.7	Dry lake (Ivanpah Dry Lake)	Barren	None	None	None

Table B-1 (continued)
Inventory of Biological Resources

Link No.	Milepost		Topographic Conditions	Vegetation Type ^b	Wildlife ^c	Legally Protected and Special-Interest Species ^d	
	Begin	End Length				Plants	Wildlife

Footnotes:

^aBiological resources within and adjacent to corridors are shown on Figure 4-2 (Vegetation) and 4-3 (Wildlife).

^bCBS = creosote bush scrub, dominated by Laurea tridentata and other species as described in the text or as noted; SS = saltbush scrub; BS = blackbush scrub; DW = desert wash scrub; RW = riparian woodland; JTW = Joshua tree woodland; all vegetation types described in text.

^cCommon and characteristic species of each habitat type described in text.

^dLegally protected and special-interest species described in text. Density class for desert tortoise habitat represents estimate of density (individuals per square mile) based on available literature and BLM file data.

TABLE C-1
PLANT SPECIES OBSERVED DURING FIELD SURVEYS

Species List Number	Scientific Name
---------------------	-----------------

APPENDIX C

PLANT SPECIES OBSERVED DURING FIELD SURVEYS

Grassia purpurascens
Lupinus albus
Poa annua
Polypodium vulgare
Senecio jacobaea
Ononis spinosa
Quercus agrifolia
Salix herbacea
Spartanum contractum

Asclepias tuberosa
Asperula cynosuroides
Phlox paniculata
Scilla maritima
Thalictrum flavum
Urtica dioica
Verbena officinalis
Viola tricolor
Yucca filamentosa
Zinnia mexicana
Albizia julibrissin
Chenopodium album
Cucumis melo
Echinacea purpurea
Epilobium ciliatum
Geranium robertianum
Hamamelis virginica
Hebe pinnatifida
Hydrangea arborescens
Impatiens capensis
Juniperus communis
Lonicera xylosteum
Malva sylvestris
Monarda mollis
Nicotiana glauca
Opuntia basilaris
Passiflora ligularis
Penstemon digitalis
Physalis peruviana
Rosa rugosa
Sedum spectabile
Sparganium angustifolium
Staphylea trifolia
Taxus canadensis
Thymus occidentalis
Ulmus americana
Verbena stricta
Viola blanda
Yucca filamentosa

APPENDIX C

PLANT SPECIES OBSERVED DURING FIELD SURVEYS

TABLE C-1
PLANT SPECIES OBSERVED DURING FIELD SURVEYS

<u>Species List Number</u>	<u>Scientific Name</u>
C-1	<u>Ambrosia dumosa</u> <u>Bouteloua spp.</u> <u>Chrysothamnus spp.</u> <u>Erioneuron pulchellum</u> <u>Krameria parviflora</u> <u>Larrea tridentata</u> <u>Pectis papposa</u> <u>Psilostrophe cooperi</u> <u>Oenothera deltoides</u> <u>Opuntia basilaris</u> <u>Opuntia echinocarpa</u> <u>Salsola iberica</u> <u>Sporobolus contractus</u>
C-2	<u>Acacia greggii</u> <u>Ambrosia dumosa</u> <u>Atriplex canescens</u> <u>Bouteloua spp.</u> <u>Chrysothamnus spp.</u> <u>Echinocactus polycephalus</u> <u>Echinocereus spp.</u> <u>Encelia spp.</u> <u>Ephedra californica</u> <u>Eriogonum deflexum</u> <u>E. fasciculatum</u> <u>E. inflatum</u> <u>Erioneuron pulchellum</u> <u>Ferocactus acanthoides</u> <u>Hilaria rigida</u> <u>Krameria grayi</u> <u>Larrea tridentata</u> <u>Opuntia acanthocarpa</u> <u>O. basilaris</u> <u>Physalis spp.</u> <u>Penstemon spp.</u> <u>Sphaeralcea ambigua</u> <u>Stephanomeria spp.</u> <u>Yucca schidigera</u>

Table C-1 (continued)
Plant Species Observed During Field Surveys

<u>Species List Number</u>	<u>Scientific Name</u>
C-3	<u>Acacia greggii</u> <u>Ambrosia dumosa</u> <u>Chilopsis linearis</u> <u>Chrysothamnus</u> spp. <u>Echinocactus polycephalus</u> <u>Echinocereus</u> spp. <u>Encelia virginensis</u> <u>Ephedra californica</u> <u>Eriogonum deflexum</u> <u>E. fasciculatum</u> <u>E. inflatum</u> <u>Ferocactus acanthoides</u> <u>Gutierrezia</u> spp. <u>Hymenoclea salsola</u> <u>Krameria grayi</u> <u>Larrea tridentata</u> <u>Opuntia acanthocarpa</u> <u>O. basilaris</u> <u>O. echinocarpa</u> <u>O. ramosissima</u> <u>Salizaria mexicana</u> <u>Stephanomeria</u> spp. <u>Yucca schidigera</u>
C-4	<u>Coleogyne ramosissima</u> <u>Ephedra californica</u> <u>Eriogonum</u> spp. <u>Eurotia lanata</u> <u>Grayia spinosa</u> <u>Juniperus osteosperma</u> <u>Pinus monophylla</u> <u>Thamnosma montana</u>
C-5	<u>Agave utahensis</u> ssp. <u>nevadensis</u> <u>Ephedra viridis</u> <u>Yucca baccata</u> <u>Y. brevifolia</u> <u>Y. schidigera</u>

Table C-1 (continued)
Plant Species Observed During Field Surveys

<u>Species List Number</u>	<u>Scientific Name</u>
C-6	<u>Acacia greggii</u> <u>Atriplex hymenelytra</u> <u>A. polycarpa</u> <u>Brickellia spp.</u> <u>Chilopsis linearis</u> <u>Chrysothamnus spp.</u> <u>Echinocactus polycephalus</u> <u>Encelia spp.</u> <u>Eriogonum deflexum</u> <u>E. fasciculatum</u> <u>E. inflatum</u> <u>Eurotia lanata</u> <u>Ferocactus acanthoides</u> <u>Haplopappus spp.</u> <u>Hymenoclea salsola</u> <u>Larrea tridentata</u> <u>Lycium cooperi</u> <u>Menodora spinosa</u> <u>Nicotiana trigonophylla</u> <u>Opuntia basilaris</u> <u>O. echinocarpa</u> <u>O. ramosissima</u> <u>Phoradendron spp.</u> <u>Psilostrophe cooperi</u> <u>Salizaria mexicana</u> <u>Yucca brevifolia</u> <u>Y. schidigera</u>
C-7	<u>Ambrosia dumosa</u> <u>Atriplex hymenelytra</u> <u>Eriogonum fasciculatum</u> <u>E. inflatum</u> <u>Grayia spinosa</u> <u>Haplopappus spp.</u> <u>Larrea tridentata</u> <u>Opuntia basilaris</u> <u>O. ramosissima</u> <u>Psilotrophe cooperi</u>

Table C-1 (continued)
Plant Species Observed During Field Surveys

<u>Species List Number</u>	<u>Scientific Name</u>
C-8	<u>Abronia villosa</u> <u>Ambrosia dumosa</u> <u>Astragalus</u> spp. <u>Chrysothamnus</u> spp. <u>Croton californica</u> <u>Cucurbita foetidissima</u> <u>Ephedra californica</u> <u>Eriogonum fasciculatum</u> <u>E. inflatum</u> <u>Grayia spinosa</u> <u>Hilaria spinosa</u> <u>Hymenoclea salsola</u> <u>Larrea tridentata</u> <u>Menodora spinescens</u> <u>Nicotiana trigonophylla</u> <u>Oenothera</u> spp. <u>Oryzopsis</u> spp. <u>Prunus fremontii</u> <u>Psilostrophe cooperi</u> <u>Salizaria mexicana</u> <u>Yucca brevifolia</u>
C-9	<u>Ambrosia dumosa</u> <u>Encelia</u> spp. <u>Echinocactus polycephalus</u> <u>Echinocereus</u> spp. <u>Eriogonum fasciculatum</u> <u>E. inflatum</u> <u>Eurotia lanata</u> <u>Ferocactus acanthoides</u> <u>Gutierrezia</u> spp. <u>Hymenoclea salsola</u> <u>Larrea tridentata</u> <u>Opuntia basilaris</u> <u>O. bigelovii</u> <u>O. ramosissima</u> <u>Psilostrophe cooperi</u> <u>Yucca schidigera</u>

Table C-1 (continued)
Plant Species Observed During Field Surveys

<u>Species List Number</u>	<u>Scientific Name</u>
C-10	<u>Ambrosia dumosa</u> <u>Atriplex canescens</u> <u>A. polycarpa</u> <u>Baccharis</u> spp. <u>Brickellia</u> spp. <u>Chilopsis linearis</u> <u>Chrysothamnus</u> spp. <u>Gutierrezia</u> spp. <u>Haplopappus</u> spp. <u>Populus fremontii</u> <u>Tamarix</u> spp.
C-11	<u>Ambrosia dumosa</u> <u>Ephedra californica</u> <u>Eriogonum fasciculatum</u> <u>E. inflatum</u> <u>Hymenoclea salsola</u> <u>Larrea tridentata</u> <u>Opuntia</u> spp. <u>Prunus fremontii</u>
C-12	<u>Ambrosia dumosa</u> <u>Atriplex canescens</u> <u>A. polycarpa</u> <u>Chrysothamnus</u> spp. <u>Distichlis spicata</u> <u>Ephedra californica</u> <u>Eriogonum</u> spp. <u>Gutierrezia</u> spp. <u>Grayia spinosa</u> <u>Hymenoclea salsola</u> <u>Lycium cooperi</u> <u>Populus fremontii</u> <u>Salsola iberica</u> <u>Tamarix</u> spp.

Table C-1 (continued)
Plant Species Observed During Field Surveys

<u>Species List Number</u>	<u>Scientific Name</u>
C-13	<u>Ambrosia dumosa</u> <u>Encelia spp.</u> <u>Ephedra californica</u> <u>Eriogonum inflatum</u> <u>Grayia spinosa</u> <u>Hymenoclea salsola</u> <u>Larrea tridentata</u> <u>Lycium cooperi</u> <u>Opuntia bigelovii</u>
C-14	<u>Ambrosia dumosa</u> <u>Atriplex canescens</u> <u>A. confertifolia</u> <u>A. hymenelytra</u> <u>A. polycarpa</u> <u>Eurotia lanata</u> <u>Grayia spinosa</u> <u>Lycium cooperi</u> <u>Prunus fremontii</u> <u>Salsola iberica</u>

^aSpecies list numbers keyed to observation notes in Table B-1.

